4.5 **PHYSICS (232)**

4.5.1 Physics Paper 1 (232/1)

1. 5.32 cm (1 mark)

- 2. magnitude of the force
 - The perpendicular distance between the force and the pivot. (1 mark)
- 3. Patmosphere = Pmercury + pair enclosed;

(3 marks)

4. (a) F = Ke;

$$20 = 0.5 \text{ K};$$

 $K = 40 \text{ Ncm}^{-1}$

(2 marks)

(b) $F = 40 \times 0.86 =$ = 34.4 N;

(1 mark)

- 5. Weight of object in air
 - Weight of object when fully immersed in fluids

(2 marks)

6. Upthrust = weight in air - weight of object in fluid.

(1 mark)

7. Wood is a poor conductor of heat; hence heat is used to burn paper, while most heat is conducted away by copper; hence paper takes long to burn.

(2 marks)

8. Clockwise moments = anticlockwise moments;

$$0.18x = 1(50 - x) + 0.12(100 - x)$$

$$0.18x = 50 - x + 12 - 12x$$

$$0.18x = 62 - 1.12x$$

$$7.30x = 62$$

$$x = 47.69 \text{ cm};$$

(3 marks)

9. Air is compressible; so the transmitted pressure is reduced;

(2 marks)

10. The high velocity of the gas causes a low pressure region;

Atmospheric pressure is higher;

Pressure difference draws air into the region;

(3 marks)

11. Water molecules have a high adhesion forces; With glass molecules and hence rise up the tube while mercury molecules have greater cohesion;

Forces within than adhesion with glass hence do not rise up.

(2 marks)

12. Allow for expansion;

Water expands on cooling between 4° C and 0° C;

(1 marks)

13. Diffusion of the ink molecules;

(1 mark)

SECTION B

- 14. (a) increasing the angular velocity;
 - Reducing the radius of the path;

(2 marks)

(b) (i) Tension in the string;

(1 mark)

(ii) Arrow to centre of circle;

(1 mark)

- (iii) Direction of motion of object changes and causes the velocity to change with time; (1 mark)
- (iv) $F = \frac{MV^2}{r}$; = $\frac{0.5 \# 8^{-2}}{2}$ = 16N; (3 marks)
- (c) (i) $V^2 = u^2 + 2as$; $0 = u^2 - 2 \# 10 \# 100$ $u = \sqrt{2000}$ 44.72 ms^{-1} ; (2 marks)
 - (ii) V = u + at; 0 = 44.72 - 10 # t t = 4.472Total time = 2 # 4.472= 8.94s; (2 marks)
- 15. (a) Quantity of heat required to convert 1 kg of ice at 0° C to water without change in temperature; (1 mark)
 - (b) (i) E = Pt; = 60 # 5 # 60; = 18000 J; (3 marks)
 - (ii) Mass of water = 190 130 = 60g; $ml_f = Pt$.

$$\frac{60}{1000} I_f = 60 \# 60 \# 5;$$

$$I_f = 3 \# 105 \text{ J/Kg};$$
 (4 marks)

(iii) Heat from the surrounding melts the ice; (1 mark)

16. (a)
$$F = Ma;$$

 $F=2 \# 5$
 $= 10N;$
friction force = 12 - 10

= 2N; (3 marks)

(b) (i) OA - the ball bearing decelerates; as the upthrust increases to a maximum;

(2 marks)

(4 marks)

AB - ball attains terminal velocity; when upthrust = weight; (2 marks)

(c) (i)
$$VR = 2$$
 (1 mark)

- (ii) To change direction of effort; (1 mark)
- (iii) Efficiency = $\frac{MA}{VR}$ # 100; $80 = \frac{MA}{2}$ # 100% MA = 1.6; $1.6 = \frac{L}{500}$

$$L = 500 \text{ X } 1.6$$

= 800 N; (3 marks)

- 17. (a) (i) F = mg = 10 # 10 = 100 N ; Additional pressure $= \frac{1000 \text{ M}}{1000 \text{ M}^2} = 1 \text{ Ncm}_{-2}$; new reading = 10 + 1 = 11 N;
 - (ii) Pressure has increased; because, when the volume reduces, the collisions between the gas molecules and walls of the container increases; (2 marks)
 - (b) (i) Pressure = 11 Ncm^{-2} (1 mark)
 - (ii) $\frac{P_1}{T_1} = \frac{P_2}{T_2}$ $\frac{1}{300} = \frac{11}{T_2}$; $T_2 = 300 + 11 = 330k$; $T_2 = 57^{\circ} \text{ C}$ (4 marks)

- 18. (a) (i) (I) Reading decreases on spring balance;
 - (II) Reading on weighing balance increases.
 - (ii) As the block is lowered, upthust increases; and hence it apparently weighs less;

(4 marks)

- (b) (i) Upthrust weight in air weight in water
 - = 2.7 2.46
 - = 0.24 N;

Reading in weighing balance = 2.8 + 0.24

= 3.04 N;

(2 marks)

(ii) Relative density = weight in air;

upthrust

 $=\frac{27}{0.24}$

=11.25;

Density = R.d # density of water

= 11.25 # 1000

= 11250 kgm⁻³;

(3 marks)

(c) The hydrometer sinks more;

The density of the water is reduced;


(2 marks)

4.5.2 Physics Paper 2 (232/2)

SECTION A

- 1. angle of incidence = angle of reflection = 0 (1 mark)
- 2. larger hole acts as many small holes (1 mark) many overlapping images of same object (1 mark)
- 3. Within the magnet, N and S poles of the dipoles cancel out but at the end of the poles they don't. (1 mark)
- 4. (a) 2V (1 mark)
 - (b) 1.6V (1 mark)

5.



Object at the intersection of incident ray;

(1 mark)

Incident rays;

(2 marks)

- 6. Ray totally reflected by face AC (1 mark) $i = 60^{c}$ hence $r = 60^{c}$ (1 mark)
- 7. a = 1 and b = 0 (1 mark)

x = neutron (1 mark)

8. $\frac{Ns}{Np} = \frac{Vs}{Vp}$ (1 mark)

 $\frac{Vs}{10} = \frac{Vs}{12} \qquad (1 \text{ mark})$ $Vs = 6V \qquad (1 \text{ mark})$

9. Each lamp on full voltage (1 mark)
Failure of one lamp does not affect the others (1 mark)

10. X rays ionise air molecules between plates (1 mark)

Ions move to plates of opposite sign

(1 mark)

11. Sun being hotter produces short wavelength infrared waves which penetrate glass; burning wood produces long wavelength infrared waves which do not penetrate glass.

(1 mark)

- 12. K=E-T (1 mark)
- 13. Arsenic shares 4 of its 5 electrons with germanium. (1 mark)

the extra electron is free for conduction.

(1 mark)

SECTION B

- 14. (a) $f_A = 10cm$ (1 mark)
 - (b) (i) to produce a magnified real image (1 mark)
 - (ii) to produce a magnified virtual image of the 1st image. (1 mark)
 - (c) (i) move A so that the object is slightly outside f_A (1 mark)
 - (ii) move B so that the real image is within f_B . (1 mark)
 - (d) (i) $m = \frac{24}{16}$ = $\frac{3}{2}$ (2 marks)
 - (ii) $m = \frac{28}{4}$ = 7 (2 marks)
- 15. (a) Negative charges flow from earth to cap. (1 mark)
 - Negative charge neutralizes the positive. (1 mark)
 - (b) (i) $\frac{}{c}$ $\frac{}{a}$ $\frac{}{a}$ (1 mark)
 - $=\frac{1}{3} + \frac{1}{6}$ (1 mark)

 $=\frac{1}{2}$

C = 2nF (1 mark)

(ii)
$$Q = cV$$
 (1 mark)

$$=2x4$$

$$=$$
 8*nC* (1 mark)

(iii)
$$Q = 8nC$$
 (1 mark)

(c)



- radical field;

- Correct dirrection; (2 marks)

- 16. (a) (i) Energy = QV (1 mark)
 - (ii) Power = $\frac{E}{t} = \frac{QV}{t}$ (1 mark)
 - (iii) $I = \frac{Q}{t}$ (rate of flow of charge) (1 mark)

$$P = \frac{Q}{t}.V$$

$$P = I.V$$
 (1 mark)

(b) Power =
$$VI = 20 \times 60 (1 \text{ mark})$$

$$240 \text{ x I} = 1200 \text{ W} \text{ (1 mark)}$$

$$I = \frac{1200}{240}$$

$$= 5A$$
 (1 mark)

4A15A hence fuse will blow. (1 mark)

- 17. (a) (i) Thermionically by cathode (1 mark)
 - (ii) causing fluorescence on screen (1 mark)
 - (iii) (i) control brightness of fluorescence (1 mark)
 - (ii) to focus the electron beam (1 mark)

(b) 1 wavelength =
$$2 cm$$
 (1 mark)

period =
$$2 \# 2 \# 10^{-3} s_{(1 \text{ mark})}$$

$$= 4 # 10^{-3} s$$

$$f = \frac{1}{T}$$


(1 mark)

$$= \frac{1}{4 \# 10^{-3}}$$

(1 mark)

(1 mark)

18. (a)



- curved waves - converging before focus

(1 mark)

(1 mark)


- diverging after focus.
- (b) (i) O cm trough and crest interference (2 marks)
 - (ii) +10 crest and crest interference (2 marks)
- (c) (i) Waves produced are reflected at the fixed ends. (1 mark)


 Incident and reflected waves interfer constructively at antinodes. (1 mark)
 and destructively at nodes. (1 mark)

(ii)
$$m = \frac{2}{3} # 1.5$$

= 1m (1 mark)

19. (a) (i)





- (b) coil moves to and fro (1 mark) force on coil varies direction as current varies in direction. (1 mark)
- (c) (i) dilute sulphuric acid (1 mark)
 - (ii) (I) Zinc ions go into acid leaving electrons on the plate (1 mark)
 - (II) Give up electrons to discharge hydrogen Ions. (1 mark)
 - (iii) Electrons flow from zinc plate to the copper plate. (1 mark)

4.5.3 Physics Paper 3 (232/3)

1.

PART A

(c)

Distance d (cm)	70	60	50	40
Time t for 20 oscillations(s)	24.3	25.8	26.7	27.5
t = 20 (s)	1.22	1.29	1.34	1.38
$T^4(S^4)$	2.22	2.77	3.22	3.57
$d^2(cm^2)$	4900	3600	2500	1600

(3 marks)

(1 mark)

(1 mark)

(1 mark)

Table 1 (6 marks)

(d) (i) See graph (5 marks)

Scale and axis

Plotting

Line

(1 marks)

(2 marks) (1 mark)

(ii) Slope =
$$(42^{2.50}18)^{3.50}_{2}$$
;
= $_4.2 \# 10 \bar{4} \text{ S}_4\text{cm}_2$;


(3 marks)

(iii) K =
$$\sqrt{\frac{4}{4.2 \# 10^{-4}}}$$
;

 $= 963 \text{ S}^4\text{cm}^{-2};$

(3 marks)

d (i)



1. PART B

(e)
$$l = 0.1 \text{ m}$$

 $b = 0.01 \text{ m}$ (1 mark)

$$(f) m = 0.06 \text{ kg} (1 \text{ mark})$$

(g)
$$p = \frac{0.06}{3}(0.1^2 + 0.01^2)$$

= 2.02 # 10⁻⁴ (2 marks)

(i) (I)
$$t = 75s$$
 (1 mark)

$$(II) T = 7.5s (1 mark)$$

(2 marks)

(III)
$$7.5 = 2r \sqrt{\frac{2.02 \# 10^{-4}}{G}}$$

$$G = 1.42 \# 10^{-4}$$

unit not required.

2. PARTA

(b)
$$V_0 = 3.0V$$
 (1 mark)

(d)

Voltage(V)	2.5	2.25	2.0	1.75	1.5	1.25
Time(s)	1.7	2.6	3.9	4.8	6.5	7.9

(e) (i) see graph (5 marks)

(ii) $t_{\frac{1}{2}} = 6.4 \text{ S}$ (1 mark)

(f) $R = \frac{6.4 \# 10^{6}}{0.693 \# 2200}$

$$= 4200 X \tag{1 mark}$$

PART B

(h) (i)
$$L_1 = 47.4 \text{ cm}$$
 (1 mark)

(ii)
$$W_1 = \frac{0.474 \# 0.05 \# 10}{0.35}$$

$$= 0.68 \,\mathrm{N} \tag{1 mark}$$

357

(II)
$$W_2 =$$

(i) (I)
$$L_2 = 28 \text{ cm}$$
 (1 mark)

 $= 0.4 \text{ N} \tag{1 mark}$


(j) $T1 = 26^{\circ}C$

(iii) Accept (18285-40).05 # 10 (1 mark)

(k) (i) $L_3 = 28.5 \text{ cm}$ (1 mark)

(ii) $T_2 = 83^{\circ}C$ Accept (60 - 95°C) (1 mark)

0.35



$$= \frac{0.28 - 0.27}{0.27 \# 57}$$
$$= 6.5 \# 10^{-4} \text{K}^{-1}$$

(2 marks)

(e) (i)