

|                            |   |     |     |     |     |     |
|----------------------------|---|-----|-----|-----|-----|-----|
| Volume V(cm <sup>3</sup> ) | 5 | 5.8 | 6.4 | 7.0 | 7.2 | 7.8 |
|----------------------------|---|-----|-----|-----|-----|-----|

- (i) Plot a graph of volume V against temperature.  
(ii) From the graph, determine the volume of the gas at 0°C.  
(iii) Determine the slope of the graph.  
(iv) The equation of the line obtained is of the form  $V = kT + c$ . What is the value of k and c?
6. (a) State **Charles' law**  
(b) A mass of gas occupies a volume of 150cm<sup>3</sup> at a temperature of -73°C and a pressure of 1 atmosphere. Determine the 1.5 atmospheres and the temperature 227 °C
7. In an experiment to verify Boyle's law, two quantities were advised to be kept constant  
(a). State the quantities.  
(b). the results of experiment to verify Boyle's law were recorded in the table below.

|                       |      |       |       |       |       |
|-----------------------|------|-------|-------|-------|-------|
| Pressure(atmospheres) | 1.0  | 1.2   | 1.4   | 1.6   | 1.8   |
| Volume (litres)       | 0.62 | 0.521 | 0.450 | 0.391 | 0.351 |

Plot a suitable graph to verify the law.

- (c). Determine the volume of the gas when the pressure is two atmospheres.


## SECTION 1- ANSWERS

### Measurement I

1. (a) *Density is the mass per unit volume of a substance, while relative density is the number of times a substance is denser than water- 2mks of each is defined properly*  
(b) *By law of floatation,*  
(c) *Mass of the ship = mass of water displaced*  
*Mass of water displaced = 1300000kg*  
*Volume of water displaced =  $\frac{\text{mass}}{\text{Density}}$*   

$$= \frac{1,300,000\text{kg}}{1025\text{kg/m}^3}$$

$$= 1268.3\text{m}^3$$
- (ii) *Weight of ship – weight of cargo = upthrust in fresh water*  
*13,000,000kg – W = weight of water displaced in fresh water*  
*13000000 – W = (1268.3 x 1000) x 10\*TEZ\**  
*W = 13,000,000 – 12, 683,000*  
*W = 31,7000N*  
*Cargo removed = 317tonnes*
- (c) Apparatus  
- *Measuring cylinder, water, test tube, sand and a weighing balance*



#### Procedure

1. *A measuring cylinder is half-filled with water and the level recorded*

2. Then a clean dry test tube is placed into the cylinder and some sand is added to it so that it floats upright. The new level of water is recorded.
3. the volume of water displaced is then noted, the test tube is then removed from the cylinder, it is dried and its weight determined
4. The experiment is repeated four times, adding a little more sand each time

Observation  $\frac{1}{2}$ mk for correct observation

The test-tube sinks deeper with each addition of sand. Weight of test-tube with its contents is equal to weight of water displaced.

Conclusion –  $\frac{1}{2}$ mk for correct conclusion

A floating object displaces its own weight of the fluid in which it floats. This is the fluid in which it floats. This is the law of floatation

2. a) Define relative density

The ration of density of substance to the density of water.

Or Ration of mass of a substance to the mass of equal volume of water.

3.  $A_1V_1 = A_2V_2$   
 $200 \times 2.5 \times 10^{-6} \times V = 15 \times 10^{-4} \times 5$   
 $V = \frac{15 \times 10^{-4} \times 5}{200 \times 2.5 \times 10^{-6}}$   
 $= \frac{75 \times 10^{-4}}{500 \times 10^{-6}} = \frac{7500}{500}$   
 $= 15\text{m/s}$

4. i) Relation density =  $\frac{\text{weight in air}}{\text{Up thrust in water}}$

$$= \frac{20}{20-18} = \frac{20}{2} = 10$$

$$\text{Density} = 10 \times 1000$$

$$= 10000\text{kgm}^{-3}$$


- (ii) R.d =  $\frac{\text{upthrust in liquid}}{\text{Upthrust in water}}$

$$= \frac{20-16}{20-18} = \frac{4}{2} = 2$$

$$\text{Density} = 2 \times 1000$$

$$= 2000\text{kg}^{-3}$$

5. Volume of ice =  $72-50 = 22\text{cm}^3$   
 Volume of water =  $70-50 = 20\text{cm}^3$   
 Mass of water = mass of ice = volume x density  
 $= 20\text{cm}^3 \times 1\text{gcm}^{-3}$   
 $= 20\text{g}$   
 Density of ice =  $\frac{m}{V} = \frac{20}{22} = 0.909$   
 $= 0.91\text{cm}^{-3}$



## Force

1. The mass of the balloon fabric is 2kg and the density of air is  $1.25\text{kgm}^{-3}$   
 mass of gas =  $3 \times 0.9 \text{ kg} = 2.7\text{kg}$   
 Total weight of balloon  
 $10 \times (2 + 2.7) = 47\text{N}$   
 Mass of air displaced.

$$1.25 \times 3 = 3.75$$

*Wt of air displaced*

$$1.25 \times 3 = 3.75N$$

$$\text{Tension} = U - W$$

$$= 37.5N - 22.7N = 14.8N$$

- i) Determine the tension in the string  
 ii) If the string is suddenly cut, calculate the acceleration of the balloon upwards

$$F = M\alpha$$

$$14.8 = m\alpha \quad \text{where } m = 2.27 \text{ kg}$$

$$14.8 = \frac{2.27}{2.27} \alpha$$

$$\alpha = 6.5198 \text{ m/s}^2$$

- iii) What is the maximum mass of the equipment the balloon can lift up at a constant velocity  
 maximum mass that the balloon can carry

$$\frac{14.8N}{10N/kg} = 1.48kg$$

- c) State and explain two features of a hydrometer that make it sensitive in its function.

- The stem is thin. This makes the hydrometer sensitive such that a small change in density of liquid causes a large change on the stem.
- The bulb is large to make it float.  
The bulb is heavy to make it float a upright.

2.  $\text{Static friction} = mg \sin\theta$   
 $= 5 \times 10 \sin 25^\circ$   
 $= 5 \times 10 \times 0.4226 = 21.13$

- 3.
- Increase in pressure
  - Addition of impurities

4. In water the cohesion forces between water molecules is lower than the adhesive forces between water and glass. Which in mercury the cohesion forces between mercury molecules are greater than adhesive forces between mercury and glass.

5. a) Pushing a wall/anything that does not move when force is applied

(b) (i)  $\text{work done} = \text{Area under the graph}$   
 $= (40 \times 20) + (20 \times 10) = (80 \times 40)$   
 $= 800 + 200 + 3200 = 4200J$


(ii)  $\text{work done} = mgh$   
 $= 25 \times 10 \times 12 = 3000J$   
 Energy wasted =  $(4200 - 3000)J = 1200J$

- (iii) Friction force between the surfaces. Some work is done against friction

6. For a system in equilibrium the sum of clockwise moments about a point is equal to the sum of anticlockwise moments about the same point;

7. - the latitude of the location  
 - The altitude of the location

8. i)



- ii) When side  $x$  of the film is broken, surface tension acts only on one side  $Y$  of the film;  $\sqrt{1}$  surface tension of the film tends to make the surface area to be minimum  $\sqrt{1}$  hence it pulls the string to make a smooth curve
9. Explanation- Water rises higher in a glass tube with narrow bore than the one with larger bore because more water molecules get in contact with glass molecules because of greater adhesive force between glass molecules and water molecules, then in the one with large bore.
10. (a) If no external force acts on the system of colliding bodies  
 (b) (i) – Viscosity acting  $\sqrt{1}$  upwards- each forces  
 - Upthrust acting upwards and correct directions  
 - Weight acting downwards  
 (ii) Correct curve and position  $\sqrt{1}$  above graph (a)  
 (iii) Viscosity is directly proportional to radius  $\sqrt{1}$ . Hence small ball has low friction leading to higher speed of fall and higher terminal velocity  
 (c) (i) Tension = force on car  

$$F = ma \sqrt{1}$$

$$= 1000 \times 0.5 = 500N \sqrt{1}$$


(ii) Retardation =  $\frac{F}{m} = \frac{2000}{1000} = 2m/s^2$   

$$n = \frac{36 \times 1000}{60 \times 60} = 10m/s$$

$$v^2 = u^2 + 2as$$

$$0 = (10)^2 + 2(-2)s \quad \checkmark 1$$

$$S = \frac{100}{4} = 25m \quad \checkmark 1$$



11. When the surface is dry, the frictional force between the tyres and the surface is higher  $\sqrt{1}$  than when wet, hence there is less skidding
- 12.

|   |  |
|---|--|
| Limiting friction                           | Dynamic friction                             |
| Friction between objects just before moving | Friction between surfaces in relative motion |

13. Cohesive force between mercury molecules is stronger than the adhesive force between mercury molecules and the glass side; – (correct differentiation of forces) (2mks)
14. Acceleration (a) =  $\frac{M_2g}{(M_1 + M_2)}$   

$$a = \frac{4 \times 10}{2 + 4}$$

$$= \frac{40}{6}$$

$$= 6.66$$

$$= 6.7ms^{-2}$$

## Pressure

1. Because of its low density

2. Atmospheric pressure is the pressure exerted on the surface of the surface of the earth by the weight of the air column

$$3. \quad h_w \rho_w g = h_a \rho_a g$$

$$\therefore h_w \rho_w = h_a \rho_a \quad \checkmark 1$$

$$\text{Density of alcohol} = \left( \frac{16 \cancel{\text{cm}} \times 1 \text{g/cm}^3}{20 \cancel{\text{cm}}} \right) \times 1000$$

$$= 800 \text{ kgm}^{-3} \quad \checkmark 1$$

$$4. \quad P = h \rho g$$

$$= \frac{90 \text{ m} \times 13600 \text{ kgm}^{-3} \times 10 \text{ Nkg}^{-1}}{1000} \quad \checkmark 1 \text{ mk}$$

$$= 12240 \text{ Nm}^{-2} \quad \checkmark 1 \text{ mk}$$

5. The balls move apart since the pressure on the sides is reduced by the fast moving air. High pressure between the balls pushes them outwards.

$$6. \quad \frac{(76 - 74) \times 13600 \times 10}{100} = h \times 1.25 \times 10$$

$$H = \frac{2 \times 13600}{100 \times 1.25}$$

$$= 217.6 \text{ m}$$

7. a) This is the heat energy required by a unit mass of a solid to change to liquid state at constant temperature.

b) i) The heat absorbed by the melting ice at  $0^\circ\text{C}$

$$H_1 = ML_f$$

$$= \frac{20 \text{ kg}}{1000} \times 334000 \text{ Jkg}^{-1} = 6680 \text{ J}$$

ii) The heat absorbed by the melted ice (water) to rise to temperature  $T$  (answer may be given in terms of  $T$ )

$$H_2 = \frac{20 \text{ kg}}{1000} \times 4200 \text{ Jkg}^{-1} \text{K}^{-1} (T - 0)$$

$$= 84(T - 0)$$

$$= 84T \text{ Joules}$$

iii) The heat lost by the warm water and the calorimeter (answer may be given in terms of  $T$ )

$$H_2 = \left[ \frac{200 \text{ kg}}{1000} \times 4200 \text{ Jkg}^{-1} \text{K}^{-1} + \frac{80 \times 900}{1000} \right] (T - 60)$$

$$= \left[ 840 + 72(60 - T) \right]$$

$$= 54720 - 912T$$

iv) The final temperature of the mixture

(Specific latent heat of fusion of ice =  $334000 \text{ J kg}^{-1}$ )

Specific heat capacity of water =  $4200 \text{ J kg}^{-1} \text{K}^{-1}$

Specific heat capacity of copper =  $900 \text{ J kg}^{-1} \text{K}^{-1}$ )

Heat lost = Heat gained.

$$6680 + 84T = 54720 - 912T$$

$$T \quad 912T + 84 = 54720 - 6680$$

$$T \quad \frac{996}{996} = \frac{48040}{996}$$

$$\tau = 48.233^0 \approx 48.2^0$$

8. *Pressure due to kerosene* =  $\rho K h g$   
 $= 800 \times 0.1 \times 10 = 800 p.a \sqrt{l}$

*Pressure due to water* =  $w h w g$   
 $= 1000 \times 0.2 \times 10 = 2000 p.a \sqrt{l}$

*Atmospheric pressure* =  $103,000 p.a$

*Total pressure* =  $800 + 2000 + 103000$   
 $= 105800 p.a \sqrt{l}$

9. *Pressure applied at one part in a liquid is transmitted equally to all other parts of the enclosed liquid.*

10. *Elastic PE* =  $\frac{1}{2} F e$   
 $= \frac{1}{2} \times 1.5 \times 0.01;$   
 $= 7.5 \times 10^{-3} J;$

11. *Pressure on* =  $Lfg;$   
*Solid at c* =  $(0.02 \times 1000 \times 10) + (0.04 \times 800 \times 10);$   
 $= 200 + 320$   
 $= 520 N/m^2 ;$

12. *Difference in the level of water should be 20cm*

13. *Pressure of the gas* = *Atmospheric pressure* +  $ehg;$   
 $= 1.0 \times 10^5 + \frac{20 \times 1000 \times 10}{100}$   
 $= 1.0 \times 10^5 + 2.0 \times 10^3 N/m^2$   
 $= 1.02 \times 10^5 Pa;$

14. - *Rubber is elastic; and when a nail is pushed through it stretches and grips firmly the nail without allowing air leakage;*  
*or – Valve effect pressure from inside causes tyre rubber to press firmly on the nail;*

15. (a) – *Increasing the force (weight)*  
 (b) *Slanting sides increase the area supporting the weight of the liquid, hence its effect on the bottom of the container*

$\sqrt{1}$

17. *In the narrow section of the funnel, air moves with high velocity hence followed by 10N pressure and when they emerge into the wider section, they spread, hence more min-low velocity resulting to high pressure. The high pressure below the ball lifts the ball up to the neck of the funnel.*

18. *Max pressure* =  $\frac{\text{Force}}{\text{Min Area}} \sqrt{l}$   
 $= \frac{3N}{0.1 \times 0.05} \sqrt{l}$   
 $= 600 N/m^2 \sqrt{l}$

19. (a) – *Incompressible*  
 – *Not corrosive*  
 – *Has low freezing point and high boiling point* (any one)

20.  $h_1 p_1 g = h_2 p_2 g$   
 $\frac{h_2 = h_1 p_1}{p_2}$   
 $= \frac{0.7 \times 13600 Kg/m^3}{p_2}$

$$1000\text{kgm}^{-3}$$


$$= 9.52\text{m}$$

21.  $\text{Pressure} = \frac{\text{Force}}{\text{Area}}$

$$= \frac{2500}{4 \times 0.025}$$

$$= 250,000\text{Pa}$$


22. a)
- i) Atmospheric pressure  $1.05 \times 10^5 \text{N/M}^2$
  - ii) Any water vapour available is near its condensing point. Intermolecular forces are therefore appreciable  $\checkmark$ , so it does not behave like an ideal gas
  - iii) - Fix a millimeter scale to read the length (L) of air column B  $\checkmark$  and the difference in height (h) between the levels A and C  $\checkmark$
  - Adjust the level of C by adding more mercury a little at a time and record the corresponding values of L and h each time  $\checkmark$
  - A graph of L against h represents Boyle's law  $\checkmark$



- (b) i) Increase in temperature causes gas molecules to move faster (increases in kinetic energy),  $\checkmark$  hence they generate greater/ higher impulsive force on impact  $\checkmark$
- ii) With increase in volume gas molecules are sparsely spaced  $\checkmark$  so the rate of collision is reduced/ lowered

### Current I

1. (a) natural convection – involves change in density of the fluid with temperature, whereas forced convection involves the mixing of hot and cold part of the fluid though some external stirring, like a fan or pump (2mks)



- (c) i) Copper is a good conductor of heat; hence water gets warmed faster
- ii) The glass cover does not allow the radiant heat from inside the panel to escape since this heat is lower energy than that from the sun. This heat continues being trapped inside and the temperature increases, thus boosting the heating of the water

### Particulate nature of matter

1. (a) The kinetic theory of matter states that matter is made up of tiny particles which are in a


constant random motion

(b) – Gas particles have low cohesive forces

- Gas particles have high kinetic energy

- Gas particles have low density,

(c) (i)



(ii) Experiment procedure

- A long glass tube is clamped horizontally as shown in the figure below
- A piece of cotton wool is soaked in concentrated solution of hydrochloric acid and another in concentrated ammonia solution
- Simultaneously, the soaked cotton wool pieces are inserted at the opposite ends of the horizontal glass tube and cork.
- Observations are noted

(iii) Possible observation (1mk for correct observation)

- o A white deposit of ammonium chloride forms on the walls of the glass tube in the region nearer end B

Conclusion (1mk for correct conclusion)

- Different gases have different rates of diffusion

2. Gases have weaker (small) intermolecular forces while liquids have relatively stronger (bigger) intermolecular forces ✓1mk

Or Water has stronger intermolecular forces than gases. ✓1mk

3. Brownian motion in liquids and gases

4. The rate of change of momentum of a body is directly proportional to the resultant external force producing the change and acts in the direction of the force

5. The K. E of the smoke particles reduces and hence their movements will be slower (reduces)

6. The silver coating ✓1 on the inner surfaces of the double walled glass

7. (a) The balloon expands (increases in volume) ✓1

- Evaluation reduces air pressure in the bell jar. Reduction in pressure in the jar leads to expansion of air in balloon ✓1

(b) (i) Hydrogen gas diffuses faster into the porous pot mixing with air initially in the pot, this increases pressure in the pot causing air to move out through the tube forming bubbles. ✓1

(ii) Hydrogen gas diffuses faster out of the pot. This reduces the gas pressure inside the pot hence higher atmospheric pressure on the surface of water in the beaker to push water up the glass tube. ✓1

(c) (i)  $P_1V_1 = P_2V_2$

$$P_1L_1 = P_2L_2 \quad \checkmark 1$$

$$P_1 = 750\text{mmHg}, P_2 = (750 + 50) = 800\text{mmHg}$$

$$L_1 = 40\text{mm}$$

$$750 \times 40 = 800 \times L_2$$



$$L_2 = \frac{750 \times 40}{800} \quad \checkmark 1$$

$$= 37.5 \text{ mm} \quad \checkmark 1$$

(ii) The pressure on trapped air is higher when the tube is vertical than when it is horizontal. Increase in pressure leads to reduction in volume in order to increase the number of collisions per unit time between the air particle and the walls of the glass. This increases the air pressure to balance the increased external pressure.

8. Diffusion is faster in vacuum  $\checkmark 1$  since there are no air particles to interfere with motion  $\checkmark 1$

9. Energetic molecules gain heat energy from the substance in which the liquid is in contact and escapes. This causes cooling of the latter

10. (a) (i) Air molecules/particles which were in a state of continuous random motion collided with smoke particles

(ii) They are light hence move significantly when bombarded by air molecules

(iii) There would be increased rate of movement

(b) (i) Volume of oil drops = volume of patch

$$\frac{4}{3} R^3 = \frac{d^2 t}{4} \quad t = \text{thickness}$$

$$\frac{4}{3} \times (7 \times 10^{-4})^3$$

$$= \frac{(0.75)^2 t}{4}$$

$$5.7166 \times 10^{-11} = 0.1406t$$

$$\text{Thickness, } t = \frac{5.7166 \times 10^{-11}}{0.1406}$$

$$= 4.066 \times 10^{-10} \text{ m (accept other units other than metres)}$$

(b) (ii) Assumptions- Oil drop forms a perfect sphere (1mk)

- Patch formed is a perfect circle (1mk) (any 2)

11. The particles making up gases are further apart than those in liquids  $\checkmark$

### Thermal expansion

1.  Concrete  
Steel

- The beam expands linearly

- The beam remains straight but longer than before heating

- **Both concrete and steel have same rates of expansion**

- Their value of linear expansivity is  $11 \times 10^{-6}$

2. a) - Alcohol.  $\checkmark 1$  mk

- Mercury.  $\checkmark 1$  mk

b) i) below  $-40^\circ\text{C}$  alcohol  $\checkmark \frac{1}{2}$  because it has a low freezing point of  $-115^\circ\text{C}$ .

Mercury freezes  $\checkmark \frac{1}{2}$  at  $-39^\circ\text{C}$ .

ii)  $150^\circ\text{C} \rightarrow$  mercury  $\checkmark \frac{1}{2}$  because it has a high boiling point of  $357^\circ$ ,

alcohol boils at  $78^\circ\text{C}$ .  $\checkmark \frac{1}{2}$


3. - This is the temperature of pure melting ice at standard/normal atmospheric pressure;  
(Both pure and standard pressure mentioned;

4. - Using a thin walled bulb  $\checkmark 1$

- Using a narrower capillary tube  $\checkmark 1$

5. On cooling, the brass contracts more than iron, hence become shorter than iron and forms upwards curve, making the marble to roll and settle at the centre of the curve.

6.



## Measurement II

- Exact diameter reading – 0.11
  - Find the density of the ball bearing giving your answer correct to three significant figures

$$V = \frac{4}{3} \pi \left(\frac{D}{2}\right)^3$$

$$P = m/v$$

- Determine the extension of the system

6N purchase 2cm extension

$$50N = \frac{2 \times 50}{6} = 16.667 \text{ cm}$$

|   |                 |
|---|-----------------|
| Total extension = 16.667X2 + $\frac{16.667}{Z}$ | 33.33           |
|   | 8.33            |
|   | <u>41.66 cm</u> |

- Water in a dam falls through a height 24.5m. If we assume that there are no energy losses, calculate the new temperature of the water as it strikes the lower end, given that its initial temperature at the top of the dam is 18.9°C.

Take specific heat capacity of water = 4200Jkg<sup>-1</sup> K<sup>-1</sup>

$$mgh = mC\Delta\theta$$

$$24.5 \times 10 = 4200 \text{ J/kg K} \times (\theta - 18.9)$$

$$\frac{24.5}{420} = \theta - 18.9$$

$$420$$

$$\theta = 18.9583^\circ \text{C or K}$$

- The powder around the hot needle moves away.

Reason

- The high temperature of the needle lowers the surface tension of the water around it.
- High surface tension on the sided pits the powder away

- (a) (i) Volume of one drop =  $\frac{26.00 - 25.2}{50}$

$$= \frac{0.8}{50} = 0.016 \text{ cm}^3$$

$$(ii) \text{ Thickness of oil molecule} = \frac{\text{vol. of drop}}{\text{Area of drop}}$$

$$= \frac{0.016}{3.142 \times 3.5 \times 3.5} = 4.15 \times 10^{-4} \text{ cm}$$

$$3.142 \times 3.5 \times 3.5 = 4.15 \times 10^{-4} \text{ cm}$$

(iii) – The patch is even

- Oil drop forms a monolayer

(iv) – To show the circular patch formed by the oil drop

$$(b) \text{ Vol. of molecule} = 1.5 \times 10^{-9} \times 0.6 \times 10^{-9} \times 0.6 \times 10^{-9}$$

$$= 0.54 \times 10^{-27} = 5.4 \times 10^{-28} \text{ m}^3$$


$$1 \text{ litre} = 1000 \text{ cm}^3$$

$$= 1.0 \times 10^{-3} \text{ m}^3$$

$$\text{No. of molecules in 1 litre} = \frac{1.0 \times 10^{-3}}{5.4 \times 10^{-28}}$$


$$= 0.18515 \times 10^{25}$$

$$= 1.8519 \times 10^{24} \text{ molecules}$$

5. *-ve error is added*
- 
- $5.99$   
 $\underline{0.06}$   
 $6.05\text{mm}$

6. *Actual reading = 3.21 + 0.06; = 3.27cm;*  
of 0.5mm

7. (i) *Reading = actual reading = zero error*  
 $= 2.32\text{mm} + -0.01$   
 $= 2.31\text{mm} \quad (1\text{mk})$



9. *Reading, mass = 2.75Kg √1*  
*Density = mass/volume = 2.75Kg/3x10<sup>-4</sup>m<sup>3</sup> √1*  
 $= 9.167 \times 10^3 \text{Kg/m}^3 \sqrt{1}$   
 $= 9167 \text{Kg/m}^3$

10. *Main scale reading = 3.1cm = 3.1cm*  
*Vernier scale reading = (4x 0.01) = 0.04cm*  
*Diameter of the marble = 3.13 x 10<sup>-2</sup>m*

11. *Volume of the marble = 4/3 π r<sup>3</sup>*  
 $= 4 \times 3.14 \times 1.565 \times 1.565 \times 1.565 \times 10^{-6} \quad \checkmark 1$   
 $= 10.0476 \times 10^{-6} \text{m}^3 \quad \checkmark 1$   
*Mass of the marble = 2.0 x 10<sup>-3</sup>kg*  
*Density of the disc = mass/volume*  
 $= \frac{2.0 \times 10^{-3}}{10.0476 \times 10^{-6}}$   
 $= 0.1246 \times 10^3$   
 $= 12.46 \times 10^2 \text{Kg/m}^3$

12.  $\frac{1}{2} \times 3 \sqrt{\quad} = 1.5\text{cm} \sqrt{1}$

13. a) *Volume of drop = 4/3 r<sup>3</sup>*  
 $V = \frac{4}{3} \times 22/7 \times (1.36/1000)^3 \sqrt{1}$   
 $= 1.054 \times 10^{-8} \text{m}^3$

b)  $\frac{4}{3}r^3 = R^2 t \sqrt{1}$

$$t = \sqrt[4]{\frac{1.36 \times 10^{-3}}{(4.0 \times 10^{-1})^2}} \sqrt{1}$$

$$t = \sqrt[4]{\frac{1.36 \times 10^{-9}}{4.0^2 \times 10^{-2}}}$$

$$= 0.2096 \times 10^{-7}$$

$$= 2.096 \times 10^{-8} \text{ m } \sqrt{1}$$

3 mks

c) *Lycopodium powder makes the film outline clearly visible*  $\sqrt{1}$

1 Mk

- d) - The film/ patch is a perfect circle  
 - The film is a monolayer  
 - There is no space between the molecules

An


14. Zero error + 0.04  
 Reading diameter = 0.93 - 0.04

15. Main scale = 5.5mm  
 Head scale coincidence =  $\frac{23\text{mm}}{100}$

Reading = 5.50  
 - 0.23  
 5.73mm

Actual reading = 5.73mm - 0.01mm = 5.72mm

16.



a ward 1m for the correct diagram showing the correct reading.  $\sqrt{1}$   
 No mark if the student uses 50 disks on the thimble scale.

### Turning effect of a force

1. Sum of clockwise moments = sum of anticlockwise moments

$$60\text{cm} \times 200\text{g} + 50\text{cm} \times Mg = 40\text{cm} \times 400\text{g} \text{ *TEZ*}$$

$$12000\text{cmg} + 50\text{mcmg} = 16000\text{cmg}$$

$$50M\text{cmg} = 4000\text{cmg}$$

$$50M = 4000$$

$$M = \frac{4000}{50} = 80\text{g}$$

$$w = 0.8\text{N}$$

2. Calculate the force F1

$$40 \times 5 = f \times 60 + 3 \times 10$$

$$200 = 60F + 30$$


$$60F = 170$$

$$F = \underline{170}$$

$$6\theta$$

$$= 2.833N$$

3. (a) Balloon and air will expand therefore up thrust on balloon increases thus clockwise moment increases ✓1  
 (c) (ii) Volume under water =  $\frac{3}{4} \times 40,000$   
 $= 30,000\text{cm}^3$  ✓
4. For a system in equilibrium the sum of clockwise moments about a point is equal to the sum of anticlockwise moments about the same point;



$$0.8 \times 2.5 = d \times 10$$

$$d = \frac{(0.8 \times 25)}{10} = 2.0 \quad (1mk)$$

$$\text{length} = (0.8 + 2) = 2.8 = 5.6\text{cm} \quad (1mk)$$

6.  $actm = ctm$   
 $x(0.3) + 2.0 \times 0.1 = (30 \times 0.2) + 2 \times 0.1 \quad \checkmark 1$   
 $0.3x = 6.2 - 2.0 \quad \checkmark 1$   
 $x = 14N \quad \checkmark 1$
7. Clockwise moments = Anticlockwise moments  
 $1.5 \times 0.1 = W \times 0.15 \quad \checkmark$   
 $W = \frac{0.15}{0.15}$   
 $W = 1N \quad \checkmark$

## Equilibrium and centre of gravity


1. a) This is the point of application of the resultant gravitational force on a body.  
 ▪ Appoint at which the weight of the body seems to cut  
 b) The figure below shows a wine glass  
 State how the stability of the glass is affected if it is filled with wine  
 ▪ The glass becomes more unstable since the c.o g is now raised
2. - Stability reduces /is lowered/becomes deny unstable less;  
 - position of C.D.C is raised / there is more mass at the atop; (Deny unstable)
3. - Increasing the base area/ making base heavier ✓1  
 - Increasing COG/ making base heavier ✓1
4.  $10w + (1.0 \times 6.0) = 2.0 \times 40 \quad \checkmark 1$   
 $10w + 60 = 80$   
 $W = \frac{20}{10} = 2N \quad \checkmark 1$
5. It remains vertical because the shape generated by the rotation is always symmetric, so its centre of gravity falls directly above the point of support. It therefore remains vertical for sometime.
6. As the wine fills the glass, the centre of gravity rises and this lowers the stability ✓1
7. Bisect any two sides and join the bisector to the apex where lines meet if the C.O.G

8. Clockwise Moment = Anticlockwise Moment

$$\begin{aligned} \frac{20 \times 30}{600} &= 30(f + s) \\ 600 &= 30f + 150 \\ 30f &= 450 \\ f &= 15 \text{ N} \end{aligned}$$

## Fluid flow


1.



2. The sum of potential energy, kinetic energy and pressure per unit volume of non-viscous fluid flowing is constant

3.  $A_1 V_1 = A_2 V_2;$   
 $7 \times 5 = 21 \times V_2;$   
 $V_2 = 1.667 \text{ m/s};$

4.



5. (i) For a non-viscous, non compressible fluid, in continuous flow, the pressure at any part of it is inversely proportional to the velocity of the fluid (1mk)

(ii) The gas jets in the barrel reduces pressure in the barrel hence the higher atmospheric pressure outside the barrel pushes air through the hole

6. Air flows over the top faster than air flow underneath; so air pressure on the top is less than air pressure underneath  $\sqrt{1}$

The difference in pressure causes the lift effect  $\sqrt{1}$


7. High speed of air reduces pressure above the mouth of the container  $\sqrt{1}$  higher pressure below the body pushes it up

8. Volume flux = Velocity x cross-sectional area  
 $= 5 \text{ m/s} \times 3.142 \times 10^{-4} \text{ m}^2$   
 $= 1.571 \times 10^{-3} \text{ m}^3/\text{s}$

## Hook's law

|                               |             |             |             |             |             |             |             |             |
|-------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Mass attached (kg)            | 0.15        | 0.30        | 0.45        | 0.60        | 0.75        | 0.90        | 1.05        | 1.20        |
| Force (load) on the spring(N) | <b>1.5</b>  | <b>3.0</b>  | <b>4.5</b>  | <b>6.0</b>  | <b>7.5</b>  | <b>9.0</b>  | <b>10.5</b> | <b>12.0</b> |
| Extension of spring (m)       | <b>0.04</b> | <b>0.08</b> | <b>0.12</b> | <b>0.16</b> | <b>0.20</b> | <b>0.25</b> | <b>0.30</b> | <b>0.34</b> |

1.




$$(c) \text{ slope} = \frac{0.16 - 0.05}{6.0 - 1.9} = \frac{0.11}{4.1} = 0.02683$$

$$\text{Spring constant} = \frac{1}{0.02683} = 37.27 \text{ Nm}^{-1}$$

(e) - Force  
- Acceleration

2.



$$3. \quad F_1 = Ke_1 = 40 = Ke_1$$

$$e_1 = \frac{40}{K}$$

$$F_2 = \frac{Ke_2}{K} = \frac{20}{K} = Ke_2$$

$$e_2 = \frac{20}{K}$$

$$\text{but } e_1 + e_2 = 20$$

Mocks Topical Analysis  $\checkmark 1$

$$\frac{40}{K} + \frac{20}{K} = 20\text{cm}$$

$$60 = 20k$$

$$K = 3\text{N/cm} \quad \checkmark 1$$

4. Diameter of coils/ Thickness/ No. of turns per unit length / length of spring are different  $\checkmark 1$

5. Upper springs,  $e = \frac{3.6}{3}$   
 Middle springs,  $e = \frac{3.6}{2} = 1.8\text{cm}$   
 Lower springs,  $e = \frac{3.6}{1} = 3.6\text{cm}$

$$\text{Total extension} = 1.2 + 1.8 + 3.6 = 6.6\text{cm}$$

6. a) Load on each spring  $= \frac{60}{3} = 20\text{N} \checkmark$   
 Extension (e) in one spring  $= \frac{F}{K} \checkmark$  for one spring  
 $= \frac{20}{50}$   
 $= 0.4\text{m} \checkmark$

b) The effective constant (K)  
 $= K_1 + K_2 + K_3 \checkmark$   
 $= 3(50)$   
 $= 150\text{N/M} \checkmark$

7. A load of (25 – 20)N causes extension of (17 – 15) cm.  
 i.e. 5N causes extension of 2 cm  $\checkmark 1$

$$20\text{N} = ?$$

$$\frac{20\text{N}}{5\text{N}} \times 2\text{cm} = 8\text{cm} \quad \checkmark 1$$

When no mass is hung.  
 Length of the spring  $= 15\text{cm} - 8\text{cm}$   
 $= 7\text{cm} \quad \checkmark 1$

a) Uniform velocity :- The change in displacement for equal time intervals is the same.  
 Uniform acceleration:- Change in velocity for equal time intervals is the same.  
 b) Determine the acceleration of the trolley pulling the tape

$$V_a = \frac{2}{0.02} = 100\text{cm/s} \quad V_b = \frac{3}{0.02} = 150\text{cm/s} \quad a = \frac{V-U}{t}$$

$$= \frac{(150 - 100)}{(7 \times 0.02 - 0.02)}$$

$$a = 416.67\text{ cm/s}^2$$

c) i) Determine the motion of the ball relating it to its different positions along the following  
 I AB the body is projected upwards and at B  $V = 0$   
 II BC the body falls back to the starting point (moving in the opposite direction)  
 III CE the body rebounds on the ground (at starting point) and starts moving up again

ii) From the graph calculate the acceleration due to gravity

$$a = \frac{v-u}{t} \quad \left| \quad \begin{array}{l} a = -10\text{m/s}^2 \\ = 10\text{m/s}^2 \end{array} \right.$$

$$= \frac{0-20}{2}$$



## Linear motion

1. a) *Uniform velocity* :- The change in displacement for equal time intervals is the same.  
*Uniform acceleration* :- Change in velocity for equal time intervals is the same.  
 b) Determine the acceleration of the trolley pulling the tape

$$V_a = \frac{2}{0.02} = 100 \text{ cm/s} \quad V_b = \frac{3}{0.02} = 150 \text{ cm/s} \quad a = \frac{V-U}{t}$$

$$= \frac{(150 - 100)}{(7 \times 0.02 - 0.02)}$$

$$a = 416.67 \text{ cm/s}^2$$

- c) i) Determine the motion of the ball relating it to its different positions along the following  
 I AB the body is projected upwards and at B  $V = 0$   
 II BC the body falls back to the starting point (moving in the opposite direction)  
 III CE the body rebounds on the ground (at starting point) and starts moving up again

ii) From the graph calculate the acceleration due to gravity

$$a = \frac{v-u}{t} \quad \left| \quad \begin{array}{l} a = -10 \text{ m/s}^2 \\ = 10 \text{ m/s}^2 \end{array} \right.$$

$$= \frac{0-20}{2}$$

## 2. Conduction

Free electrons at the heated end gain more kinetic energy and spread the heat energy to other parts of the rod

3. (a) Momentum before collision = momentum after collision

$$\frac{150 \times 80}{1000} = 2.65 \times V$$

$$16 = 2.65V$$

$$V = \frac{16}{2.65}$$

$$= 6.0377$$

But  $\frac{1}{2} mV^2 = mgh$

$$h = \frac{V^2}{2g} = \frac{(6.0377)^2}{2 \times 10}$$

$$h = \frac{36.4538}{20}$$

$$= 1.82269 \text{ m}$$


(b) The block will be deformed

4. Total distance = Area under graph;  
 $= \frac{1}{2} (12 + 5) \times 20$ ; OR  
 $= 170 \text{ m}$ ;

$$\text{Average speed} = \frac{170 \text{ m}}{12 \text{ s}}$$

$$= 14.17 \text{ m/s}$$

5.



✓1

6. (i) Acts as a thermostat  
(ii) On closing, the switch, current becomes complete, the current flows causing heating effect, the bimetallic strip bends downwards and contents separates.  
when the room becomes cool the strip bends upward completing the current and the process repeats itself on and off regulating the temperature  
- Weight of the fluid in which it floats

7. Clockwise moments = anticlockwise moments at equilibrium

$$0.6 \times 0.7 = W \times 0.3;$$

$$W = 0.6 \times 0.7$$

$$0.3$$

$$= 1.4N;$$

$$0.6 + R = 1.4$$

$$R = 0.8N;$$


8. (a) Length = area under curve

$$= 10 \times (32-18);$$

$$= 10 \times 14 = 140m;$$

(b)  $10-25 = -15; = -1.875ms^{-1}$

$18-10 \quad 8 \quad Decal = 1.875ms^{-1};$



(ii) Keep the train in circular motion;

(iii) Friction force between the wheels and rails;

(d) (i)  $F = \frac{m(v-u)}{t}$

$$= \frac{90(0-45)}{1.2};$$

$$1.2$$

$$= -3375N;$$

(ii) Kinetic energy – Heat + sound + P.E(deformation);

(iii)  $E = \frac{1}{2} Mv^2;$

$$= \frac{1}{2} \times 90 \times 45^2;$$

$$= 91,125J;$$

9. (a) (i) When a body is projected vertically upwards, it undergoes a uniform retardation due to the gravitational pull. The body thus slows down, comes to rest then starts falling with an increasing velocity (in opposite direction)

(ii) Acc of free fall = gradient / slope of the graph

$$= \frac{5-0}{3-0} = \frac{5}{3} = 1.66ms^{-2}$$

$$3-0 \quad 3$$

(iii) Total distance = Area under the curve

$$(\frac{1}{2} \times 5 \times 3) + (\frac{1}{2} \times 2 \times 3.3)$$

$$\frac{15}{2} + \frac{10}{3} = \frac{30}{6} + \frac{20}{6} = \frac{50}{6} = 25 = 8\frac{1}{3}m$$

$$2 \quad 3 \quad 6 \quad 6 \quad 3$$

(iv) -  $Wt$  in the moon =  $mg = 2kg \times \frac{5}{3} = \frac{10}{3} = 3\frac{1}{3}N$

(v) - It will accelerate faster at  $10ms^{-1}$  from the graph

- It will attain a maximum height after  $\frac{1}{2}$  second

(b)  $V_1 = \frac{1.5cm}{0.025} = 75cms^{-1} = V_2 = \frac{3.0cm}{0.02s} = 150cms^{-1}$

$$0.025$$

$$0.02s$$

$$a = \frac{V_2 - V_1}{t} = \frac{150 - 75}{0.02 \times 4} = \frac{75}{0.08} = 937.5 \text{ cm}^{-2} \text{ or } 9.375 \text{ ms}^{-2}$$

$$S = ut + \frac{1}{2} at^2$$

$$= (0.75 \times 4) + \frac{1}{2} \times 9.375 \times 4^2 = 3 + 75 = 78 \text{ m}$$

10. a) The volume of a fixed mass of gas is inversely proportional to its pressure provided temperature is kept constant.

(b) (i) The bubble expands as it comes up finally bursts when at the surface

$$(ii) p_1 V_1 = P_2 V_2$$

$$(76 + 30) \times 3 = (76 + 5) V_2$$

$$106 \times 3 = 81 \times V_2$$

$$V_2 = \frac{106 \times 3}{81}$$

$$= \frac{81}{81}$$

$$= 3.93 \text{ cm}^3$$

(c)  $100^\circ\text{C} - 0^\circ\text{C} = 98 - 11$

$$1 \text{ division} = \frac{87}{100}$$

$$= \frac{87}{100}$$

$$\text{Reading} = \frac{8 \times 56}{1000}$$

$$= \frac{448}{1000}$$

$$= 48.72^\circ\text{C}$$

11. Distance traveled  $\sqrt{l}$

12. a) A body continues with its state of rest or uniform motion unless acted upon by some external forces  $\sqrt{l}$

b) i)  $s = \frac{1}{2} gt^2$

$$720 = \frac{1}{2} \times 10 \times t^2 \sqrt{l}$$

$$t^2 = 144$$

$$t = \sqrt{144} = 12 \text{ sec } \sqrt{l}$$

ii) Range =  $ut$

$$= 120 \times 12 \sqrt{l}$$

$$= 1440 \text{ m } \sqrt{l}$$

c) i) – Centripetal force acting on  $M_1 \sqrt{l}$

- Weight ( $M_2 g$ ) acting on  $M_2 \sqrt{l}$

ii)  $M_2$  moves upwards;  $\sqrt{l}$

When the speed of  $M_1$  increases centripetal force remains the same, the radius of the circle described by  $M_1$  increases  $\sqrt{l}$

iii) Centripetal force = weight of  $M_2$

$$M_1 V^2 / r = M_2 g$$

$$0.020 V^2 / l = 0.050 \times 10 \sqrt{l}$$

$$V^2 = 0.5 / 0.02 = 25 \sqrt{l}$$

$$V = \sqrt{25} = 5 \text{ m/s } \sqrt{l}$$

13. (a) Constant rate of change of displacement with time  $\sqrt{l}$

OR- A body is said to be moving with uniform velocity if its rate of change of displacement with time is constant

(b) (i) For one correct tangent

$$\text{Velocity } t = 1 \text{ s} = \frac{42 - 20}{2 - 0.5}$$

$$= \frac{22}{1.5}$$

(correct reading from graph and expression)

$$= 14.67 \text{ m/s}$$

$$\begin{aligned} \text{Velocity at } t = 4s &= \frac{67.5 - 30}{5 - 0.5} \quad (\text{correct reading from graph and expression}) \\ &= 8.33\text{m/s (accuracy)} \quad \checkmark 1 \end{aligned}$$

$$\begin{aligned} \text{(ii) } a &= \frac{V - u}{t} = \frac{8.33 - 14.67}{4 - 1} = \frac{6.34}{3} \quad \checkmark 1 \\ &= 2.11\text{m/s}^2 \quad \checkmark 1 \end{aligned}$$

$$\begin{aligned} 14. \quad S &= \frac{1}{2}gt^2 \quad \text{since } u = 0 \\ &= \frac{1}{2} \times 10 \times 4 \times 4 \\ &= 80\text{m} \end{aligned}$$

$$\begin{aligned} 15. \quad \text{b) } \quad \text{i) } t &= \frac{v - u}{g} \quad \checkmark \\ &= \frac{0 - 60}{-10} \quad \checkmark \\ &= 6 \text{ secs} \quad \checkmark \\ \quad \quad \text{ii) } h &= ut - \frac{1}{2}gt^2 \quad \checkmark \\ &= 60 \times 6 - \frac{1}{2} \times 10 \times 6^2 \quad \checkmark \\ &= 360 - 180 \\ &= 180\text{m} \quad \checkmark \\ \quad \quad \text{iii) } V^2 &= U^2 + 2aS \quad \checkmark \\ 0 &= U^2 + 2 \times -10 \times 5 \quad \checkmark \\ 0 &= U^2 - 100 \\ U &= 10\text{m/s} \quad \checkmark \end{aligned}$$

c) Resistance/ friction with air is negligible

$$\begin{aligned} 16. \quad \text{a) Length of nine dots} &= 6.9\text{cm} \quad \checkmark \\ \text{Time taken} &= \frac{1}{50} \times 9 = 0.02 \times 9 \\ &= 0.185 \quad \checkmark \\ \text{Velocity} &= \frac{6.9\text{cm}}{0.18\text{s}} \quad \checkmark \\ &= 38\text{cm/s or } 0.38\text{m/s} \quad \checkmark \end{aligned}$$

$$\begin{aligned} \text{b) } Ft &= 0.8 \times 0.5 - 0.8 \times 0.38 \quad \checkmark \\ Ft &= 0.096 \\ F &= \frac{0.096}{0.18} = 0.533\text{N} \end{aligned}$$

17. a) Equations of linear motion.

$$\begin{aligned} \text{i) } V &= u + at. \quad \checkmark 1 \\ \text{ii) } V^2 &= u^2 + 2as \quad \checkmark 1 \\ \text{iii) } S &= ut + \frac{1}{2}at^2. \quad \checkmark 1 \end{aligned}$$

$$\text{b) } \frac{100 \text{ km/h} \times 10}{36} = 27.78\text{m/s}$$

$$\text{In } 0.5 \text{ sec the driver covers } 27.78\text{m/s} \times 0.55 = 13.89 \text{ M} \quad \checkmark 1$$

After applying brake

$$a = -4\text{m/s}^2$$

$$u = 27.78\text{m/s.}$$

$$v = 0$$

$$\therefore v^2 = u^2 + 2as \quad \checkmark 1$$

$$-2as = u^2. \text{ since } v = 0$$

$$S = u^2 = (27.78\text{m/s})^2 = 96.47\text{M} \checkmark 1$$

$$-2a \quad (-2) \quad (-4\text{m/s}^2)$$

$$\text{Total distance covered} = (13.89 + 96.476)\text{M} = 110.36 \text{ M} \checkmark 1$$

c) (i) See the graph paper

(ii)

$$(i) \text{ Average speed of the whole journey} = \frac{\text{Total distance covered}}{\text{Total time taken}}$$

Distance = Area under the graph

$$= (20\text{m/s} \times 50\text{s}) + (\frac{1}{2} (20 + 25) \times 10) + \frac{1}{2} (50 + 65) \times 25$$

$$= 1000\text{m} + 225\text{m} + 1437.5\text{m} = 2662.5\text{m} \checkmark 1$$

Total time = 125s

$$\text{Speed} = \frac{2662.5}{125\text{s}} = 21.3 \text{ m/s} \checkmark 1$$


$$= 21.3 \text{ m/s} \checkmark 1$$

(ii)  $a = v - u$

$$= \frac{25 - 20}{10 \text{ s}} \text{ m/s} \checkmark 1$$

$$= 0.5\text{m/s}^2 \checkmark 1$$

18.



## Machines & inclined planes

1.  $M.A = \frac{600\text{M}}{650\text{M}} = 0.92307$

$$V.R = \frac{1}{\sin 25} = 2.366$$

$$\eta = \frac{M.A}{V.R} = \% 100$$

$$= \frac{0.92307}{2.366} \times 100 = 39.01\%$$

2. (a)  $V.R = 5$

(b)  $\frac{M.A}{E} = L \checkmark$

$$= \frac{4000}{1000} \checkmark$$

$$= 4 \checkmark$$

(c)  $\text{eff.} = \frac{M.A}{V.R} \times 100\%$

$$= \frac{4}{5}$$

$$= 80\%$$

(d) Some work is done overcoming friction or lifting the moving parts

(e)  $W = F \times d$

$$= 40,000 \times 0.05 = 2000\text{J}$$

$$\begin{aligned}
 3. \quad &VR = 4 \\
 &A = L = 100 \\
 &E = 48 \\
 &\therefore u = \frac{M.A}{V.R} \times 100\% \\
 &= \frac{25}{48} \times 100 = 52.08\%
 \end{aligned}$$

4. (a) (i) Velocity ratio is the distance moved by the effort to the distance moved by the load in the same time  
(ii) - Pulley belts  
- Gears (any one)

(b) Graph

$$\begin{aligned}
 (c) \text{ (i) } V.R &= \frac{R^2}{r^2} = \frac{7 \times 7}{1.4 \times 1.4} = 25 \\
 \text{Efficiency} &= \frac{M.A \times 100\%}{V.R} \\
 M.A &= r \times V.R = \frac{80}{100} \times 25 = 20 \\
 E &= \frac{KL}{M.A} = \frac{100}{20} \times 10 = 50N
 \end{aligned}$$

$$\text{(ii) } EH = \frac{\text{work output} \times 100\%}{\text{Work input}}$$


$$\begin{aligned}
 \text{Work output} &= mgh \\
 &= 100 \times 10 \times 2.5 \\
 &= 2500J
 \end{aligned}$$

$$80 = \frac{2500 \times 100}{\text{Work output}}$$

$$\text{Work output} = \frac{2500 \times 100}{80} = 3125J$$

$$\begin{aligned}
 \text{Energy lost} &= 3125 - 2500 \\
 &= 625J
 \end{aligned}$$

5. i)



$$\begin{aligned}
 \text{ii) } E &= L/M.A \\
 &= 60/3 \\
 &= 20N \checkmark
 \end{aligned}$$

$$\begin{aligned}
 \text{Total work done by effort} &= E \times \text{Distance moved by effort} \checkmark \\
 &= 20 \times 9 \times V.R \\
 &= 20 \times 9 \times 4 \\
 &= 720J \checkmark
 \end{aligned}$$

## Newton's law

1. (a) Newton's first law:  
- A body remains in its state of rest or uniform motion in a straight line unless acted upon by an external force \*TEZ\*
- (b) Elastic collision is one in which both kinetic energy and momentum are conserved, while inelastic collision is one in which momentum is conserved, but kinetic energy is not (2mks)
- (c) Momentum before collision = momentum after collision (1mk)  

$$(2000 \times 10) + (1000 \times 0) = (2000 + 1000)V$$

$$3000V = 20000$$

$$V = 6\frac{2}{3} \text{m/s}$$
- (ii)  $d = \text{Velocity} \times \text{time}$  \*TEZ\*  

$$= \frac{20}{3} \times 20 = \frac{400\text{m}}{3} = 133\frac{1}{3}\text{m}$$
- (iii) Impulse = change in momentum  

$$= 2000 (10 - 6\frac{2}{3}\text{m/s}) - \text{for the minibus}$$

$$\text{Or} = 1000 (6\frac{2}{3}\text{m/s} - 0) - \text{for the car}$$

$$= 6667\text{NS}$$
- Impulse of force =  $\frac{\text{Impulse}}{\text{Time}}$   

$$= \frac{6667}{2} = 3333.5\text{N}$$
- (iv) K.E before collision =  $\frac{1}{2} \times 2000 \times 10^2 = 100,000\text{J}$   
 K.E after collision =  $\frac{1}{2} \times 3000 \times (6\frac{2}{3})^2 = 66,666.7\text{J}$   
 Change in K.E =  $(100,000 - 66666.7) \text{J}$   

$$= 33,333.3\text{J}$$

2. The rate of change of momentum is directly proportional to the external force acting on a body it is in the direction of force
3. A body continues in its initial state rest or uniform motion unless compelled by an external force to make it behave differently.

4.  $F = Ma$   
 $900 = 600a$   
 $100a = \frac{9000}{10} = 900\text{ms}^{-2} \checkmark 1$  (1mk)

but  $a = \frac{v-u}{t}$

$$at = v-u$$

$$(900 \times 0.1) = v = 90\text{ms}^{-1} \checkmark 1$$
 (1mk)

✓1

5. For every action there is an equal and opposite reaction ✓1
6. (a) The rate of change of momentum of a body is directly proportional to the resultant external force producing the change and takes place in the direction of force.

- (b)  $m_1u_1 - m_2u_2 = (m_1 + m_2)V$   

$$V = \frac{m_1u_1 - m_2u_2}{M_1 + m_2}$$

$$= \frac{(2500 \times 40) - (3500 \times 20)}{2500 + 3500}$$

$$= \frac{30000}{6000} = 5\text{ms}^{-1}$$

## Circular motion

1. The direction of the speed keeps changing hence the velocity at each point on the circular path is given by the tangent to the path at that point
2. a) This is the rate of change of angular displacement with time.  
 b) The tension  $T$ , on the string was measured for various values of angular velocity,  $\omega$ . The

distance  $r$  from the centre was maintained at 30cm. The results are as shown below

|   |      |      |      |      |      |
|---|------|------|------|------|------|
| Angular velocity $\omega$ ( $\text{rad s}^{-1}$ ) | 2.0  | 3.0  | 4.0  | 5.0  | 6.0  |
| Tension $T$ (N)                                   | 0.04 | 0.34 | 0.76 | 1.30 | 1.96 |
| $\omega^2$ ( $\text{rad s}^{-2}$ )                | 4    | 9    | 16   | 25   | 36   |

i) Plot the graph of  $T$  ( $y$  - axis) against  $\omega^2$

ii) From the graph, determine the mass,  $m$ , of the body given that

$$T = \omega^2 r - C$$

Where  $C$  is a constant

$$\text{Gradient} = Mr$$

$$M = \frac{\text{gradient}}{r}$$

$$\text{Gradient} = 0.76 - 0.04$$

$$16 - 4$$

$$= 0.06 \text{ N} / (\text{rad}^2 \text{ s}^{-2})$$

$$= \frac{0.06}{\frac{30}{100}} = \frac{0.6}{3} = 0.2 \text{ kg}$$

iii) Determine the constant  $C$  and suggest what it represents in the set up

$C$  is the  $Y$ -intercept

$$C = -0.2 \text{ N}$$

3. (a) It keeps changing direction and hence must experience centripetal acceleration

(b) (i)  $f = 0.5 \text{ Hz}$

$$T = \frac{1}{f}$$

$$= \frac{1}{0.5}$$

$$= 2 \text{ sec}$$

(ii)  $\omega = \frac{2\pi}{T}$ ;

$$T$$

$$= \frac{2 \times 3.142}{2}$$

$$= 3.142 / \text{sec};$$

(iii)  $V = r\omega$ ; or

$$= 6 \times 3.12 = 18.852 \text{ m/s}$$

(iv)  $F = \frac{MV^2}{r}$

$$r$$

$$= 30 \times \frac{(18.852)^2}{6}$$

$$6$$

$$= 1776.99 \text{ N}$$

OR  $F = mr\omega^2$

$$= 30 \times 6 \times (3.142)^2 = 1776.99 \text{ N}$$

✓1

(c) Graph

(iii) centrifugal

5. (a) (i) The direction of the particle is tangential to the path of any given point. There is instantaneous change in direction of velocity, this causes acceleration of the particle.

(ii)  $F = \frac{mV^2}{r}$

$$r$$

$$= \frac{1.5 \times 10^3 \times 10 \times 10}{20} \quad \checkmark 1$$

$$= 7.5 \times 10^3 \text{ N} \quad \checkmark 1$$

(b) (i) The value of  $r$  increases. Increase in speed leads to increase in centripetal force on the bob. This leads to increase in radius of path (centripetal force is directly proportional to radius)



(c) The cyclist leans inwards in order to have enough component of the contact force to provide adequate centripetal force.

6. a) The rate of change of angular displacement with time  $\sqrt{1}$

$$\begin{aligned} \text{b) i) } T_A &= \frac{mv^2}{r} - mg \quad \sqrt{1} \\ &= \frac{0.2 \times 5^2}{0.5} - 2.0 \quad \sqrt{1} \\ &= 8N \quad \sqrt{1} \end{aligned}$$

$$\begin{aligned} \text{ii) } T_B &= \frac{mv^2}{r} \\ &= \frac{0.2 \times 5^2}{0.5} = 10N \quad \sqrt{1} \end{aligned}$$

$$\begin{aligned} \text{iii) } T_C &= \frac{Mv^2}{r} + mg \quad \sqrt{1} \\ &= \frac{0.2 \times 5^2}{0.5} + 2 \quad \sqrt{1} \\ &= 12N \quad \sqrt{1} \end{aligned}$$

c) At point C where tension is maximum  $\sqrt{1}$

$$\begin{aligned} \text{d) i) } T \cos Q &= mg \quad \sqrt{1} & T(0.3/0.5) &= 1.5 & T &= 2.5N \sqrt{1} \\ T \cos Q &= \frac{150}{1000} \times 10 \quad \sqrt{1} & T &= \frac{1.5 \times 5}{3} \end{aligned}$$

$$T \cos Q = 1.5$$

ii) Speed governor  $\sqrt{1}$

7. (a) The direction of its velocity is continuously changing (1mk)

$$\text{(b) (i) } Fr = mw^2r \quad (1mk)$$

$$0.4 = 0.2 \times w^2 \times 0.08$$

$$w^2 = \underline{0.4}$$

$$0.2 \times 0.08 \quad (1mk)$$

$$w^2 = 25 \text{ and } 25^{-2}$$

$$w = 5 \text{ rad } 6^{-1} \quad (1mk)$$

$$\text{(ii) } F = mw^2r$$

$$= 0.2 \times 5 \times 5 \times 0.12$$

$$= 0.6N \text{ (must be shown)}$$

(c) The block will slide (1mk)

Frictional force (0.4N) is less than the force required to maintain it in uniform circle (1mk)

$$8. \quad \alpha = rw^2 = 0.2(16\pi)^2 \quad \sqrt{1}$$

$$= 505.3 \text{ m/s}^2 \quad \sqrt{1}$$

9. a) i) Centrifugal force  $\sqrt{1}$

ii) The gradient of the graph = mass x radius  $\sqrt{1}$

$$T = MW^2r - C$$

From  $Mr = \text{gradient}$

$$Mr = \frac{1.30 - 0.76}{25 - 16} = 0.06 \quad \sqrt{1}$$

$$Mr = 0.06$$

$$M \times 0.3 = 0.06 \quad \sqrt{1}$$

$$M = \frac{0.06}{0.3} = 0.2 \text{ kg} \quad \sqrt{1}$$

iii) y - intercept = -0.2N

$$-0.2 = -C$$

$$0.2 = C\sqrt{v}$$

Frictional force  $\propto v$

10. (a) at the highest point of the circle

$$T = \frac{Mv^2}{r} - mg$$

$$\frac{Mv^2}{R} = Fe = MW^2R$$

$$\therefore T = Mw^2r - mg$$

$$\text{But } w = 2\pi f \checkmark 1 \text{ mk}$$

$$T = (2\pi \times 10)^2 \times 2\text{kg} \times 0.5 - (2 \times 10)$$

$$= 400\pi^2 - 20 = 3927.84 \text{ N } \checkmark 1 \text{ mk}$$

b) T at the lowest point

$$T = Fe + Mg$$

$$= Mw^2r + Mg$$

$$= 400\pi^2 + 20$$

$$= 3967.84 \text{ N } \checkmark 1 \text{ mk}$$

### Quality of heat

1. a) Specific heat capacity is defined as the quantity of heat required to raise the temperature of a unit mass of a substance by one Kelvin.  $\checkmark 1 \text{ mk}$

b) (i) heat gained by calorimeter

$$Q_E = C\Delta\theta$$

$$= 400\text{Jk}^{-1} \times (34 - 20)\text{k} \checkmark 1 \quad = 560\text{j} \checkmark 1$$

(ii) Heat gained by water

$$Q_W = MC\Delta\theta = 200 \text{ kg} \times 4200\text{kg}^{-1}\text{k}^{-1} \times (34 - 20)\text{k}$$

$$= 11760\text{j} \checkmark 1$$

(iii) Heat lost by the metal = heat gained by the calorimeter + heat gained by Water.

$$= 560\text{j} + 11760\text{j} = 12320\text{j} \checkmark 1$$

(iv) Specific heat capacity of the metal.

$$C = \frac{Q}{M\Delta\theta} \checkmark 1$$

$$= \frac{12320\text{j}}{0.3 \text{ kg} \times (100 - 34)}$$

$$= \frac{12320\text{j}}{0.3\text{kg} \times 66 \text{ k}} = 622.22 \text{ jkg}^{-1}\text{k}^{-1} \checkmark 1$$

2. (a)

| Boiling   | Evaporation                                    |
|---|--|
| - Takes place at constant temperature             | - Takes place at all temperatures              |
| - Not affected by surface area of liquid exposure | - Increases with increase in surface area      |
| - Vigorous, visible process all over the liquid   | - Slow invisible process at the liquid surface |

(each for any two correct)

(b) (i) A – Temperature of the solid is increasing;

AB- solid is melting;

(ii) Heat supplied = Heat gained

$$p \times t = MC\Delta\theta$$

$$200 \times 100 = 0.2 \times C \times 100$$

$$20 \times 100 = 0.2 \times C \times (350 - 250)$$

$$C = 1000 \text{JKg}^{-1} \text{K}^{-1};$$

(iii) Heat applied = heat gained; OR

$$p \times t = Mlf;$$

$$200 \times 200 = 0.2 \times lf;$$

$$lf = 200,000 \text{J/Kg};$$

3. (a) The quantity of heat required to raise the temperature of a given mass of a material by one Kelvin.

(b) (i)  $C = \frac{Q}{\theta}$

$$Q = C\theta$$

$$= 40 \times (34-25)$$

$$= 40 \times 9 = 360 \text{J}$$

(ii)  $M_w C_w D = (0.1 \times 9 \times 4200) = 3780 \text{J}$

(iii) Heat lost = heat gained by calorimeter + heat gained by water  
 $= 3780 + 360 = 4140 \text{J}$  or  $4.14 \text{KJ}$

(b)  $\frac{150}{1000} \times C_m \times \Delta\theta = 4140$   
 $0.15 \times (100 - 34) C_m = 4140$   
 $9.9 C = 4140$   
 $C_m = \frac{4140}{9.9} = 48.18 \text{JKg}^{-1} \text{K}^{-1}$

(b)

| Boiling                               | Evaporation                    |
|---------------------------------------|--------------------------------|
| - Temperature is constant             | - Temperature can vary         |
| - Affected by impurities and pressure | - Not affected with impurities |
| - vigorous with bubbles               | - not vigorous, no bubbles     |
| - takes place in the whole liquid     | - takes place at the surface   |

4. (a) Evaporation is change of liquid to gas at any temperature while boiling is change of liquid to gas at constant temperature;

(i) Heat  $= ml_v + MC\Delta\theta$   
 $= 0.44 \times \frac{2.26}{1000} \times 10^6 + 0.44 \times \frac{4200}{1000} \times 100;$   
 $= 994.4 + 184.8 = 1179.2 \text{J};$

(ii)  $Q = \frac{mLf}{t}$   
 $= 4.4 \times 10^{-3} \times 3.34 \times 10^5$   
 $= 1336 \text{J};$

(iii) - Heat is absorbed from the surrounding since the can is unlagged;

(c) A = Vacuum;

B = Silvered (shiny) surfaces/polished/smooth;

(ii) - Loss of heat by conduction is reduced by cork and vacuum;

- Loss of convection is reduced by vacuum;

- Loss by radiation is reduced by silvered wall;

5. Thermometer A gives a higher reading than B;  $\sqrt{1}$  Black surface is better absorber of heat than silvery surface  $\sqrt{1}$

5. Water at the top part of the boiling tube boils while the ice remains unmelted. This is because water is a poor conductor of heat and hot air less dense remain at the top.

6. – Evaporation is silent while boiling is a vigorous visible process  
 - Evaporation takes place at all temperatures while boiling takes place at a specific temperature  
 - Evaporation takes place on the surface of the liquid but boiling takes place in the whole liquid
7. Steel is a good conductor of heat; therefore draws heat from your body unlike wood which is a poor conductor of heat
8.  $Pt = MC\Delta T$   $\Delta T = 35 - 15 = 18K$   
 $T = \frac{MC\Delta T}{P}$   
 $= \frac{5kg \times 400JKg^{-1}K^{-1} \times 18K}{1000Js^{-1}}$   
 $= 36s$
9. (a) It is the quantity of heat required to convert a unit mass of the substance from the solid to the liquid state without change of temperature (1mk)  
 Heat lost by naphthalene  
 $T = 100 - 80 = 20k$   
 $H_1 = 0.5kg \times 2100JKg^{-1}K^{-1} \times 20K = 21000J$   
 $L_f = mL_f = 0.5kg \times 170000JKg^{-1} = 85000J$   
 $T = 80 - 20 = 60k$ ,  $H_2 = 0.5kg \times 2100JKg^{-1}K^{-1} \times 60k = 63000J$   
 Heat lost by aluminium  
 $T = 100 - 20 = 80k$   
 $H = 0.4kg \times 900JKg^{-1}K^{-1} \times 80k = 28800J$   
 Total heat lost  $= 169000J + 28800J = 197800J$   
 $= 197.8KJ$   
 $= 198KJ$
- (c) Blowing wind over the surface of the liquid increases the kinetic energy of the liquid molecules  
 - It can also be made to evaporate faster by bubbling air through the liquid as it increases the surface area from which the liquid molecules may escape
- (d) (i) – Charcoal is a black body and therefore a better absorber of heat  
 - Sprinkled with water, so that it takes latent heat from the air around the air around and evaporates, causing cooling in cabinet  
 (ii) Heat is conducted by the by the metallic tank and the wire mesh to the surrounding air
10. Loss of heat through radiation is reduced/ minimized ✓

### **Work, energy and power**

1. (a) The law of conservation of energy states that the sum of kinetic energy and potential energy of a system is a constant
- (b) (i) At P and T potential energy is a maximum and kinetic energy is a minimum. Hence velocity is zero (2mks)  
 (ii) At Q and S P.E has reduced by 0.1J. This equals the K.E  
 $K.E = \frac{1}{2} MV^2$   
 $0.1 = \frac{1}{2} \times 0.8V^2$   
 $0.1 = 0.4V^2$   
 $0.1 = V^2$   
 $V^2 = \frac{1}{4} = 0.25$   
 $V = 0.5m/s$
- (iii) At R, auP.E has been converted to K.E velocity now is a maximum  
 So,  $0.2 = \frac{1}{2} MV^2$   
 $0.4 = V^2$   
 $\sqrt{V} = 0.4m/s$

$$V = 0.64\text{m/s}$$

$$(c) (i) M.A = \frac{L}{E} = \frac{280\text{N}}{40\text{N}} = 7 \text{ (2mks)}$$

$$(ii) V.R = \frac{P}{R} = \frac{70}{5} = 14 \quad (2\text{mks})$$

$$(iii) n = \frac{M.A}{V.R} \times 100\%$$

$$= \frac{7}{14} \times 100\%$$

$$= 50\% \quad (2\text{mks})$$

$$2. a) (i) CR = \frac{2\pi R}{2\pi r} = \frac{\text{No of teeth draw}}{\text{No. of teeth of driven}} \quad \checkmark 1$$

$$\therefore \frac{16.5 \text{ cm}}{r} = \frac{44}{16}$$

$$r = \frac{16.5 \text{ cm} \times 16}{44} \quad \checkmark 1$$

$$R = 6 \text{ cm} \quad \checkmark 1$$

$$(ii) V.R. = \frac{R}{r} \quad \checkmark 1$$

$$= \frac{16.5 \text{ cm}}{6 \text{ cm}} \quad \checkmark 1$$

$$= \underline{\underline{2.75}} \quad \checkmark 1$$

$$b) (i) M.A. = \frac{L}{E} \quad 120\text{kg} \times 10\text{N/kg} = 1200\text{N} \quad \checkmark 1$$

$$= \frac{1200\text{N}}{300\text{N}} \quad \checkmark 1$$

$$= 4 \quad \checkmark 1$$

$$(ii) \text{ Its efficiency of:}$$

$$D = \frac{M.A \times 100\%}{V.R.} \quad \checkmark 1$$

$$= \frac{4 \times 100\%}{6}$$

$$= 66.67\% \quad \checkmark 1$$

3. (a) Work is said to be done when the body on which a force is applied moves in the direction of force; S.I unit if the Joule, J or (Nm);

$$(b) (i) \text{ Work done} = P.E \text{ gained ;}$$

$$= mgh$$

$$= 500 \times 4 \times 10;$$

$$= 20,000\text{J};$$

$$(ii) \text{ Power} = \frac{\text{work done}}{\text{time taken}}$$

$$= \frac{20,000}{8}$$

$$= 2.5\text{KW} ; \text{ or } (2500 \text{ watts});$$

$$(iii) \text{ Efficiency} = \frac{\text{work output} \times 100}{\text{work input}}$$

$$= 2.5 \times 100;$$

$$2.8$$

$$= 89.29\%;$$

- (iv) – Friction between movable parts  
 - Sound due to moving parts  
 - - heat –some of the electrical energy is converted to unnecessary heat

### Floating and sinking

1. a)(i) R.d. = Weight of solid

$$\frac{\text{Upthrust in water}}{= 3N}$$

$$\frac{(3 - 0.22)N}{2.78} \checkmark 1$$

$$= 1.079 \checkmark 1$$

- (ii) Its apparent weight in a liquid of density  $800 \text{ kgm}^{-3}$ . R.d of the liquid = Upthrust in the liquid

$$\frac{\text{Upthrust in water}}{\text{R.d of the liquid}} = \frac{800 \text{ kgm}^{-3}}{1000 \text{ kgm}^{-3}} = 0.8 \checkmark 1$$

$$0.8 = \frac{u}{2.78 \text{ N}}$$

$$u = 2.78 \times 0.8$$

$$= 2.224$$

$$\text{Upthrust } u = 2.224 \text{ N } \checkmark 1$$

$$\text{Apparent weight of liquid} = \text{weight in air} - \text{upthrust in liquid}$$

$$= 3.0 - u - 2.224 \text{ N} = 0.776 \text{ N } \checkmark 1$$

2.  $P_1 V_1 = P_2 V_2 \checkmark 1$

$$P_1 = A + h \rho g = 100 \text{ 000 Nm}^{-2} + (0.5 \text{ m} \times 1000 \text{ kgm}^{-3} \times 10 \text{ N/Kg}) \checkmark 1$$

$$P_1 = 105000 \text{ Nm}^{-2}$$

$$P_2 = 100 \text{ 000 Nm}^{-2} \text{ i.e only Atmospheric pressure}$$

$\therefore$  Volume is density proportional to  $R^3$ .

$$\therefore P_1 R^3 = P_2 R_2^3$$

$$R^3 = \frac{P_1 R^3}{P_2} = \frac{105000 \text{ Nm}^{-2} (1 \times 10^{-3})^3}{100000 \text{ Nm}^{-2}} \checkmark 1$$

$$R^3 = 1.05 \times 10^{-9} \text{ m}$$

$$R = \sqrt[3]{1.05 \times 10^{-9}} = 1.0164 \times 10^{-3} \text{ m}$$

$$D = 2.0328 \times 10^{-3} \text{ m or } 2.0328 \text{ mm } \checkmark 1 \text{ mk}$$

3. (a) When a body is wholly or partially inverted in a fluid, it experiences an upthrust force equal to the weight of fluid displaced

- (b) (i) Clockwise moments = anticlockwise moments

$$0.02 \text{ N} \times 0.3 = F \times 0.4$$

$$F = \frac{0.02 \times 0.3}{0.4} = 0.015 \text{ N}$$

$$\text{Upthrust} = \text{weight} - F$$

$$=(90.25 - 0.015)N = 0.235N$$

(ii) Upthrust = weight of liquid displaced  
= 0.235N

$$\text{Mass of liquid} = \frac{\text{weight}}{g}$$

$$= \frac{0.235}{10} = 0.0235\text{kg}$$

$$\text{Vol. of liquid} = \text{vol. of solid} = \frac{\text{mass}}{\text{Density}}$$

$$= \frac{0.025}{200} = 1.25 \times 10^{-4}\text{kgm}^{-3}$$

$$\text{Density of liquid} = \frac{\text{Mass of liquid}}{\text{Vol. of liquid}} = \frac{0.0235}{1.25 \times 10^{-4}}$$

$$= 1880\text{kgm}^{-3}$$

(ii) tension = upthrust - weight  
Weight = mass x gravitational  
= density x volume x gravitational force  
= 0.167 x 1000 x 24 x 10 = 40080  
Tension = 264000 - 40080 = 223920N

4. Needle floats in water due to surface tension. Needle sinks when detergent is added because it reduces surface tension

5. c (ii) Volume under water =  $\frac{3}{4} \times 40,000$   
= 30,000cm<sup>3</sup> ✓

6. (a) (i)  $T = \frac{1}{f} = \frac{1}{100} = 0.01\text{sec}$ ;  
average Vol.  $u = \frac{0.5}{0.01}$ ; = 50cm/s;

(ii) Average Vol.  $V = \frac{2.5}{0.01}$ ; = 250cm/s;

(iii)  $a = \frac{v-u}{t}$  ;  
=  $\frac{250 - 50}{0.01 \times 4}$  ;  
= 5000cm/s<sup>2</sup> ;

(b)  $F = ma$   
= 0.5 x 50 N = 25N;

7. (a) When a body is wholly or partially immersed in a fluid, it experience and upthrust equal to the weight of the fluid displaced;

(b) (i) Shape;  
- Space between 1.0 and 1.1 is larger than that between 1.1 and 1.2


(ii) - Law of floatation which states that floating object displaces its own weight.

(c) Upthrust = Weight of fluid  
= Volume of fluid x density x density x g  
=  $V\rho g$ ;

8. (a) It is the number of times a substance is denser than an equal amount of water

(b) (i) Weight = mass x gravity weight of water displaced

$$p = \frac{M}{V}$$



$$M = \rho \times V$$

$$= (800 \times 12 \times 0.06)$$

$$W = Mg = 576 \text{Kg} \times 10$$

$$= 5760 \text{N}$$

(ii) Upthrust = Weight of liquid displaced

$$= \rho_2 \times V_1 \times g$$

$$= 1.0^3 \times 10^3 \times 0.06 \times 4 \times 10 = 2472 \text{ N}$$

(iii)  $5760 - 2472 = 3288 \text{N}$

c (i)  $(W_2 - W_3)$

(ii)  $R.d = \frac{\text{weight of cork in air}}{\text{weight of equal vol. of water}}$

$$= \frac{W_2 - W_1}{W_2 - W_3}$$

9. (a) A floating body displaces its own weight of the fluid in which it floats  
 (b) The length (x) of block in water increases (block sinks more). Warm water is lighter; hence the blocks must displace more water in order to balance the same weight of the block  
 (c) (i) Upthrust – weight of air displaced

Volume of air = 200  
 Mass of air =  $(200 \times 1.2)$  ✓1  
 Weight of air displaced =  $200 \times 1.2 \times 10$  ✓1  
 $= 2400 \text{N}$  ✓1

(ii) Resultant upward force =  $(2400 - 2200)$  ✓1  
 $= 200 \text{N}$

wt of 1 rat =  $\frac{200}{1000} \times 14 = 2 \text{N}$

$(2 \times n) = 200$  ✓1  
 $n = \frac{200}{2} = 100 \text{ rats}$  ✓1

10. a) When a body is partially or fully/ wholly immersed in a fluid, it experiences an up thrust which is equal to the weight of the fluid displaced ✓1 1 Mk

- b) i) The measurement of weight registered reduces as the brick is lowered into the water ✓  
 Because of increase in up thrust ✓1

ii) Up thrust = weight in air – weight in water (apparent weight)  
 $= (100 - 80) \text{ N}$   
 $= 20 \text{N}$  ✓1

From Archimedes principle

$$20 = V \times S \times g \quad \checkmark 1$$

$$V = \frac{20}{1000 \times 10}$$

$$V = 2 \times 10^{-3} \text{m}^3 \quad \checkmark 1$$

- c) i) To increase sensitivity  
 ii) It displaces more liquid that provides an up thrust to make the hydrometer float  
 iii) To keep the hydrometer upright

11. (a) A floating body displaces its own weight of fluid in which it floats (1mk)

(b) (i)  $p = h\rho g$   
 $= \frac{90}{100} \times 1000 \times 10$   
 $= 9000 \text{Pa or } 900 \text{N/M}^2$



- (ii) – Upthrust force  
 - Weight  
 - tension on the string (for atleast 2 correct)  
 Upthrust = weight + tension on the string

(iii) Upthrust = weight + tension  
 Tension = Upthrust – weight  

$$= \frac{(50 \times 40 \times 20 \times 1000 \times 10)}{1000000} - \frac{(50 \times 40 \times 20 \times 600 \times 10)}{1000000}$$

$$= 400 - 240 = 160N$$

12. Weight of glass = weight of mercury displaced  
 $0.25 \times g = V \times 13.6 \times 10^3 \times g$   
 $V = 0.25$

$$= \frac{13.6 \times 10^3}{1000} \times 0.25$$

$$= 1.838 \times 10^{-5} m^3 (18.4 cm^3)$$

13. a) A floating object displaces its own weight of the fluid in which it falls ✓ 1

b) Up thrust on balloon = weight of air displaced  
 $= mg = Pvg$   
 $= 80m^3 \times 1.2 Kg/m^3 \times 10N/Kg$   
 $= 960N \checkmark 1$

Lifting force = Up thrust – weight of helium  
 $= 960 - (80 \times 0.18 \times 10) \checkmark 1$   
 $= 960 - 144$   
 $= 816N \checkmark 1$

c) i) Mass of water displaced by glass =  $52 - 32 = 20g \checkmark 1$   
 Volume of water displaced = Volume of glass =  $\frac{20g}{1g/cm^3} = 20cm^3 \checkmark 1$

ii) Mass of acid displaced by glass =  $52 - 18 = 34g \checkmark 1$   
 Volume of acid displaced by glass =  $20cm^3 \checkmark 1$   
 Density of acid =  $\frac{34g}{20cm^3} = 1.7g/cm^3 \checkmark 1$

## GAS LAWS


1. (a) It states that the pressure of a fixed mass of gas is inversely proportional to its volume provided temperature is kept constant (PV = K)

(b)  $P_1L_1 = P_2L_2$   
 $86 \times 5 = 75 \times L_2$   
 $= 5.73cm$

(c) (i) When steam condenses, the pressure inside the container will be lower than the atmospheric pressure on the outside. The excess atmospheric pressure acting on the lid exerts a force on the lid thus making it difficult to open the lid.

(ii) The lift pump depends only on atmospheric pressure which can only support a column of water 10m long. The force pump uses force and therefore can lift water to the length greater than 10m.

2. a) Describe how the apparatus may be used to verify pressure law  
 Plotting pressure against absolute temp we get a straight line graph



Conclusion

Pressure of infixed mass of a gas indirectly proportional to its absolute temperature if volume is kept constant

b) i) Given that the relationship between pressure,  $P$  and temperature,  $T$  in Kelvin is of the form

$$P = kT + C$$

Where  $k$  and  $C$  are constants, determine from the graph, values of  $k$  and  $C$

$$K = \text{gradient} \\ = \frac{(8-0) \times 10^4 \text{ Nm}^{-2}}{200 - 0}$$

$$K = 400 \text{ N m}^{-2} \text{ K}^{-1}$$

$$C = 0$$

ii) Why would it be possible for pressure of the gas to be reduced to zero in practice?

- The gas liquefies at low temperature before reaching zero Kelvin

c) A gas is put into a container of fixed volume at a constant volume at a pressure of  $2.1 \times 10^5 \text{ Nm}^{-2}$  and temperature  $27^\circ\text{C}$ . The gas is then heated to a temperature of  $327^\circ\text{C}$ . Determine the new pressure

$$\frac{P_1}{T_1} = \frac{P_2}{T_2} \\ P_1 = 2.1 \times 10^5 \text{ Nm}^{-2} \\ P_2 = ? \\ T_1 = 27 + 273 \\ = 300\text{K}$$

$$T_2 = 273 + 327 \\ = 600\text{K} \\ P_2 = \frac{P_1 T_2}{T_1} \\ = \frac{(2.1 \times 10^5) \times 600}{300} = 4.2 \times 10^5 \text{ Nm}^{-2}$$

3. a) The volume of a fixed mass of gas is inversely proportional to its pressure provided temperature is kept constant.

(b) (i) The bubble expands as it comes up finally bursts when at the surface

$$(ii) p_1 V_1 = p_2 V_2 \\ (76 + 30) \times 3 = (76 + 5) V_2 \\ 106 \times 3 = 81 \times V_2 \\ V_2 = \frac{106 \times 3}{81} \\ = 3.93 \text{ cm}^3$$

(c)  $100^\circ\text{C} - 0^\circ\text{C} = 98 - 11$

$$1 \text{ division} = \frac{87}{100}$$

$$\text{Reading} = \frac{8 \times 56}{1000} \\ = 48.72^\circ\text{C}$$

4. a) The volume of a fixed mass of a gas is inversely proportional to the pressure provided that temperature is kept constant  $\sqrt{1}$

|                                |     |     |     |     |
|--------------------------------|-----|-----|-----|-----|
| $1/v \text{ (mm}^{-3}\text{)}$ | 0.5 | 0.4 | 0.2 | 0.1 |
|--------------------------------|-----|-----|-----|-----|

b) Labeling the axes

Scale

Plotting (3,4) pts

2 points

Below 2 points

Smooth curve / straight line

$$\begin{aligned}
 \text{c) Gradient} &= \frac{\Delta y}{\Delta x} \\
 &= \frac{400 - 160}{(0.5 - 0.2) \times 10^{-9}} \quad \checkmark 1 \\
 &= \frac{340}{0.3 \times 10^{-9}} \\
 &= 1133.33 \times 10^{-9} \\
 &= 1.1333 \times 10^{-6} \text{ KNM}
 \end{aligned}$$

$$\begin{aligned}
 \text{d) } \frac{P_1 V_1}{T_1} &= \frac{P_2 V_2}{T_2} \quad \checkmark 1 \\
 \frac{1 \times 10^5 \times V_1}{285} &= \frac{P_2 \times V_1}{373} \quad \checkmark 1 \\
 P_2 &= 6.95 \times 10^4 \text{ Pa} \quad \checkmark 1
 \end{aligned}$$

5. a) *Boyles Law: States:-*

(i) - *The pressure of a fixed mass of a gas is inversely proportional to its volume, provided the temperature is kept constant. ✓1*

(ii) *Charles Law states:*

- *The volume of a fixed mass of a gas is directly proportional to its absolute temperature at constant pressure. ✓1*

b) (i)

(ii) *at 0°C, v = 4.7 cm<sup>3</sup> ± 0.1 ✓1*

(iii) *Slope =  $\frac{DV}{DT}$*

$$= \frac{(6.4 - 5.0) \text{ cm}^3}{(60 - 10)^\circ\text{C}} \quad \checkmark 1 = 0.028 \text{ cm}^3/^\circ\text{C} \pm 0.002 \quad \checkmark 1$$

(iv) *V = KT + C.*

$$K = \text{Slope} = 0.028 \text{ cm}^3/^\circ\text{C} \pm 0.002 \quad \checkmark 1$$


*C = V intercept when T = 0 and ✓1*

$$= 4.7 \text{ cm}^3 \pm 0.1$$

## SECTION II QUESTIONS

### Pressure


1. In the diagram below, the U-tube contains two liquids; X and Y which do not mix. If the density of liquid Y is  $900\text{Kg m}^{-3}$  and that of X is  $1200\text{Kg m}^{-3}$ , calculate the height of liquid Y



### Current II

✓ 1


1. A battery is rated 120 AH. How long will it work if it steadily supplies a current of 4A.
2. The current capacity of an accumulator is 40Ah. Find the amount of current flowing if the accumulator is used for 600 minutes
3. (a) A student hung a magnet next to a coil of wire to make a door chime as shown in **figure 6:-**



When the switch **S** was put on, the magnet hit the chime bar which made some noise.

- (i) Explain how the current made the magnet move towards the chime bar
- (ii) What should the student do to make the magnet hit the chime bar harder?
- (iii) The student was asked to describe the energy changes inside the device. State the changes:

- (b) A coil of wire is connected in series with a battery, a rheostat and a switch as shown in **figure 7:**




- (i) Draw on the diagram, the shape of the magnetic field inside and outside the coil when the switch is closed

If the jockey **J** on the rheostat is moved towards **Q** what's the effect on:-

- (ii) The resistance of the circuit
- (iii) The current through the coil
- (iv) The magnetic field in the coil

- (i) Explain why a transformer will only transform alternating voltages and not direct current voltage
- (ii) Explain why transformers are widely used throughout the national grid system

4. Determine the current passing through  $L_1$  in the figure shown below, given that 0.8A passes through the battery, 0.28A through  $L_2$  and 0.15A through  $L_3$ .



5. State **two** advantages of generating an alternating current (a.c) to direct current (d.c) in a power station.

**Thermal expansion**


1. Aquatic animals and plants are observed to survive in frozen ponds. Explain this observation

**Light**

1. The length of a pinhole camera is 20cm. Determine the height of a storey building 300m away from the pin hole if the image formed on the screen of the pin hole camera is 2.5cm high

**Electrostatics**

1. (a) An earthed pointed conductor **C** is placed near an insulated conductor **X** charged positively as shown below.




State and explain what happens to charges on x finally.

- (b) A spherical metal sphere is charged positively and brought to contact with the inside surface of a hollow conductor it is then transferred to the cap of the telescope. State and explain what is observed.
- (c) On the axes below sketch a graph of charge against time for charging capacitor.



- (d) State **two** applications of capacitors.
2. The figure below shows an uncharged pith ball under the attraction of a charged ball.




- State and explain what would be observed after the two pith balls touch
3. a) Two metal cons **A** and **B** of different sizes rest on two identical gold leaf electroscope as shown.




Compare the divergence of the gold leaves of the two electroscopes. Explain your answer

- b) Two identical spheres **A** and **B** each standing on an insulated base are in contact. A negatively charged rod is brought near sphere **A** as shown below




- In what way will **A** differ from **B** if separated while the rod is near?
4. In the **figure 1** below, explain what happens when one of the metal balls comes into contact with a negatively charged rod




5. A gold leaf electroscope is positively charged as shown in the diagram in figure 1 where **C** is the

cap and **L** is the gold leaf. State and explain what happens to **L** when a positively charged rod is brought near **C** without touching it.




- You are provided with a charged electroscope, an insulator and a conductor. Describe how you would use these apparatus to distinguish in the insulator from the conductor
- Two identical metal spheres **A** and **B** each standing on an insulating base are in contact. A negatively charged rod is brought near sphere **A** as shown in the figure below.



In what way will sphere **A** differ from **B** if it is separated while the rod is near?

### Measurement II

- The figure below shows a micrometer screw gauge being used to measure the diameter of a rod. The thimble scale has 50 divisions.




### Magnetsim

- (a) Two pins are attached to each of the magnets as shown below.




Explain the behaviour of pins in each case.

- (i) Draw the magnetic field pattern around the magnets below.




Soft iron block  
educationgroup.com

- (ii) Give **one** application of this behaviour of soft iron.
- 2 Give a reason why attraction in magnetism is not regarded as a reliable method of testing for polarity
- 3 (a) The diagram **figure 7** below shows an electromagnet made by a student of Nyamogo Girls secondary school, in the laboratory. The magnet was meant to pick up and release a metal object




- (i) Name giving reasons a suitable material for part **X**
- (ii) The electromagnet will just lift a metal of mass 150g. Taking  $g = 10\text{N/Kg}$ , what will be the least force exerted by the magnet to do this
- (iii) State the changes which the student should make so that a heavier metal object could be lifted by the magnet
- (iv) Explain why the strength of the above magnet cannot be increased indefinitely
- (b) The diagram figure 8 below shows one method of making a magnet. Complete the diagram to make both ends **A** and **B** of the cores be North poles




4. Use the domain theory to differentiate between hard magnetic materials and soft magnetic materials
5. Two similar pins were placed one on a wooden block and the other on an iron block. The two blocks were placed near a magnet. State and explain the observations noted
6. Draw the magnetic field pattern in the figure below and indicate the direction of the force.

**Figure 5**




7. The figure below shows uncharged pith ball under the attraction of a charged ball






State and explain what would be observed after the two pith balls touch


8. The diagram below shows a magnetic field patterns between magnets **S** and **R**. use it to answer questions below:-




- (a) Identify the poles **A** and **B**  
 (b) State which of the two magnets **R** and **S** is stronger. Explain
9. You are provided with a two metal bars; one is magnetized while the other is un-magnetized. Describe briefly how you can identify the two bars without using repulsion method
10. A steel bar can be magnetized and not an aluminum bar. Explain
11. (a) State the **two** laws of electromagnetic induction  
 (b) State **one** way through which energy is lost in a transformer and give a remedy for it.  
 (c) The resistance of a length of power transmitting cable is  $20\Omega$  and is used to transmit  $12\text{KV}$  at a current of  $1\text{A}$ . If the voltage is stepped up to  $18\text{KV}$  by a transformer, determine the power loss. (Assume the transformer is ideal)  
 (d) Using a well- labeled diagram explain how a moving coil meter works  
 (e) Suggest **one** method of decreasing the sensitivity of a moving coil meter
12. The diagram below shows a ferromagnetic material being magnetized by the method shown



On the axes given below, sketch a graph to show how the strength of the magnet being created varies with the number of strokes




13. Arrange the following types of waves in order of increasing frequency:-Ultra-violet radiation, visible light, radio waves and x-rays
14. a) State Lenz's law of electromagnetic induction  
 b) The figure 6 below shows a diagram of a simple electric generator



- i) Label the parts **P** and **Q**
- ii) Identify the polarities of the poles **X** and **Y**
- iii) State **two** ways of increasing the voltage – output in this generator
- c) A transformer supplies a current of 13.5A at a voltage of 48v to a device from a.c. main supply of 240V. Given the transformer is 80% efficient; calculate
  - i) The power supplied to the transformer
  - ii) Current in the primary coil

15. The figure below shows a magnetic material being magnetized




Complete the diagram showing the windings on the magnetic material so as to produce polarities at **A** and **B** both south poles

16. Explain why repulsion method is the surest test for polarity of a magnet as opposed to attraction
17. A nail is electrically magnetized. It attracts an increasing number of iron pins as the magnetizing current increases. After sometime, the nail can no longer attract any more pins. Explain this observation.
18. The table below shows the type of radiation, detection method and uses of electromagnetic radiations. Complete the table:

| Type of radiation | Detection method      | Use              |
|-------------------|-----------------------|------------------|
| Ultraviolet       | Photographic paper    |                  |
|                   | Blackened thermometer | Warmth sensation |
| Radio waves       |                       | Communication    |

19. The figure 3 shows the effect on the magnetic field when two materials **A** and **B** are placed in the magnetic field.



State the difference between A and B.

20. A nail is electrically magnetized. It attracts an increasing number of iron pins as the magnetizing current increases. After sometime, the nail can no longer attract any more pins. Explain this observation.
21. Use the domain theory to explain the process of magnetization

### Reflection at curved surfaces and spherical surfaces

1. (a) The table below shows the object distance  $u$  and the corresponding image distance  $v$  of an object placed in front of a convex lens.

|                                |    |      |    |      |      |      |
|--------------------------------|----|------|----|------|------|------|
| $u$ cm                         | 20 | 25   | 30 | 40   | 50   | 70   |
| $v$ cm                         | 20 | 16.7 | 15 | 13.3 | 12.5 | 11.5 |
| $\frac{1}{u}$ $\text{cm}^{-1}$ |    |      |    |      |      |      |
| $\frac{1}{V}$ $\text{cm}^{-1}$ |    |      |    |      |      |      |


(i) Complete the table by giving your answer to 3 d.p.

(ii) Plot a graph of  $\frac{1}{v}$  (y axis) against  $\frac{1}{u}$


(iii) From the graph, determine the focal length of the lens.

(b) State any **two** differences between human eye and the camera.


2. A battery is rated 120 AH. How long will it work if it steadily supplies a current of 4A.
3.
  - a) Distinguish between mechanical and electromagnetic waves
  - b) What is the relationship between periodic time and the frequency of a progressive wave
  - c) Complete the diagram below to show the shape of the wave fronts after passing the gap



4. Complete the ray diagram to show the position of the image




5. The figure below shows two mirrors inclined at an angle of  $30^\circ$  to each other. A ray of light is incident on one mirror as shown



Sketch the path of the ray to show its reflection on the two mirrors

6. What is meant by the term **spherical aberration**?
7. Complete the ray diagram below by showing the position of the image




State **one** application of the set up above

8. A plain sheet of paper and a plane mirror both reflect light yet only the plane mirror forms images. Explain why the paper cannot form images.
9. Give **one** advantage and **one** disadvantage of using a convex mirror as a driving mirror
10. The table below shows the image distance **V** and the corresponding magnification, **M** for an object placed in front of a concave mirror.

|                             |     |     |     |     |     |     |     |
|-----------------------------|-----|-----|-----|-----|-----|-----|-----|
| <b>Magnification M</b>      | 0.5 | 1.0 | 2.0 | 3.0 | 4.0 | 5.0 | 6.0 |
| <b>Image distance V(cm)</b> | 15  | 20  | 30  | 40  | 50  | 60  | 70  |

- (a) Plot a graph of **V (y-axis)** against the magnification **M**
- (b) From the graph, determine the focal length of the mirror
- (c) Given the image **IM**, locate the position of the object **OB**. Use arrows to show how the image is formed on a concave mirror



11. State the difference between a virtual image formed by a plane mirror and that formed by a concave mirror
12. Figure 1 shows an object O being viewed using two inclined mirrors M<sub>1</sub> and M<sub>2</sub>. Complete the diagram by sketching rays to show the position of the image as seen by the eye -E





Fig 1.

13. The figure below shows an object placed in front of a plane mirror. Draw appropriate rays to locate the image as seen by the observer.




14. Define the following terms as used in curved mirrors:-
  - (i) Principal focus (F)
  - (ii) Focal length (f)
- (b) By use of a ray diagram, show how a concave mirror may be a dentist mirror
- (c) An object is placed 12cm from a convex mirror of radius of curvature 20cm. Calculate the position of the image
- (d) (i) A lady holds a large concave mirror of focal length 1.8m from her face. State **two** characteristics of her image in the mirror
- (ii) A boy is standing between two cliffs **A** and **B** but nearer to cliff **A** than **B**. He stands 160m from wall **BA** and shouts once. He hears two echoes and discovers that the time between the two echoes is 0.8 seconds. Determine how far the boy is standing from cliff **B** given that the speed of sound in air is 340m/s

### Linear motion

1. A footballer kicks a ball of mass 0.6kg initially at rest using a force of 720N. If the foot was in contact with the ball 0.1seconds, what was the take off speed of the ball?
2. A car starting from rest accelerated uniformly for 5minutes to reach 30m/s. it continues at this

speed for the next 20 minutes and then decelerates uniformly to come to a stop in 10 minutes. On the axes provided, sketch the graph of the velocity against time for the motion of the car.



### Machines & inclined planes

1. (a) Distinguish energy from work
- (b) A lady uses a ramp to lift 1500N load through a vertical distance of 10m. The ramp makes an angle  $30^\circ$  to the horizontal. If the efficiency of the ramp is 75%:
  - (i) Calculate the V.R of the machine
  - (ii) Find the mechanical advantage
  - (iii) Find the effort needed to move the load up the ramp at constant velocity


✓ 1

### Resistors

1. You are provided with the following apparatus: connecting wires, an ammeter, fixed resistors, a voltmeter, a variable resistor, a switch and two dry cells in a cell holder. Draw a circuit that can be used using the apparatus above to verify Ohm's law 2. (a) Three resistors **A**, **B** and **C** where **A** is resistor  $200\ \Omega$  **C** of resistance  $100\ \Omega$  and **B** is of unknown resistance are connected in parallel. This arrangement is then placed in a circuit and current passing through and potential difference across it measured. The table below shows the result.


|           |      |      |      |      |      |
|-----------|------|------|------|------|------|
| p.d.v.    | 3.0  | 6.0  | 9.0  | 12.0 | 15.0 |
| Current A | 0.15 | 0.30 | 0.45 | 0.60 | 0.75 |

- (i) Plot a graph of p.d. against current A
  - (ii) From the graph calculate the total resistance of the resistors.
  - (iii) What is the value of the unknown resistor?
- (b) The ammeter in the figure below shows a reading of 2A. What is the reading of the voltmeter v.?




3. a) Three resistors of resistance  $2.0\ \Omega$  and  $6.0\ \Omega$  are connected together in a circuit. Draw a circuit diagram to show the arrangement of the resistors which gives
  - i) Effective resistance of  $3.0\ \Omega$

- ii) Minimum resistance  
 b) In figure 9 the voltmeter reads 2.1V when the switch is closed, the voltmeter reads 1.8v and the ammeter 0.1 A




**Determine:**

- i) The e.m.f of the cell
  - ii) The internal resistance of the cell
  - iii) The resistance of the of the lamp
- c) One reason why the earth pin is longer than the neutral and the live pins is to open the shutters of the socket, state **one** other reason why it is longer  
 d) Why are lamps in a house always connected in parallel but not in series?
4. State **two** other factors, which would affect the resistance of a metal conductor other than the temperature  
 b) The ammeter in the circuit below has negligible internal resistance. The cell has internal resistance of  $0.5\Omega$  and an electromotive force of  $3.0V$




Determine the value of current the ammeter registers when switch S is closed

5. a) State **Ohms law**
6. In the configuration of resistors given below, determine the current through the  $5\Omega$  resistor




7. **Figure 5** is a circuit diagram of three resistors connected to a  $6V$  battery




**Figure 5**

- Determine the potential difference across the  $3\Omega$  resistor  
 8. (a) Study the circuit diagram shown below.


Figure 7



- (i) Determine the reading of the voltmeter V.
  - (ii) Determine the reading of the ammeter A.
  - (iii) Explain the effect on the reading of the voltmeter if the  $3\Omega$  resistor is altered to be  $6\Omega$
- (b) A transformer is designated to work from a 240V a.c. mains and to give a supply of 8V to ring house bells. The primary has 4800 turns.
- (i) What type of transformer is this? Give a reason.
  - (ii) Why is the iron core laminated?
  - (iii) Calculate the secondary turns if the efficiency is 100%.
9. The diagram in the figure below shows a wheat stone bridge




- K, L, M and N** are four resistors joined as shown. The value of resistance of resistor **K** is unknown.  $I_1, I_2, I_3$  and  $I_4$  are the amount of current passing through **K, M, L** and **N** respectively. It is also provided that **L** is a variable resistor.
- (i) Explain how the set-up is used to determine the value of unknown resistance of **K**
  - (ii) State why wheat stone bridge is more accurate in measuring resistance than the voltmeter-ammeter method
- (b) In an experiment to determine the resistance of a nichrome wire using the metre bridge, the balance point was found to be at 38cm mark. If the value of the resistance in the right hand gap needed to balance the bridge was  $25\Omega$ , Calculate the value of resistance of nichrome wire
10. Four capacitors of capacitance,  $3\mu\text{f}$ ,  $4\mu\text{f}$ ,  $5\mu\text{f}$  and  $3\mu\text{f}$  are arranged as shown below. Find the




11. In the circuit diagram below, X is a fixed resistor while Y can be varied between 0 and  $100\Omega$  using a sliding jockey






Calculate:

- (i) The minimum possible current in the circuit (2mks)
- (ii) The maximum possible current in the circuit (2mks)
- (c) The following figure shows an electric circuit in which five resistors are connected to a battery of e.m.f 4.0V and negligible internal resistance



**Determine:**


- (i) The total resistance of the circuit
  - (ii) The potential difference between **Y** and **Q**
  - (d) Explain **two** factors that affect the resistance of a metallic conductor
12. Three identical cells of e.m.f. 2.0v and of negligible internal resistance are connected as shown in figure below. Determine the ammeter reading.



### Refraction of light


1. The refractive index of paraffin is 1.47 and that of glass is 1.55. Determine the critical angle of a ray of light travelling from glass to paraffin
2. The diagram **figure 1** below shows a ray of light incident on glass air boundary:

fig. 1



A second ray strikes the boundary at the same point C at an angle of incident greater than  $a^\circ$ .

- (i) On the diagram, draw the second ray before and after striking the boundary
- State Snell's law
  - When does total internal reflection occur?
  - The figure below represents a ray of light falling normally on the curved surface of a semi-circular glass block A at an angle of  $32^\circ$  at O and emerging into air at an angle of  $48^\circ$



Calculate the absolute refractive index of the glass of which the block is made. (Assume air is a vacuum)

4. **Figure 2** below shows a ray of light traveling from glass to water





figure 2


figure 2

Calculate the angle  $\theta$  if the refractive index of glass and water are  $\frac{3}{2}$  and  $\frac{4}{3}$  respectively (3mks)


5. Figure 3 shows light rays moving from medium 1 to medium 2. If the refractive index of medium 1 is  $\frac{4}{3}$  and that of medium 2 is  $\frac{3}{2}$ . Calculate angle  $r$



6. (a) The diagram below shows a glass prism and an incident ray striking the face marked AB. The critical angle of the glass is  $42^\circ$ . Use it to answer the questions that follow:-



- (i) Complete the diagram showing the path of the emergent ray  
 (ii) Calculate the angle of refraction of the resultant emergent ray
7. (a) (i) What is a critical angle as used in refraction of light?  
 (ii) State **one** condition under which total internal reflection occurs  
 (b) Calculate the value of the critical angle  $c$  in the figure below



- (c) (i) Show that  $m = \frac{v}{f} + 1$


where  $m$  = linear magnification,  $V$  = Image distance and  $f$  is the focal length of lens

- (ii) In the table below shows readings obtained out of an experiment to determine focal length of a converging lens

|                         |      |      |    |    |    |
|-------------------------|------|------|----|----|----|
| Image distance $V$ (cm) | 17.1 | 18.3 | 20 | 23 | 30 |
| Object distance $u$     | 40   | 35   | 30 | 25 | 20 |

Plot a graph of  $\frac{1}{V}$  against  $\frac{1}{u}$  and determine the focal length of the lens from the graph. (Use the graph paper provided).


8. a) The Fig.9 shows a ray of sunlight incident to face AB of a glass prism. ••  
 Fig. 9



- i) Complete the diagram showing the observation on the screen.
  - ii) Explain the observation on the screen.
  - iii) State why the spectrum formed above is not pure.
- b) i) You are provided with four equilateral prisms and four convex lenses. Sketch a diagram showing how all the eight can be arranged to make a simple prism binoculars.
- ii) State **one** reason why prisms produce better optical instruments than plane mirrors.

## Sound II


1. The human ear can distinguish two sounds as separate only if they need it at least 0.1 seconds apart. How far from a wall must an observer be in order to hear an echo when he shouts. (Speed of sound = 330m/s)
2. A girl standing 220m from the foot of a high wall claps her hands and the echo reaches her 1.29 seconds later. Calculate the velocity of sound in air using this observation
3. A boy standing in front of a cliff blows a whistle and hears the echo after 0.5 seconds. He then moves 17m further away from the cliff and blows the whistle again. He now hears the echo after 0.6seconds. Determine the speed of the sound
4. Sound tends to travel over longer distance at night. Explain
5. You are given two tubes  $T_1$  and  $T_2$ , a clock and a hard wall. Explain how you can use the apparatus to demonstrate reflection of sound



6. State **two** conditions necessary for total internal reflection to occur
7. A student carrying out an experiment discovered that it took 2 seconds for sound wave traveling through a telephone line to cover a distance  $d$  metres and 20 seconds for the same sound traveling through air to cover a similar distance. Determine the ratio of the speed of sound in air to that in the wire.
8. State **one** factor that affects the velocity of sound in air


## Thin lenses

1. The figure below shows how rays from a distant and near objects are focused inside a human eye with a certain defect



Name the defect and state **two** causes of the defect


2. (a) The figure below shows an object **O** placed in front of an objective lens  $L_o$  whose focal length  $f_o$  is less than  $f_e$ , the focal length of the eyepiece  $L_e$ . Complete using ray construction how the arrangement would produce a compound microscope




- (b) A nail is placed 25cm from the objective lens of focal length 15cm. On the other side of the objective lens another converging lens of focal length 30cm is placed as the eyepiece. The distance between the two lenses is 52.5cm

**Find:** (i) the position of the first image  
(ii) the position of the final image from the eye piece lens

3. (a) The figure below shows a set-up consisting of a mounted lens,  $L_1$ , a screen **S**, a metre rule and a candle




- (i) Describe how the set-up can be used to determine the focal length,  $f$  of the lens.  
(ii) Explain why the set-up would not work if the lens was replaced with a diverging lens
- (b) The graph in the diagram in figure below shows the relationship between  $\frac{1}{u}$  and  $\frac{1}{v}$  for a converging lens where  $u$  and  $v$  are the object and image distances respectively. From the graph, determine the focal length  $f$  of the lens




- (c) An object placed 15cm from a convex lens forms an image twice the size of the object.  
Determine the focal length of the lens

4. The graph below represents a graph of stopping potential  $V_s$ , V against frequency  $f$ , Hz



- (a) Use the graph to determine:
- The threshold frequency of the metal
  - Plank's constant
  - Work function of the metal


(b) **Figure 8** below shows a mercury vapour lamp, which emits ultraviolet light held over a negatively charged electroscope:



- (i) What happens to the leaf after the lamp is switched on?
  - (ii) Explain why it happens
  - (iii) If the experiment is repeated with equally bright red light held the same distance from the plate in place of the mercury vapour lamp, what effect would this have on the leaf?  
Give a reason
  - (iv) What does photoelectric effect suggest about the nature of light?
- 5.
- (a) Describe briefly a simple method of estimating the focal length of a convex lens.
  - (b) Define linear magnification of a lens.
  - (c) In an experiment to determine the focal length of a converging lens, the following readings were obtained

|   |      |      |      |      |      |
|---|------|------|------|------|------|
| <b>Image distance <math>V</math> cm</b> | 14.3 | 16.0 | 17.7 | 21.0 | 31.0 |
| <b>Magnification <math>m</math></b>     | 0.4  | 0.60 | 0.80 | 1.10 | 2.10 |

- (i) Plot a graph of  $m$  against  $V$ .
  - (ii) From the graph determine the focal length of the lens.
  - (d) Which eye defect is corrected by a diverging lens? Show using a diagram how this is achieved
- 6.
- a) Describe with the aid of a labeled diagram an experiment to determine the focal length of the lens when provided with the following;
    - An illuminated object screen
    - A convex lens
    - A lens holder
    - A plane mirror
    - A meter rule
  - b) A small vertical object is placed 28cm in front of a convex lens of focal length 12cm. In the space below, draw a ray diagram to locate the image and find its magnification.  
(use a scale: 1cm represents 4cm)
  - c) The figure below shows a human eye with a certain defect




- i) Name the defect
- ii) On the same diagram, sketch the appropriate lens to correct the defect and sketch rays



to show the effect of the lens

7. An object of height 10cm is placed in front of a diverging lens of focal length 25 cm and at a distance of 20 cm from the lens. Calculate the height of the image formed
8. (a) The figure below shows an object, **O**, placed in front of an objective lens **L<sub>o</sub>** whose focal length, **f<sub>o</sub>** is less than the focal length of the eye piece lens; **L<sub>e</sub>** Complete using ray construction how the arrangement would produce a compound. Microscope



- (b) A thin converging lens of focal length 30cm is used to form a real image on a screen 90cm from the lens, Determine :-
  - (i) The object distance
  - (ii) The magnification
9. Figure 2 shows an object **O** placed in front of a concave ion with principal foci **F** and **F<sup>1</sup>**. Construct a ray diagram to locate the position of the image




Fig 2


10. Use a ray diagram to show how short sightedness in a human eye can be corrected.
11.
  - a) An object is placed 15 centimeters in front of a diverging lens of focal length 20 cm. Use a ray diagram to determine the image distance and its magnification.
  - b) A nuclide **F** has a half life of 5 hours. What percentage of the original number of atoms of the isotope would have decayed after 30 hours?
  - c) A current of 1.5A flows through a conductor in 5 seconds. Determine the number of electrons that pass through the conductor (charge on an electron =  $1.6 \times 10^{-19} \text{ C}$ )
12. Calculate the wavelength of the KBC f.m radio waves transmitted at a frequency of 95.6 mega Hertz

## Quality of heat


- 500g of a metal is heated to 100°C and then placed in a 200g mass of water at 15°C. If the final temperature rises to 21°C, calculate the specific heat capacity of the metal.  
(Specific heat capacity of water = 4200J<sup>-1</sup>kg<sup>-1</sup>)

## Waves II

- Distinguish between stationary and progressive waves
    - State a reason why a closed tube or pipe produces less quality sound than an open one
  - The figure below shows a piston inside a glass tube




- Air inside is made to vibrate producing a fundamental note. Find the fundamental frequency to be produced
  - If the glass tube is made open by removing the piston and opening the other end, what is the new fundamental frequency?
- The figure below shows circular waves approaching a straight reflector. Complete the sketch to show what happens when the waves hit the reflector.




State **one** difference between the way sound waves and electromagnetic waves are transmitted.


- A mine worker stands between two vertical cliffs 400 m from the nearest cliff. The cliffs are **X** distance apart. Every time he strikes the rock once he hears two echoes, the first one comes after 2.5 sec. while the 2<sup>nd</sup> follows 2sec. later. From this information; Calculate;
  - Speed of sound in air.
  - The value of **X**.
- The figure below shows waves starting from two coherent sources  $S_1$  and  $S_2$ .




- What would be observed at **P** if the waves are  
 (i) light waves. (ii) Sound waves.  
 (c) State the conditions for diffraction of light to occur. *✓1 mk*
4. The diagram below represents plane wave fronts produced in a ripple tank.




- Given that the distance **AB** is 15 cm, determine the wavelength of the wave
5. (a) The figure below shows a wave profile with velocity 340m/s



- Determine:** (i) The frequency of the wave  
 (ii) The period of the wave
- (b) On the same diagram in (b) above sketch a wave profile of another wave with same frequency, greater amplitude but  $180^\circ$  out of phase with the one in (b)
6. Figure 3 shows the displacement of a particle in a progressive wave incident on a boundary between deep and shallow region




- Complete the diagram to show what is observed after bounding. (Assume no loss of energy)
7. (a) Explain the difference between progressive waves and stationary waves  
 (b) State any **two** conditions necessary for the establishment of a stationary wave  
 (c) (i) The figure below shows two loudspeakers  $L_1$  and  $L_2$  connected to a signal generator




- An observer walks along the line  $0-01$  (equidistant from  $L_1$  and  $L_2$ ) and another along the Line  $AA_1$ . Explain the observation made by each and give reasons to your answer
- (d) If a wave is propagated at a velocity of  $50\text{m/s}$ ; determine its frequency if the distance travelled by the wavelength in 2 cycles is  $1.25\text{m}$   
 (e) State **one** condition necessary for interference to occur


8. Give conditions necessary for diffraction of waves to occur  
 9. Distinguish between diffraction and refraction of waves  
 10.




11. Determine the frequency of the wave  
 The figure below shows a wave profile.




12. The following shows a diagram of displacement against time curve of a standing wave reflected its own path




13. A student touches the surface of water in a big pan at point A at regular intervals and observes the ripples as in the diagram below. Explain the pattern of the ripples observed.



14. In the figure below shows a series of plane waves approaching a gap. Complete the diagram to show the wire after passing through the gap if




15. (a) (i) State the difference between mechanical and electromagnetic waves  
(ii) Give **one** example of each of the above waves  
(b) The figure below shows water waves crossing a boundary between deep and shallow water




Complete the diagram to show the wave in the deep region

- (c) (i) Complete the diagram below to show the shape of the wave fronts after passing through the deep gap



- (ii) What is the relationship between periodic time and frequency of a progressive wave?  
(iii) Explain why radio waves reception is better than T.V reception in mountainous regions

16. a) i) Distinguish between stationary waves and progressive waves. In terms of their propagation.  
 ii) State a reason why a closed pipe produces less quality sound than an open pipe.  
 b) The Fig. 10 represents an oscillation taking place at a particular point while a sound wave in a gas passes the point. The vertical axis is labeled displacement.



- i) Explain what is meant by displacement in this context.  
 ii) From the figure determine: I. The period.  
 II. The frequency
- c) Calculate the wavelength of the sound wave in the figure. Take the velocity of sound in the gas to be 34m/s  
 d) State **two** factors that can. increase the speed of sound in. solids

### Work, energy and power


- Given that a lamp is rated 45W 240V. Calculate the resistance of the heating element.
- An electric bulb is rated 40W, 240V. What is the resistance of its filament?
- An electrical immersion heater is rated 3kW, 250V. Choose a suitable fuse from 3A, 5A, 10A, 12A, and 20A that can be used in such an appliance.
- An electric kettle is rated 3KW, 250V. Determine the resistance of the coil
- An electric kettle rated 3.0Kw, 240V is filled with water. If the water boiled after 8 minutes of heating, determine the energy used in boiling the water.
- (a) An electrical heater is rated 3.45KW. The heater is immersed in 2.4kg of water. Calculate the minimum time it takes for the temperature of the water to rise from 23.0°C to 69.0°C. (*Specific heat capacity of water = 4.2Jg<sup>-1</sup>K<sup>-1</sup>*)

### Floating and sinking


- Define thermionic emission
- (a) State:- (i) Archimedes's principle  
 (ii) The law of floatation  
 (b) The solid of mass 12kg, weighs 100N and 94N when fully immersed in water and liquid **L** respectively. Calculate:-  
 (i) The density of the liquid in S.I units  
 (ii) The density of the solid

## Photoelectric effect

1. (a) Define the following:
  - (i) Photoelectric effect
  - (ii) threshold wavelength
- (b) The variation of frequency  $f$  with the maximum kinetic energy  $E_k$  of the emitted electrons is shown on the graph below:



- (ii) the value of the Planck's constant  $h$
  - (iii) the work function,  $W_0$
  - (c) On the same graph in (b) above, draw a line to show the variation of frequency,  $f$ , with the maximum kinetic energy,  $E_k$ , of the emitted electrons from a second metal which has a lower work function than that used in (b)
2. **Figure 8** below shows a mercury vapour lamp, which emits ultraviolet light held over a negatively charged electroscope:




- (i) What happens to the leaf after the lamp is switched on?
- (ii) Explain why it happens
- (iii) If the experiment is repeated with equally bright red light held the same distance from the plate in place of the mercury vapour lamp, what effect would this have on the leaf?  
Give a reason

- (iv) What does photoelectric effect suggest about the nature of light?
3. Calculate the wavelength of Green light whose energy is  $3.37 \times 10^{-19} \text{ J}$ .  
 ( $h = 6.63 \times 10^{-34} \text{ JS}$ ,  $C = 3.0 \times 10^8 \text{ m/s}$ )
4. a) Define the term *work function*  
 b) Name **one** factor that determines the velocity of photoelectrons produced on a metal surface when light shine on it  
 c) In a photoelectric effect experiment, a certain surface was illuminated with radiations of different wavelengths and stopping potential determined for each wavelength. The table below shows the results obtained.

|   |      |      |      |      |      |
|---|------|------|------|------|------|
| Stopping potential, $V_s$                   | 1.35 | 1.15 | 0.93 | 0.62 | 0.36 |
| Wave length, ( $\times 10^{-7} \text{ m}$ ) | 3.77 | 4.04 | 4.36 | 4.92 | 5.46 |

- i) On the grid provided plot a graph of stopping potential (Y –axis) against frequency  
 ii) From your graph determine:  
 a) The threshold frequency  
 b) The plank's constant,  $h$   
 ( $e = 1.6 \times 10^{-19} \text{ Coulomb}$ ,  $C = 3.0 \times 10^8 \text{ m/s}$ )
5. a) State the role of the Grid in a cathode ray tube  
 b) Explain why a magnetic field is used in the TV deflection system instead of an electric field  
 c) The time base of a CRO is 25ms/div while its gain is 2.5V/div. Use this information to answer the questions that follow:  
 i) Calculate the frequency of the signal  
 ii) What is the peak voltage of the signal
6. The graph below shows the relation between the stopping potential,  $V_s$  and the frequency of radiation when a certain surface is illuminated with light of different frequencies




From the graph determine:-


- (i) The threshold frequency  
 (ii) The value of plank's constant ( $e = 1.6 \times 10^{-19} \text{ C}$ )  
 (III) The work function of the material



- 7 a) State **one** reason why a C.R.O is a more accurate voltmeter than a moving coil voltmeter  
 (b)The diagram below represents a cathode ray oscilloscope (CRO)




- i) Name the parts labeled **A** and **B**  
 ii) What are the functions of **C** and **D**?  
 iii) State how electrons are produced
8. a) What is meant by the term photo electric effect  
 b) In an experiment using a photo cell, ultra violet light of varying frequency strikes a metal surface. The maximum Kinetic energy ( $KE_{max}$ ) of the frequency  $F$  is measured. The graph below shows how the maximum kinetic energy varies with frequency  $F$




Use the graph to determine:-

- i) Threshold frequency **F**  
 ii) The plank's constant, **h**  
 iii) Work function of the metal

9. (a) The diagram fig 9 below shows a photo cell; connected in a circuit:-  
fig. 9



- (i) Complete the diagram by indicating the correct polarities in the gap for current to flow in the circuit
  - (ii) State and explain the effect of using light of different wave lengths on the amount of current flowing in the circuit given that the distance of the source of light remains the same
- (b) Two fixed resistors one of  $100\Omega$  and the other of unknown resistance are connected in parallel. The combination is placed in a circuit and current passing through the combination was measured for various p.d. The graph in figure 10 below drawn to scale shows the results:-



- (i) From the graph, calculate the total resistance of the combination
  - (ii) Determine the value of the unknown resistance
- (c) (i) Explain the cause of eddy currents and how they are minimized in a transformer
- (ii) A transformer with 4200 turns in the primary coil operates a 240V mains supply and gives an output of 8.0V. Determine the number of turns in the secondary coil (assuming it is 10% efficient)


10. State **one** factor that affects photoelectric effect
11. a) i) What is photoelectric effect?  
 ii) You are provided with the following; a photo cell; a source of UV light, a rheostat, a source of e.m.f, a millimeter, a voltmeter and connecting wires. Draw a circuit diagram to show how photoelectric effect may be demonstrated in the laboratory
- b) In a photoelectric effect experiment, a certain surface was illuminated with radiation of different frequencies and stopping potential determined for each frequency. The following results were obtained:

|                                      |      |      |      |      |      |
|--------------------------------------|------|------|------|------|------|
| Frequency (f) ( $\times 10^{14}$ Hz) | 7.95 | 7.41 | 6.88 | 6.10 | 5.49 |
| Stopping potential, ( $V_s$ ), (V)   | 1.35 | 1.15 | 0.93 | 0.62 | 0.36 |


- i) Plot a graph of stopping potential (Y-axis) against frequency
- ii) Determine plank's constant, h and the work function of the surface given that  $eV_s = hf - hf_0$ , where  $hf_0 = Q_e = 1.6 \times 10^{-19} \text{ C}$
- c) A surface whose work function  $Q = 6.4 \times 10^{-19} \text{ J}$  is illuminated with light of frequency  $3.0 \times 10^{15} \text{ Hz}$ . Find the minimum K.E of the emitted photo electrons (use value of h obtained in **b(ii)** above)

### ELECTRICITY & Electronics


1. State **one** advantage of a lead-acid accumulator over a dry cell
2. State **one** defect of a simple cell and explain how it can be corrected.
3. Study the circuit below:




- Determine the current flowing in the circuit
4. When the time base of a cathode ray oscilloscope is turned on, there is a horizontal trace across the screen as shown in the figure:-




- (i) An alternating potential difference of constant frequency and constant amplitude is then connected to the Y-input of the oscilloscope. Sketch on the same diagram above the trace which might be obtained
- (ii) The time base is switched off but the alternating potential difference is left connected. Describe what would be observed on the screen
5. The figure below shows the wiring in a modern mains appliance




- Identify the wires **Y** and **Z**
6. State **two** ways of decreasing capacitance
7. (a) The figure below represents part of an electric cooker coil.




- (i) State why the part labeled **W** is coiled
- (ii) State the property of material **X** that makes it suitable for its use
- (b) State the advantage of transmitting power at:-
- (i) Very high voltage
- (ii) Alternating voltage
- (c) Aluminium wires are commonly used in power transmission than copper wires. Give **two** advantages of aluminum as transmission lines
- (d) The diagram below shows a wrongly wired three pin plug. Black




- (i) Indicate in the diagram above the correct colors for the wiring
- (ii) State the use of device marked **X**
- (e) A household uses a 1.5Kw water heater for 2 hours a day for 30 days. If the cost of electricity is shs.6.70 per Kwh, how much will they pay for this consumption?
8. The diagram below shows a simple cell:-



- (a) (i) Name **z** and solution **y**
- (ii) Name and explain the defect that occurs at plate **z**
- (iii) Give **one** method of preventing the defect that occurs at the copper plate
- (b) (i) Explain how P-type semi-conductor is formed
- (ii) The figure below shows a circuit diagram for full wave rectification




- (I) Draw the diodes  $D_3$  and  $D_4$  on the diagram to complete the circuit  
 (II) On the axes below sketch a voltage –time graph observed when a C.R.O is connected at points




- (iii) On the circuit diagram (b) (ii) above, draw a capacitor which can be used to smoothen the output voltage

9. Explain how conductivity of a semi conductor changes with increase in temperature  
 10. With the time base switched on; the following trace was obtained on the screen of a CRO as shown in the figure below:



Draw a circuit diagram that can be used to produce the wave above


11. **Figure 4** below shows a circle with two diodes **P** and **Q** and a cell:-




**Figure 4**

Explain the observation which would be made if **S** is closed

12. Explain why eight 1.5V cells arranged in series to give a total of 12V cannot be used to start a car. But car battery of 12V starts a car
13. a) i) Distinguish between a **p-type** and an **n-type** extrinsic semi conductors  
 ii) The figure below shows a bridge rectifier




A capacitor has been connected across the resistors as shown. Sketch on the axes below the wave form when a C-R-O is connected across the resistor; R




Sketch on the same axes above the wave form when a C-R-O is connected across the resistor R and capacitor c removed


iii) Figure shows a voltmeter connected across the cell. The voltmeter reads 1.5V when the switch S, is open and 1.25V when the switch is closed.



- i) What is the e.m.f of the cell?
- ii) What is the terminal voltage of the cell?
- iii) Calculate the internal resistance of the cell
14. What is the use of a fuse in an electric circuit?
15. Distinguish between **Topping** and **Dopping**
16. The figure below shows the set up for a simple cell.




- a) Name the Electrode **A** and the solution **B**
- b) State **two** reasons why the bulb goes off after a short time
17. The figure 2 shows a simple cell made of copper and zinc electrodes dipped in dilute sulphuric acid




- a) Identify the cathode  
 b) If a voltmeter is connected across the rods the reading is observed to reduce with time.  
 State **two** causes of this observation
18. State **one** reason why colour televisions have a higher power rating than black and white televisions
19. Explain **two** factors that affect the capacitance of a parallel-plate capacitor
20. a) A girl opened up a used up dry cell and found the following:  
 i) The zinc casing was 'eaten away'  
 ii) The cell was watery  
 Name the cell defect
- b) Three identical bulbs are connected in series with a battery of dry cells. At first the bulbs shine brightly but gradually become dimmer. Using the same cells, explain how you would increase the brilliance of the bulbs
21. Figure 9 below shows a diagram of an n – p – n transistor.  
 (a) Complete the diagram by showing the connections of two batteries **suitable for biasing** the transistor in the common- emitter mode.

Figure 9




22. State the **purpose** of introducing an impurity in a semi conductor.
23. In an attempt to establish the relationship between current through a junction diode and the p.d across it, a student connected a diode to an e.m.f source as in **figure 3** below:-




- (a) State whether the diode is forward biased or reverse biased  
 (b) Briefly describe how she obtained her readings  
 (c) Sketch a graph to represent the relationship between current (y-axis) and the p.d across the diode

24. Figure 8 shows a circuit where a battery of emf 4.5V, switches A and B, two capacitors  $C_1 = 0.3 \mu\text{F}$  and  $C_2 = 0.5 \mu\text{F}$  and a voltmeter are connected




- a) Determine the charge on  $C_1$  when switch A is closed and switch B is open  
 b) What is the effective capacitance  $C_T$  when both switches A and B are closed?  
 c) State what is observed on the voltmeter when;  
 i) Switch A is closed and switch B is open  
 ii) Switch A is closed and opened and then B is closed  
 iii) Explain the observation made in **c(ii)** above
25. (a) Define capacitance  
 (b) Two aluminium plates **A** and **B** of same dimensions are each mounted on an insulating stand. Plate **A** is charged to high voltage and connected to uncharged electroscope while plate **B** is earthed. The two plates are placed side by side as in the diagram *figure 4* below:-



- (i) Indicate on the diagram the position of the leaf and charge distribution on the electroscope  
 (ii) State and explain the observation on the electroscope when the distance ( $x$ ) of separation between the plates is increased while keeping the area of overlap the same  
 (c) A  $12 \mu\text{F}$  capacitor is charged with a 200V source then placed in parallel with uncharged  $8.0 \mu\text{F}$  capacitor as shown in fig 5 below:-

*fig. 5*




Determine:

- (i) The initial charge on the  $12 \mu\text{F}$  capacitor  
 (ii) The final charge on each capacitor



(d) The diagram **figure 6** below shows a pear shaped charged conductor on an insulating stand (charges not shown on the diagram)


**fig. 6**



Part **A** is touched using a proof-plane and then the proof-plane is brought next but not touching the cap of a leaf electroscope (not shown on the diagram). The same experiment is repeated for part **C** of the conductor.


- (i) State the expected observation in the above experiments
- (ii) Explain the observations made in **(d) (i)** above
- (iii) Name any **one** application of the above phenomenon

26. a) State **two** factors that affect the capacitance of a parallel plate capacitor  
 b) The diagram below shows an arrangement of capacitors in a circuit




- i) Determine the total charge in the circuit

27. a) What is doping as used in electronics  
 b) Distinguish between intrinsic and extrinsic semi-conductors.  
 c) What would be observed in the diagram below when switch **S** is closed, **B<sub>1</sub>** and **B<sub>2</sub>** are identical torch bulbs




28. a) Define Eddy currents  
 b) The diagram below shows the north pole of a magnet approaching a solenoid




- i) Using Lenz's law, indicate the direction of current through the galvanometer  
 ii) Explain the observation made when:

- I The magnet is moved away from the solenoid  
 II The magnet is placed stationary in the solenoid

- c) A transformer is designed as shown in the figure below. If the primary coil has 2400 turns and the secondary has 200 turns calculate the p.d across BC assuming there are no energy losses in the transformer




- d) The figure shows a cross-section of a bicycle dynamo. The wheel is connected by an axle to a permanent cylindrical magnet and is rotated by the bicycle tyre



- i) Explain why the bulb lights  
 ii) How can the bulb be made brighter
29. A car battery requires topping up with distilled water occasionally. Explain why this is necessary and why distilled water is used
30. Draw appropriate symbol of a circuit diagram of a junction diode in reverse bias


31. a) In the circuit diagram shown in Fig.5 each cell has an e.m.f of 1.5v and internal resistance of  $0.5\Omega$ . The capacitance of each capacitor is  $1.4\mu\text{F}$ .

Fig.5



- i) When the switch S is closed determine the ammeter reading.  
 ii) When the switch S is closed determine the charge on each capacitor.  
 b) The diagram in Fig. 6 represents two parallel plates of a capacitor separated by a distance  $d$ . Each plate has an area of a square unit. Suggest **two** adjustments that can be made so as to increase the effective capacitance.


Fig.6



- c) Complete the table to describe the function of the parts of a lightning conductor.

| Part                | Function |
|---------------------|----------|
| Spike               |          |
| Thick copper rod    |          |
| Earthed metal plate |          |


32. The circuits in Fig. 7 shown are close to each other.



- a) When the switch is closed, the galvanometer shows a reading and then returns to zero. When the switch is then opened, the galvanometer shows a reading in the opposite direction and then returns to zero. Explain these observations.  
 b) Energy losses in a transformer are reduced by having a laminated soft iron core. State and explain **two** other ways of reducing energy losses in a transformer.

c) The e.m.f generated as the coil of an alternating generator rotates is represented in the graph in

Fig. 8.




- i) Give reasons for the changes in the e.m.f as the coil rotates from 00 to 900 and 900 to 1800.
- ii) Sketch on the same diagram a similar graph if the generator was a direct current one.

33. State **one** advantage of:

- i) A lead-acid accumulative over a dry cell
- ii) A dry cell over lead-acid accumulator

34. Three identical cells of e.m.f. 2.0v and of negligible internal resistance are connected as shown in figure below. Determine the ammeter reading.




35. State **one** advantage of:

- i) A lead-acid accumulative over a dry cell
- ii) A dry cell over lead-acid accumulator

36. Compare the property of material used to make a fuse wire to one used to make the filament of a torch bulb.


37. State **two** reasons why the CRO is a more accurate voltmeter than a moving coil voltmeter.

38. The strip below represents part of the electromagnetic spectrum. C is the visible part of the spectrum. A is the region of the shortest wave length and F the highest



Name the sections which represent:

- (i) X-rays
  - (ii) Infra-red
  - (iii) T.V waves
39. Sketch a forward bias characteristics of a P – N junction diode in the axis below




## X-ray

- Give **one** use of X-rays in medicine
- State the factor that affects:-
  - The intensity of X-rays
  - The strength of X-rays
- An x-ray tube must be highly evacuated. Give a reason for this
- In the production of X-rays, electrons are directed at a tungsten target. State a reason why the target is made of tungsten
  - How can the intensity of the X-rays tube be increased?
- Arrange the following waves in order of increasing frequencies: microwaves, x-rays, Infra-red, ultra-violet
  - The table below shows the electromagnetic spectrum;


|            |   |              |   |           |             |
|------------|---|--------------|---|-----------|-------------|
| Gamma rays | A | Ultra violet | B | Infra red | Radio waves |
|------------|---|--------------|---|-----------|-------------|

- Identify **A** and **B**
  - State **one** use for each
- Figure 12** below shows an x-ray tube:



- Indicate on the diagram the path of x-ray beam supplied by the tube
- Why is **M** set at angle of  $45^\circ$  relative to the electron beam?
- Name a suitable metal that can be used for part **M** and give a reason for your choice
- State how the following can be controlled:-
  - Intensity
  - Penetrating power
  - The exposure to patients
- An x-ray tube is operating with an anode potential of 12Kv and a current of 10.0m.A:
  - Calculate the number of electrons hitting the anode per second
  - Determine the velocity with which the electrons strike the target
  - State **one** industrial use of x-rays
- The diagram below shows simplified diagram of an x-ray tube,


**Figure 8**



- Name the parts **A**, **B**, and **C**.
- What adjustments would be made to:


- (i) Increase the penetrating power of the x-rays produced.
- (ii) Increase the intensity of the rays produced.
- (c) Name a suitable material for the part marked **B** and give a reason for your choice.
- (d) Name a suitable material for the part marked **C** and state its purpose.
- (e) Why is it necessary to maintain a vacuum inside the tube?
- (f) State **one** use of x-rays in the following areas; -
  - (i) In medicine
  - (ii) In Industry.

8. a) The figure shows the circuit of a modern X-ray tube




- i) Indicate the path of the X-ray beam supplied by the tube
- ii) Name the part labeled **C** and state its function (2 mks)
- iii) If the tube above is operated at an accelerating potential of 100kV and only 0.05% of the energy of the electrons is converted to X – rays, calculate the wave length of the generated X-rays. (Take electric charge  $e = 1.602 \times 10^{-19}C$ , planks constant  $h = 6.63 \times 10^{-34} Js$ , and speed of light  $c = 3.0 \times 10^8m/s$ )
- iv) State **two** properties of X-rays
- v) State **one** industrial application of X-rays

9. Below is a nuclear reaction



- i) Identify radiation **K**
  - ii) Determine the value of **X** and **Y**
10. a) State the energy changes that take place in an X - ray tube
- b) Electrons in an X-ray tube are accelerated by a potential difference of 40 kV. If 20% of the electrons are converted into X- rays, determine the maximum wavelength of the emitted electrons.
- c) i) Draw a simple circuit consisting of a photocell to show the direction of flow of current
- ii) The diagram below shows a wave form displayed on a CRO screen.



If the Y — gain reads  $0.5\text{V cm}^{-1}$  while the time base is set at  $0.1\text{ ms cm}^{-1}$ , determine the amplitude and frequency of the wave.

11. The table below shows results obtained in an experiment to determine the internal resistance of a cell

|  |      |      |      |      |      |     |
|--|------|------|------|------|------|-----|
| <b>V(V)</b>  | 0.4  | 0.5  | 0.6  | 0.7  | 08   | 1.3 |
| <b>R(<math>\Omega</math>)</b>                        | 0.45 | 0.65 | 0.80 | 1.05 | 1.40 | 2.4 |
| <b><math>1/V</math> (<math>\text{V}^{-1}</math>)</b> |      |      |      |      |      |     |
| <b><math>1/R</math> (<math>\Omega^{-1}</math>)</b>   |      |      |      |      |      |     |


- Complete the table for values of  $1/V$  and  $1/R$  giving your answers to 3 d.p
- Plot a graph of  $1/V$  against  $1/R$
- Use the graph to determine the e.m.f **E** and the internal resistance **r** of the cell given that

$$\frac{E}{V} = \frac{r}{R} + 1$$

### Radioactivity

- Define radioactive decay
  - A radioactive element decays to  $1/128$  of its original activity after 49 days. Determine its half-life
- You are provided with the following:-
    - One diode
    - A load resistor
    - An a.c. source
    - One transformer
    - Using the above apparatus draw a circuit arrangement for half wave rectification
    - Explain how the circuit drawn in (a)(i) above achieves half wave rectification

- Determine the value of **x** and **y** in the nuclear equation below:-




- The half life of a radioactive element is 20minutes. The mass of the element after 120 minutes is 0.03125g. Determine the original mass of the element
  - What evidence supports the fact that gamma rays are not charged
  - Alpha particles have low penetrating power as opposed to beta particles. Give a reason for this
  - A manufacturer wishes to check the thickness of steel sheets he produces. Explain how this can be done using a radioactive source and a counter
- What is meant by radio active decay?
    - Uranium 235 was bombarded with a neutron and fission took place in the following manner:




Determine the values of **a** and **b**

- When carrying out experiments with radio active substance one is instructed that the source should never held with bare hands but with forceps. Give a reason for the instruction


- d) The diagram below shows the paths taken by three radiations **A**, **B** and **C** from a radio active isotope through an electric field



- i) State the charge on plate **Y**
  - ii) Identify the radiation **A** and **C**
  - iii) Give a reason why **C** deviates move **A**
- e)  ${}_{90}^{233}\text{Th}$  disintegrates into radium (**Ra**) by emission of two alpha and two beta particles as in equation  ${}_{90}^{233}\text{Th} \xrightarrow{\text{A}} {}_Z\text{Ra} + 2({}_2^4\text{H}) + 2({}_{-1}^0\text{e})$
- State:
- i) The atomic number of the daughter nuclide
  - ii) The mass number of the daughter nuclide
  - f) One of the application of Beta emission (**B**) is controlling thickness gauge. Explain how they are used for this purpose?
4. The following is a nuclear reaction for a fusion process resulting from the reaction of polonium with loss of beta particles



- (i) Determine the values of **S** and **T**
  - (ii) State the source of the energy released ✓ 1
5. The expression below is an equation for radioactive element **A**. Element **B** and **C** are the daughter nuclides. **A**, **B** and **C** are not the actual symbols of any of the elements



- (a) State what type of radioactive decay this is.
  - (b) What is the value of:  
**X**..... **Y**.....
6. Arrange the following in order of increasing frequency: Red light, Infrared radiation, X-rays, UV radiation, Short –radio waves, TV and Fm radio waves, Am radio waves and Long radio waves.
7. Radium -222 is a radioactive element with a half-life period of 38 sec. What fraction of the mass of a sample of this element remain after 380 sec.
8. (a) Define the term half-life of a radioactive material
- (b) (i) Use the table below to plot a graph of activity against time
- |                                   |     |     |     |     |     |     |     |    |
|-----------------------------------|-----|-----|-----|-----|-----|-----|-----|----|
| Activity (Disintegration/seconds) | 680 | 567 | 474 | 395 | 276 | 160 | 112 | 64 |
| Time <b>t</b> (days)              | 0   | 1   | 2   | 3   | 5   | 8   | 10  | 14 |
- (ii) Find the half-life of the material in days
- (c) The half-life of a radio-active substance is 138 days. A sample of the substance has  $8 \times 10^{10}$  un-decayed nuclei at time  $t = 0$ . How many un-decayed nuclei will be left after 690 days?



- (d) An element x (uranium) decays by emitting two alpha particles and a beta particle to yield element Y
- (i) State the atomic number and mass number of Y
- (ii) Write down the decay equation
9. a) What is meant by radioactive decay?
- b) A radioactive source placed 12cm from the detector produced a constant count rate of 5 counts per minute. When the source is moved close to 3cm, the count rate varies as follows;

|                   |     |    |    |    |    |
|-------------------|-----|----|----|----|----|
| <b>Time</b>       | 0   | 20 | 40 | 60 | 80 |
| <b>Count rate</b> | 101 | 65 | 43 | 29 | 21 |

- i) State the type of radiation emitted.
- ii) Explain the constant count rate when the source is 12cm away.
- iii) Plot a graph of count rate against time (Use graph paper)
- iv) Use the graph to estimate the half life of the element
10. State **one** advantage of:
- i) A lead-acid accumulative over a dry cell
- ii) A dry cell over lead-acid accumulator

### GM induction

- What is Fleming's right hand rule used for?
- State Lenze's law of electromagnetic induction

## SECTION II ANSWERS

### *Pressure*

- $$P_1 = P_2$$

$$h_x e_x g = h_y t_y g \quad \checkmark \frac{1}{2}$$

$$0.06 \times 1200 \times 10 = h_y \times 900 \times 10 \quad \checkmark \frac{1}{2}$$

$$h_y = \frac{120 \times 6}{9000}$$

$$= 0.08m$$

$$h = h_y + 3cm$$

$$= 8cm + 3cm = 11cm \quad \checkmark 1$$

✓ 1

### *Current II*

- $$120 = 4t$$

$$t = \frac{120}{4} \quad \checkmark 1 \text{ mk}$$

$$= 30 \text{ hrs} \quad \checkmark 1 \text{ mk}$$
- $$I = \frac{Q}{t} = \frac{40}{\frac{600}{60}} = \frac{40}{10} = 4A$$
- (a) (i) - When the switch is closed current flows through the coil causing a magnetic field. This repels the magnet towards the chime bar.

- The end of the coil adjacent to the south pole of the bar magnet acquires a south pole.

(ii) – Increase the number of turns in the coil.


- Increase d.c source.

- - Introduce a soft iron core in the coil.

(b) (i) Electrical – magnetic – potential – sound energy

✓ ½      ✓ ½      ✓ ½      ✓ ½

2mks



(ii) Resistance in the circuit increase. ✓1

(iii) Current reduces. ✓1

(iv) Become less stronger. ✓1


(c) (i) - In alternating current there is a change in the magnetic flux of the primary coil linked to the secondary coil. ✓1

- In direct current there's no change in the flux therefore no induction of e.m.f in it.

(ii). - Transformers step up voltage to higher value for transmission which minimizes power loss. ✓1

- They also step down higher voltage to the voltage required by the consumers. ✓1

4.



(a) Current through  $L_1$ :


$$0.8 = 0.15 + 0.28 + L_1$$

$$0.8 = 0.43 + L_1$$

$$L_1 = 0.37A$$


### Thermal expansion

1. Water freezes and the ice formed floats in water because its density is less than that of water, insulating water below it. Temperature increases down the pond because of anomalous expansion of water




## Light.

1.  $\frac{20}{30000} = \frac{2.5}{h_0} \checkmark$   
 $h_0 = \frac{30000 \times 2.5}{20}$   
 $= 3750\text{cm}$   
 $= 3.75\text{m} \checkmark$



## Electrostatics

1. a) Charges at x get neutralized.  $\checkmark 1\text{ mk}$   
C is pointed and due to point action  $\checkmark 1\text{ mk}$  charges leak off from C and are attracted to x neutralizing it.  $\checkmark 1\text{ mk}$   
b) Leaf rises.  $\checkmark 1\text{ mk}$   
A position charge is induced on the surface of hollow conductor, these repels the charges from the cap of the electroscope making the leaf to diverge with the charge.  $\checkmark 1\text{ mk}$



- d) – Smoothing waves.  $\checkmark 1\text{ mk}$   
- Reduction of sparks in induction coil  $\checkmark 1\text{ mk}$   
- In camera flash.  $\checkmark 1\text{ mk}$   
- delay circuits  $\checkmark 1\text{ mk}$  (any two)
2. After touching, the pith balls share the charge and become negative hence they repel.
3. a) – The divergence of B is greater the divergence of A  
- B has a smaller surface area than A, has low capacitance than A ( $Q= CV$ )  
b) - A will have a net positive charge while B will have a net negative charge
4. The two balls will acquire negative charge and repel
5. The gold leaf will diverge further because more positive charges will be repelled from the cap to the leaf by the positively charged rod
6. Each material is brought in turn to touch the cap. The conductor will discharge the electroscope while the insulator will not (accept bring near conductor gauge)
7. A will have a positive charge when charged rod is brought near metal A. positive charges are attracted towards it while the negative charges are repelled

## Measurement II


1. Main scale = 6.5mm  
Thimble scale = 0.34mm  $\checkmark \frac{1}{2}$   
Micrometer reading = 6.84mm  $\checkmark 1$

## Magnetsim

1. a i) The free ends repel because they have some polarity, ✓1 mk

ii) Free ends have different polarity hence attract. ✓1 mk

b i)



1mk direct✓

1mk field theory soft iron ✓

ii) magnetic shielding. ✓1 mk

2 All ferromagnetic materials are attracted by magnets or any magnetic material is attracted

3. (a) (i) Soft iron

- It is easy to magnetize and demagnetize

(ii) Least force =  $mg$

$$= \frac{150 \times 10}{1000} = 1.5N \quad \checkmark$$

(iii) – Increase the number of turns of the coil.


- Increase the amount of current / p.d ✓

(iv) When all the domains have been aligned i.e point of magnetic saturation, the strength of magnet is maximum and cannot increase beyond this point

(b) Correct coil around A

Correct coil around B

Complete correct circuit




4. Domains of soft magnetic materials are easy to arrange and disarrange while the domains of hard magnetic materials are hard to arrange and disarrange.

5. The pin or wooden block was attracted while the one on the metal block was not attracted.

Magnet induces magnetism on the pin. On the iron block which induces magnetism on the iron block. The pin on the wooden block didn't induce magnetism to the wooden block.

6. Correct direction of field



7. The two pith balls separate

Charges (-ve) are transferred from the uncharged pith ball but are not enough to neutralize the charged one. The initially uncharged pith ball now becomes positively charged hence the separation/repulsion.

8. (a) A – North pole

B – North pole

(b) R is stronger. It repels more field lines revealing its strength

9. - Supposed each bar at a time. Displace them in turn and let them come to rest. Note the direction in which they rest. Repeat 2 or 3 times for each. This one that always settles facing N-S directions a magnet

10. A steel bar has dipoles in its domains while aluminium bar does not have the dipoles (1mk)

11. (a) (i) The magnitude of the induced e.m.f is directly proportional to the rate of change of magnetic flux linkage

(ii) The direction of the induced emf is such that the current which it causes to flow produces a

magnetic effect which tends to oppose the change causing it

(b) (i) Resistance of the coil- remedy – thick copper coil )

(ii) Hysteresis loss– remedy – soft iron core

(iii) Eddy currents – remedy – laminated iron core

(iv) Poor flux linkage – Remedy winding primary coil and secondary coil on the same core

(c)  $V_p I_p = V_s I_s$


$$12000 \times 1 = 1800 \times I_s$$

$$I_s = 0.6667A$$

$$\text{Power loss} = I^2 R$$

$$= 0.6667^2 \times 20 = 8.89w \quad (3mks)$$

(d)



(correct diagram and six parts labelled correctly/ correct diagram and four parts labelled correct diagram and less than four parts labelled(1mk)


Current enters the coil thorough the hair springs and flows into the core through the rectangular coils. This causes the coil to be magnetized. The magnetic field created cuts the radial magnetic field of the magnet at right angles. This causes the core to rotate. The rotation of the core is opposed by the torque of the hair spring. When the force due to the rotation of the core is equal to the force due to the torque of the hair spring, the core comes to rest and the pointer gives the reading.

(e) (i)– Using a weak permanent magnet

(ii) Using strong hair springs

(iii) Using few turns of the rectangular coil

12.



13. Radiowaves, visible light, ultraviolet light;

14. a) Induced current flows in such away as to oppose the change producing it

b) i) P - brushless

Q- ship rings

ii) X- North

Y- South

iii)- Increasing speed of rotation of the coil


- Increasing the number of turns in the coil

- Increasing the strength of the magnet

c) i)  $\text{Efficiency} = \frac{\text{Power output}}{\text{Power input}} \times 100\%$   
 $80 = \frac{48 \times 13.5 \times 100}{\text{Power input}}$   
 $\text{Power input} = 810\text{w}$

ii)  $\text{Power input} = I_p \times V_p$   
 $810 = 240 \times I_p$   
 $I_p = 3.375 \text{ A}$

15.



16. Repulsion occur between like poles, unlike poles and magnetic materials

### Reflection at curved surfaces and spherical surfaces

1. a) i)

|               |      |      |       |       |       |       |
|---------------|------|------|-------|-------|-------|-------|
| $\frac{1}{u}$ | 0.05 | 0.04 | 0.033 | 0.025 | 0.020 | 0.014 |
| $\frac{1}{v}$ | 0.05 | 0.06 | 0.67  | 0.75  | 0.080 | 0.086 |

Any 4 correct values = 1 mk  
 Total = 3 mks

iii)  $f = 10 \pm 1 \text{ cm}$   
 Any intercept = 1 mk  
 Rec. - 1 mk  
 Arrange of Reciprocal - 1 mk


b) - Focal length of eye lens is variable while that of camera is fixed. ✓1 mk]  
 - Eye has constant image distance. ✓1 mk

2.  $120 = 4t$   
 $t = \frac{120}{4}$  ✓1 mk  
 $= 30 \text{ hrs}$  ✓1 mk


3. a) Mechanical waves – require medium for propagation  
 Electromagnetic can even travel in a vacuum (No medium required)

b) Frequency =  $1/\text{periodic time}$

c)




4.



Extrapolation of light rays backwards  
 Dotted and upright change formed between C and F  
 Reflected rays shown by arrow pointed correctly  
 angles marked as  
 $i = r = 30^\circ$  and  $i = r = 0$  (implied)

5.



6.

Spherical aberration is a situation where rays parallel but far away (distant) from the principal axis of a concave mirrors fails to pass through its focal point owing to the large radius of curvature of the mirror

7.

Complete the ray diagram below by showing the position of the image.  
 -It is used as a simple microscope or magnifying glass in the laboratory.

8.

- The reflection in a sheet of paper is irregular hence rays interfere with formation of images  
 - Reflection in a plane mirror is regular


9.

Advantage: wide field of vision  
 Disadvantages: Gives a wrong impression of the position of image due to the diminished image formed

10.


**Image V distance( Vcm)**

- Axes
- Scale ✓
- Plotting all values ✓
- Plotting at least 5pts ✓
- Line ✓
- Magnification M ✓



(b) From the graph  
 $V = mf + f$   
 $f = V - \text{intercept}$   
 $= 10\text{cm}$


OR  
 $f = \text{gradient}$   
 $= \frac{50 - 20}{4 - 1}$   
 $= \frac{30}{3}$   
 $= 10\text{cm}$



Correct rays with arrows  
 Object between P and F


11. Image formed by concave mirror is enlarged while that formed by plane mirror is same size as object

12.





13.




✓1 for real ray  
✓1 mk for virtual image

**Observer**

14. (a) (i) Point on the principal axis to which all rays originally close and parallel to the principal axis pass after reflection.  
(ii) Focal length ( $f$ ) – distance between the pole of the mirror (centre of the mirror) and the principal focus  $f$

(b)



(c) (i)  $u = +12\text{cm}, f = +10\text{cm}, V = ?$

$$\frac{1}{f} = \frac{1}{u} + \frac{1}{v}, \frac{1}{10} - \frac{1}{12} = \frac{1}{v}$$

$$\frac{12 - 10}{120} = \frac{2}{120}$$

$$\therefore V = \frac{120}{2} = 60\text{cm}$$

$\therefore$  Image distance = 60cm

- (d) (i) Enlarged, virtual, upright (any two)

(d) (ii)

$$\frac{2x - 320}{340} = 0.8$$


$$\frac{2x - 0.94}{340} = 0.8$$

$$\frac{2x}{340} = 1.74$$

$$2x = 591.6$$

$$x = 295.8\text{M}$$

Let the distance between the boy and the cliff be  $X$  and speed of sound in air is 340m/s



### Linear motion

1.  $Ft = m(V-U)$


$$720 \times 0.1 = 0.6V - 1$$

$$V = \frac{72}{0.6}$$

$$= 120\text{ms}^{-1}$$

✓ 1

2.



### Machines & inclined planes

1. (a) - Energy is the ability to do work  
- Work is done when a force applied on an object moves it through a certain distance

$$\begin{aligned}
 (b) \text{ (i) } V.R &= \frac{1}{\sin \theta} \\
 &= \frac{1}{\sin 30^\circ} \quad \checkmark 1 \\
 &= \frac{1}{0.5} \\
 &= 2 \quad \checkmark 1
 \end{aligned}$$

$$\begin{aligned}
 M.A \times V.R \\
 &= 75 \times 2 \\
 &= 150 \quad \checkmark 1
 \end{aligned}$$


$$\begin{aligned}
 (ii) \text{ Effort} &= \frac{\text{Load}}{M.A} \\
 &= \frac{1500N}{150} \\
 &= 10N
 \end{aligned}$$

### Resistors

1. - The ammeter should be in series  
- Voltmeter in parallel  
- Variable resistor in series  
- The apparatus must be workable

$$\begin{aligned}
 2. \text{ a) } ii) \text{ gradient} &= \frac{10V}{0.5A} \quad \checkmark 1 \text{ mk} \\
 &= 20 \Omega \quad \checkmark 1 \text{ mk}
 \end{aligned}$$

$$\begin{aligned}
 iii) \frac{1}{200} + \frac{1}{100} + \frac{1}{R} &= \frac{1}{20} \quad \checkmark 1 \text{ mk} \\
 \frac{1}{R} &= \frac{1}{20} - \frac{1}{200} - \frac{1}{100}
 \end{aligned}$$



$$\frac{R}{9} = \frac{20}{200} = \frac{200}{100} = 20 \Omega \checkmark 1 \text{ mk}$$

$$= 20 \Omega \checkmark 1 \text{ mk}$$

b) Current through  $10 \Omega$  resistor =  $2A$

$$p.d \text{ across } 10 \Omega \text{ resistor} = 2 \times 10 = 20v$$

$$p.d. \text{ across } 5 \Omega = 20v$$

$$\text{current} = \frac{20}{5} = 4A \checkmark 1 \text{ mk}$$


$$\text{Total current } 4 + 2 = 6A$$

Current through  $2 \Omega = 6A$


$$p.d = 6 \times 2 = 12V$$

$$\text{Total voltage} = 12 + 20 = 32V \checkmark$$

3. (i)



(ii)



b)

i)  $2.1 V$

ii)  $2.1 v - 1.8V = Ir = 0.1V$

$$r = \frac{0.3}{0.1} = 3$$

iii)  $0.1 \times R = 1.8 V$

$$R = 18R$$

4.

- Length of conductor

- Type /nature of material

- Diameter/thickness of material

b)  $E = IR + Ir$

$$3.0 = I(3.5 + 0.5) = I(4.0)$$

$$I = 0.75A$$

5.

a) - The current passing through a conductor is directly proportional to the potential difference across its ends provided temperature and other physical conditions are kept constant

6.

$$\frac{I}{R} = \frac{1}{5} + \frac{1}{2} = \frac{7}{10}$$

$$R = \frac{10}{7} \Omega$$


$$RT = \frac{10}{7} + \frac{3}{1} = \frac{31}{7} \Omega \text{ (Effective resistance)}$$

$$\text{but } I = \frac{12.4 \times 7}{31} = 2.8A$$

$$\therefore p.d \text{ across the } 3\Omega \text{ resistor} = 2.8 \times 3 = 8.4V \text{ (p.d across the } 3\Omega \text{ resistor)}$$

$$\therefore p.d \text{ across the } 5\Omega \text{ and } 2\Omega = 12.4 - 8.4 = 4.0V$$

$$\therefore \text{Current across the } 5\Omega \text{ resistor} = \frac{4.0}{5} = 0.8A$$



(answer 3mks)

7.

$$R_s = 3 + 4$$

$$R_p = \frac{7 \times 5}{7 + 5} = \frac{35}{12} \Omega$$

$$I_1 = \frac{6 \times 12}{35} = 2.057A$$

$$I \text{ through } 3\Omega \text{ resistor} = I_1$$

$$I_1 \times 5 = I_2$$

$$5(2.05 - I_2) \cdot 25 = 5I_2$$

$$10.25 - 5I_2 = I_2 \checkmark 1$$

$$6I_2 = 10.25 \Rightarrow I_2 = \frac{10.25}{6} = 1.708A \checkmark 1$$

8. a) (ii)

(ii) Effective resistance:

$$R_E = 4 + \frac{6 \times 3}{3 + 6}$$

$$= 4 + \frac{18}{9}$$

$$= 6\Omega$$

$$V = IR$$

$$I = \frac{12}{6} = 2A$$

$$\therefore V = 2 \times 4 = 8V$$

(ii)  $V = 12 - 8 \neq 4V$

$$\therefore V = IR$$

$$I = \frac{4}{6} = \frac{2}{3} = 0.667A$$

(iii) Effective resistance in parallel

$$R_E = \frac{6 \times 6}{6 + 6} = 3\Omega$$

The potential drop will increase; hence the reading of  $V$  will decrease

(c) (i) Step-down- The voltage is reduced from 240V to 8V

(ii) To reduce loss of energy due to eddy currents

$$(iii) \frac{V_P}{V_S} = \frac{N_P}{N_S}$$

$$\frac{740}{8} = \frac{4800}{V_S}$$

$$V_S = \frac{4800 \times 8}{240} \checkmark$$

$$= 20 \times 8 = 160 \text{ turns } \checkmark$$

9 (i) – Set Galvanometer to zero balance by adjusting the variable resistor  $L$ .

- P.d across  $BD$  is therefore zero  $\checkmark 1$

- P.d across  $AB = P.d$  across  $AD$

$P.d$  across  $BC = p.d$  across  $DC$   $\checkmark 1$

$I_1$  flows through  $K$  &  $L$  ( $I_1 = I_3$ )  $\checkmark 1$

$I_2$  flows through  $M$  and  $N$  ( $I_2 = I_4$ )  $\checkmark 1$

$$I_1 K = I_2 M \quad \checkmark 1$$

$$I_3 L = I_4 N$$

$$\frac{I_1 K}{I_1 L} = \frac{I_2 M}{I_2 N}$$

$$\frac{K}{L} = \frac{M}{N} \quad \checkmark 1$$

$$\frac{K}{L} = \frac{M}{N} \quad \checkmark 1$$

$\checkmark 1$

(ii) The method does not depend on the accuracy of the current measuring instrument

$$(b) \frac{R}{1} = 0.38$$

$$R = \frac{25\Omega \times 0.38}{0.62}$$

$$= 15.32\Omega \quad \checkmark 1$$

10.  $C_{11} = 4\mu f + 5\mu f = 9\mu f \quad \checkmark \frac{1}{2}$

$$\frac{1}{C_T} = \frac{1}{3} + \frac{1}{9} + \frac{1}{3}$$

$$\frac{3 + 1 + 3}{9} = \frac{7}{9}$$

$$C_T = 9 = 1.29\mu F$$

11. (i) Minimum current is when y is at max resistance, i.e.  $100\Omega$  (x and Y parallel)

current  $I = \frac{220V}{100\Omega}$

$$= 2.2A$$

(ii) Maximum current is when  $R = 500\Omega$  at y (when X and Y are parallel)

$$I = \frac{220V}{50\Omega} = 4.4A$$

(c) (i) For the upper resistors in series

$$R = 1 + 4 = 5\Omega$$

for the lower resistors in series

$$R = 2 + 3 = 5\Omega$$

For the combined resistance of the parallel sets

$$\frac{1}{R} = \frac{1}{5} + \frac{1}{5} = \frac{2}{5}$$

$$R = 2.50\Omega$$

$$\text{Total resistance} = 2.5 + 5.50\Omega = 8.00\Omega$$

(ii) Current  $I_y = 0.5A$

$$V_y = 40\Omega \times 0.25A = 1.0V$$

$$V_2 = 2\Omega \times 0.25A = 0.5V$$

$$V_{y2} = 0.5V$$

\*

(d) - Thickness/x-sectional area – Resistance is inversely proportional to the thickness of a conductor

- Length : Resistance is directly proportional to length of a conductor

## Refraction of light

1.  ${}_a n_p = 1.47$  and  ${}_a n_g = 1.55$

$${}_g n_p = {}_g n_a \times {}_a n_p$$

$$\checkmark \quad = \frac{{}_a n_p}{{}_a n_g}$$

$$= \frac{1.47}{1.5} = 0.9484$$

$$\sin C = \frac{1}{n} = \sqrt{0.9484}$$

$$C = \sin^{-1}(0.9484)$$

$$C = 71.5^\circ$$

2. (i) for incident and reflected ray  
 (ii) The ray undergoes total internal reflection. Since angle of incidence is greater than  $a^\circ$  the critical angle.
3. a) The ratio the  $\sin \theta$  of the angle of incidence to the  $\sin e$  of the angle of refraction is a constant for a pair of media  
 b) When a ray is moving from an optically dense medium to a less optically dense medium or when the angle of incidence in the optically dense medium is greater than the critical angle

$$\begin{aligned} c) a^n_g &= \frac{\sin i}{\sin r} \\ &= \frac{\sin 48^\circ}{\sin 32^\circ} \\ &= 1.40, \text{ Accept } 1.402 \end{aligned}$$

d) Separation of colours of light from white light

$$\begin{aligned} 4. \quad gnw &= gna \times anw \\ &= \frac{2}{3} \times \frac{4}{3} \\ &= \frac{8}{9} \end{aligned}$$

$$\frac{8}{9} = \frac{\sin \theta}{\sin 40}$$

$$\begin{aligned} \sin \theta &= \frac{8}{9} \sin 40 = 0.5713 \\ &= 34.84^\circ \quad \checkmark \end{aligned}$$

5. If the refractive index of medium 1 is  $\frac{4}{3}$  and that of medium 2 is  $\frac{3}{2}$ . Calculate angle  $r$

$$\begin{aligned} n_1 \sin \theta_1 &= n_2 \sin \theta_2 \\ \frac{4}{3} \sin 35 &= \frac{3}{2} \sin \theta_2 \\ \sin \theta_2 &= \frac{4}{3} \times \frac{2}{3} \sin 35 = 0.5098 \\ \theta_2 &= 30.654 \end{aligned}$$

6. a)

$$\begin{aligned} ii) n &= \frac{1}{\sin 42} \\ \frac{\sin 25}{\sin r} &= \frac{1}{R} \\ \frac{\sin 25}{\sin r} &= \sin 42 \\ \sin r &= \frac{\sin 25}{\sin 42} \\ \sin r &= \frac{\sin 25}{\sin 42} \end{aligned}$$

$$\begin{aligned} &= 0.631593 \\ r &= \sin^{-1}(0.631593) \\ &= 39.17^\circ \text{ (accept } 39.2^\circ) \end{aligned}$$

7. (a) (i) - When a ray is moving from an optically denser medium to a less optically dense medium.  
 - When the angle of incidence in the optically denser medium is greater than the critical angle (any 1)

$$(b) \sin C = \frac{n_2}{n_1} = \frac{1.3}{1.5} = 0.866$$

$$\angle C = \sin^{-1} 0.866 \quad \therefore \angle C = 60.1^\circ$$

- (c) (i) From the len's formula  $1 = \frac{1}{v} + \frac{1}{u}$  and dividing both sides by  $V$ ,

$$V = 1 + \frac{V}{u}, \text{ but } \frac{V}{u} = M$$

$$\frac{V}{f} = 1 + M \text{ and making } M \text{ the subject ;}$$

$$M = \frac{v}{f} - 1$$

(ii) Graph: - scale used (1mk)

- Labeling axis
- Straight line
- Points
- Gradient/slope

$$\frac{1}{v} = \frac{1}{u} - \frac{1}{f}$$

$$\frac{1}{f} = \frac{1}{u} + \frac{1}{v} \text{ or } \frac{1}{v} = \frac{1}{f} - \frac{1}{u}$$

Gradient = Negative

$$\frac{1}{v} \text{ Intercept} = \frac{1}{f}$$

## Sound II

1.  $2d = s \times t$   $T = 0.1s$   
 $d = \frac{s \times t}{2}$  ✓1 OR  $f = \frac{1}{0.1} = 10\text{Hz}$   
 $= \frac{330 \times 0.1}{2}$  ✓1  $c = \lambda f$   
 $= 330 = \lambda \times 10$   
 $= 16.5 \text{ m}$  ✓1  $x = 33$   
*But*  $\alpha d = \lambda = 33$   
 $d = \frac{33}{2} = 16.5$

2.  $\text{Velocity} = \frac{s}{t}$  ✓1  
 $= \frac{220 \times 2}{1.29}$  ✓1  
 $= 341.085\text{m/s}$  ✓1

3.  $\text{Difference in time between the two points} = 0.3 - 0.25 = 0.05\text{secs.}$   
 $\text{Speed} = \frac{D}{T}$  ✓1  
 $= \frac{17\text{m}}{0.05\text{sec}}$  ✓1  
 $= 340\text{m/s}$  ✓1

4. At night, the masses of air close to the ground are cooler than those higher above. Sound get refracted towards the earth ✓

5. - Place a clock near the end of one tube and point one open end towards a hard surface (wall) at an angle  $i$  ✓

- With the ear close to the end of second tube, open tube T2, listen to the reflection of the sound from the wall at different angles of  $r$  and note where the sound is loudest

- It will be observed that maximum (loudest) sound is heard when  $\angle i = \angle r$


6. - Light must travel from optically dense to less dense medium ✓

- The angle of incidence must exceed the critical angle ✓

## Thin lenses

1. Short-sightedness or myopia  
 cause- the eye-ball is too long for the relaxed focal length

2.



b) i)  $\frac{1}{F} = \frac{1}{U} + \frac{1}{V}$   
 $\frac{1}{15} = \frac{1}{25} + \frac{1}{V}$   
 $V = 37.5 \text{ cm}$

ii)  $U = 52.5 - 37.5$   
 $= 15 \text{ cm}$   
 $\frac{1}{F} = \frac{1}{U} + \frac{1}{V}$   
 $\frac{1}{V} = \frac{1}{30} - \frac{1}{15}$   
 $= -30 \text{ cm}$

3. (a) (i) - Candle placed at a distance  $u$ , from the lens and the position of the screen is adjusted until a sharp image is formed/obtained  
 - The distance,  $V$ , between the lens and screen is measured  
 - The procedure is repeated to get other values of  $V$  and  $u$   
 - For each set of  $u$  and  $v$  the value of  $f$  is determined using the formula

(ii) The image would be virtual and cannot be formed on the screen

(b) Extrapolate both sides of the graph and read-off

$\frac{1}{u}$  and  $\frac{1}{v} = 0.25 = \frac{1}{4}$

$\frac{1}{u} = \frac{1}{v}$

$\frac{1}{f} = \frac{1}{u} = f = 4$  or

$\frac{1}{f} = \frac{1}{v}$

$\frac{1}{f} = \frac{1}{v} = f = 4$

$\frac{1}{f} = \frac{1}{v}$

(c)  $M = \frac{u}{v} = 2$

$\frac{V}{15} = 2$

$15$

$v = 30 \text{ cm}$

$\frac{1}{f} = \frac{1}{15} + \frac{1}{30} \quad f = 10 \text{ cm}$

$\frac{1}{f} = \frac{1}{15} + \frac{1}{30}$

4. (a) (i) X-Intercept =  $4.5 \times 10^{14} \text{ Hz}$

(ii) Slope =  $\frac{h}{e} = h = e \times \text{slope}$

$= e \times \frac{6.6 - 0}{(6 - 4.5) \times 10^{14} \text{ s}^{-1}}$

$= 1.610^{-19} \times 4 \times 10^{-15}$

$= 6.4 \times 10^{-34} \text{ Js}$




$$\begin{aligned}
 \text{(iii) } W_0 &= hf_0 \\
 &= 6.4 \times 10^{-34} \text{ Js} \times 4.5 \times 10^{14} \text{ s}^{-1} \\
 &= 2.88 \times 10^{-19} \text{ J}
 \end{aligned}$$

- (b) (i) The leaf falls ✓ Collapses ✓1  
(ii) The electrons are repelled causing the leaf potential to decrease  
(iii) NO effect on the leaf. Light emitted by red light doesn't have enough energy to cause photoelectric effect. ✓1  
(iv) Light is a wave, it carries energy in small packets (photons). ✓1

5. (a) This distance is the focal length ✓1  
(b) Linear magnification is the ratio of image height to the object height or image distance to object distance ✓1

Eye defect – short sightedness (myopia)




6. a) By adjusting the lens' position obtain a sharp image on the screen as shown above.  
▪  $d = f$

$$\begin{aligned}
 \text{b) } v &= 5.5 \times 4 = 22 \text{ cm} \\
 \mu &= 7 \times 4 = 28 \text{ cm} \\
 M = \frac{v}{\mu} &= \frac{22}{28} = 0.7857
 \end{aligned}$$

- c) i) Long sightedness  
ii)

$$\begin{aligned}
 \text{7. } f &= -25 \text{ cm}, \quad u = 20 \text{ cm} \\
 \frac{1}{f} &= \frac{1}{u} + \frac{1}{v} \\
 \frac{1}{-25} &= \frac{1}{20} + \frac{1}{v} \\
 v &= \frac{-20 \times -25}{-25 - 20} = -11.1 \text{ cm} \quad \checkmark \\
 \frac{11.1}{20} &= \frac{h}{10} \quad \checkmark \\
 h &= 5.56 \text{ cm} \quad \checkmark
 \end{aligned}$$

8. (a)



b (i)  $f = 30\text{cm}$ ,  $v = 90\text{cm}$

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f}$$

$$\frac{1}{u} + \frac{1}{90} = \frac{1}{30}$$

$$\frac{1}{u} = \frac{1}{30} - \frac{1}{90} = \frac{1}{45}$$


$$u = 45\text{cm}$$

(ii)  $m = \frac{v}{u}$


$$= \frac{90}{45}$$

$$= 2$$

9.



10.



## Quality of heat

1. Heat lost by the metal = Heat gained by the water  
 $MmCm = MWCW$   
 $0.5 \times CM(100-21)^\circ C = 0.2 \times 4200 \times (21-15)^\circ C$   
 $Cm = \frac{5040}{39.5} = 127.59 \text{ JKg}^{-1} \text{ K}^{-1}$

## Waves II


1. (a)(i) – Stationary wave:- waveform do not move through the medium and therefore energy is transferred from the source to the same point away.  
 - Progressive wave- wave forms more though the medium and therefore energy is transferred from the source to the same point away.  
 (ii) Open pipe has both odd and even harmonics.

(b) (i) For closed pipe


$$\frac{1}{2} \lambda = 0.6$$

$$\lambda = 1.2 \text{ m}$$

$$V = f_0 \lambda, f_0 = \frac{V}{\lambda} = \frac{(340) \text{ Hz}}{1.2} = 283.3 \text{ Hz}$$



(ii) For open pipe, one end open



End correction ignored,


$$\circ \text{ If } \frac{1}{4} \lambda = 0.6$$

$$\lambda = 0.6 \times 4 = 2.4 \text{ m}$$

$$f_0 = \frac{V}{\lambda} = \frac{(340) \text{ Hz}}{2.4}$$

$$= 141.7 \text{ Hz}$$

2.



3. a) i) Sound is transmitted as a longitudinal wave electromagnetic is transmitted as a transverse wave. ✓1 mk  
 (ii)  $S = \frac{2d}{t}$  ✓1 mk  
 $= \frac{2 \times 400}{2.5} = \frac{800}{2.5}$  ✓1 mk  
 $= 320 \text{ m/s}$  ✓1 mk

iii) Let distance from mine worker to furthest cliff be  $y$ . then

$$S = \frac{2d}{t}$$

$$320 = \frac{2y}{2 \times 2.5} \quad y = \frac{1440}{2}$$

$$2y = 320 \times 4.5$$

$$2y = 1440$$

$$= 720 \text{ m}$$

$$\text{total distance} = 720 + 400$$

$$x = \underline{\underline{1120 \text{ m}}} \quad \checkmark 1 \text{ mk}$$

b) i) Bright fringe.  $\checkmark 1 \text{ mk}$

ii) Loud sound.  $\checkmark 1 \text{ mk}$

c) slit must be very narrow (less than the wavelength of light)

4.  $6\lambda = 15 \text{ cm}$

$$1 \lambda = \frac{1 \times 15}{6} \quad \checkmark 1 \text{ mk}$$

$$= 2.5 \text{ cm.} \quad \checkmark 1 \text{ mk}$$

5. (a) i)  $= \frac{8}{2} = 4 \text{ m}$

$$f = \frac{v}{\lambda}$$

$$= \frac{340}{4}$$


$$= 80 \text{ Hz}$$

ii)  $T = \frac{1}{f}$


$$= \frac{1}{80}$$

$$= 0.0125 \text{ seconds} \quad \checkmark$$

b)



6.



7. (a) In a progressive wave, all particles have the same amplitude and each particle is out of phase with the particle next to it. In stationary wave, vibrations of particles at points between successive nodes are in phase and the amplitude of particles between nodes is different

(b) Must have:- (i) Same speed

(ii) Same frequency

(iii) Same of nearly equal amplitudes

(c)  $OO_1$  – Loud sound (constructive interference) waves arrive in phase

AA1 – Loud and soft sound (Destructive and constructive interference)

(d)  $f = \frac{V}{\lambda}$  ✓

$\lambda = \frac{1.25}{2} = 0.625$

$f = \frac{50}{0.625}$   
 $= 80\text{Hz}$

(e) – Constant phase difference  
 - Nearly same amplitude

8. The width of the opening must be smaller than the wavelength of the wave.

9. - Diffraction- is the spreading waves beyond obstacle openings (1mk)  
 - Refraction- is the bending of waves when they change the medium

10. Period  $T = 16 \times 10^{-2}$  (read off)

Frequency  $= \frac{1}{T}$   
 $= \frac{1}{16 \times 10^{-2}} \text{Hz}$   
 $= 6.25 \text{Hz}$

11.  $\frac{1}{2}$  oscillations = 0.2 seconds

0.2 seconds = 0.5 oscillations

1 second  $= \frac{0.5}{0.2} = \frac{1}{0.4}$   
 $= \frac{1}{4} \times 10 = 0.25 \times 10$   
 $= 2.5 \text{Hz}$

12. Time for one complete cycle = 0.5 seconds

$\therefore T = 0.5 \text{seconds}$


$f = \frac{1}{T}$  ✓ 1

$= \frac{1}{0.5}$

$= 2 \text{Hz}$  ✓ 1

13. From A to X is shallow since the ripples are close to one another.

14.



15. (a) (i) Electromagnetic waves travel through a vacuum while mechanical waves need a medium

(ii) Mechanical waves – sound waves



- water waves

- shock

- electromagnetic waves – light waves

- radio waves

- X-rays, gamma rays (any 1)



(ii)  $f = \frac{1}{T}$  where  
 $f$  = focal length and  $T$  is the periodic time

(iii) Radio reception is better because radio waves have longer wavelength hence easily diffracted unlike TV waves which have shorter wavelength

### Work, energy and power

1.  $D = IV$  |  $P = I^2 R$   
 $I = \frac{P}{V}$  |  $R = \frac{P}{I^2}$   
 $= \frac{45}{240} \checkmark 1 \text{ mk}$  |  $= \frac{45}{(0.1875)^2} \checkmark 1 \text{ mk}$   
 $= 0.1875 \text{ A}$  |  $= \frac{45}{0.03515} = 1280 \Omega \checkmark 1 \text{ mk}$

2.  $P = \frac{V^2}{R}$   
 $R = \frac{(240)^2}{40} \checkmark$   
 $= 1440 \Omega \checkmark$

3.  $\frac{P}{V} = I$   
 $\frac{3000}{250} = 12 \text{ A}$   
 suitable fuse 13A

4. An electric kettle is rated 3KW, 250V. Determine the resistance of the coil  
 $P = IV = \frac{V^2}{R}$   
 $3000 = \frac{250^2}{R}$   
 $R = \frac{62500}{3000}$

$$= 20.83 \Omega$$

5. Energy  $E_1 = \text{Power} \times \text{time}$   
 $= 3000 \times 8 \times 60$   
 $= 1440000 \text{ J} \checkmark 1$

6.  $= 2.4 \times 4.2 \text{ Jg}^{-1} \text{ K}^{-1} \times 46 \text{ K} (1 \text{mk})$   
 $= 2.4 \times 4200 \text{ JKg}^{-1} \text{ K}^{-1} \times 46 \text{ K} (1 \text{mk})$   
 $= 463 \ 6805 \text{ J}$

Let the time be  $t$

$$\text{Energy } H = Pt = 3450 \text{ W} \times t$$

$$3450t = 463 \ 680 \text{ J}$$

$$t = 134.4 \text{ s}$$

### Floating and sinking

- Production of electrons from metal surface when suitable heat energy falls on it.
- (a) (i) The weight of a fluid displaced by a body which is partially or wholly submerged is equal to the upthrust experienced.  $\checkmark 1$

(ii) A floating body displaces its own weight of the fluid it floats on

(b) (i) Relative density  $= \frac{W_{\text{Block}} - W_{\text{liquid}}}{W_{\text{Block}} - W_{\text{water}}}$

$$= \frac{120 - 94}{120 - 100}$$

$$= 1.3$$

$$\text{Density of liquid} = 1.3 \times 1000$$

$$= 1300 \text{ Kg/m}^3$$

$$D = \frac{M}{V}$$

$$\text{Mass} = \frac{20 \text{ N}}{10} = 2 \text{ kg}$$

$$\text{Volume} = \frac{2 \text{ kg}}{1000} = 0.002 \text{ m}^3$$


$$D = \frac{2 \text{ kg}}{0.002} \checkmark 1$$

$\checkmark 1$

### Photoelectric effect

- i) – The emission of electrons from metal surface when radiation of unstable wave length falls on it  $\checkmark$
  - ii) The maximum wavelength beyond which no photoelectric effect occurs  $\checkmark$

b)



$$\begin{aligned}
 \text{ii) } EK &= hf - W \\
 h &= \text{gradient } \checkmark \\
 &= (6.2 - 2.2) \times 10^{-19} \checkmark \\
 &= (16 - 10) \times 10^{14} \\
 &= 6.667 \times 10^{-34} \text{Js } \checkmark
 \end{aligned}$$

$$\begin{aligned}
 \text{iii) } W_0 &= hf_0 \\
 &= 6.667 \times 6.4 \times 10^{14} \times 10^{-34} \checkmark \\
 &= 4.267 \times 10^{-19} \text{J} \checkmark
 \end{aligned}$$

2. (a) (i) X-Intercept =  $4.5 \times 10^{14} \text{Hz}$

(ii) Slope =  $\frac{e}{h} \checkmark$   $h = e \times \text{slope}$

$$\begin{aligned}
 &= e \times (6.6 - 0) \checkmark \\
 &= (6 - 4.5) \times 10^{14} \text{ s}^{-1} \\
 &= 1.610^{-19} \times 4 \times 10^{-15} \\
 &= 6.4 \times 10^{-34} \text{Js } \checkmark
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) } W_0 &= hf_0 \\
 &= 6.4 \times 10^{-34} \text{Js} \times 4.5 \times 10^{14} \text{ s}^{-1} \\
 &= 2.88 \times 10^{-19} \text{J}
 \end{aligned}$$

(b) (i) The leaf falls  $\checkmark$  Collapses  $\checkmark$  1

(ii) The electrons are repelled causing the leaf potential to decrease

(iii) NO effect on the leaf. Light emitted by red light doesn't have enough energy to cause

photoelectric effect.  $\checkmark$  1

(iv) Light is a wave, it carries energy in small packets (photons).  $\checkmark$  1

3. Calculate the wavelength of Green light whose energy is  $3.37 \times 10^{-19} \text{J}$ .

( $h = 6.63 \times 10^{-34} \text{Js}$ ,  $c = 3.0 \times 10^8 \text{m/s}$ )

$$\begin{array}{l|l|l}
 \lambda = \frac{c}{f} & f = \frac{3.37 \times 10^{-19} \text{J}}{6.63 \times 10^{-34} \text{Js}} & \lambda = \frac{3.0 \times 10^8 \text{m/s}}{5.083 \times 10^{14} \text{Hz}} \\
 & = 5.083 \times 10^{14} \text{Hz} & = 5.902 \times 10^{-7} \text{m} \\
 f = \frac{E}{h} & &
 \end{array}$$

4. a) This is the least radiation energy required to just dislodge an electron from a metal surface.

b) The energy of the radiation. The higher the energy the higher the velocity of photo electrons

Frequency  $\times 10^{14} \text{Hz}$     7.959    7.43    6.88    6.10    5.49

i) On the grid provided plot a graph of stopping potential (Y-axis) against frequency

Graph (diagram)

ii) From your graph determine:

The threshold frequency

$f_0 = 4.5 \times 10^{14} \text{Hz}$

b) The plank's constant,  $h$

( $e = 1.6 \times 10^{-19} \text{Coulomb}$ ,  $c = 3.0 \times 10^8 \text{m/s}$ )

$$\begin{array}{l|l|l}
 e & \begin{aligned} eVs &= hf - hf_0 \\ Vs &= h(f - hf_0) \end{aligned} & \begin{aligned} \text{gradient} &= \frac{1.15 - 0.93}{(7.43 - 6.98) \times 10^{14}} \\ &= 0.22 \times 10^{-14} \\ &= 0.55 \\ \text{gradient} &= h \\ &= 0.4 \times 10^{-14} \\ &= 0.4 \times 10^{-14} \times 1.602 \times 10^{-19} \end{aligned} \\
 & & h = 6.408 \times 10^{-34} \text{Js}
 \end{array}$$



5. a) It controls the intensity of electron leaving the electron gun controlling the brightness of the spot on the screen.  
 b) The magnetic field deflection system make electrons span the whole screen unlike the electric field deflection system.  
 c) i) Calculate the frequency of the signal

$$T = 25\text{ms/div} \times 2 \text{ div} \quad \left| \quad f = \frac{1}{50/1000} \right.$$

$$= 50 \text{ ms}$$

$$F = \frac{1}{T} \quad \left| \quad = 20\text{HZ} \right.$$

- ii) What is the peak voltage of the signal  
 ○ peak voltage =  $21 \times 2.5\text{v/div}$   
 = 5 Volts

6. (i) Graph is extrapolated to meet x-axis  
 $f_0 = 7 \times 10^{14}\text{Hz}$  ✓ 1 ✓ 1

(ii) Gradient =  $\frac{\Delta V_s}{\Delta f}$   
 $= \frac{1.75 - 0}{12 - 7}$  ✓ 1  
 $= \frac{1.75}{5 \times 10^{14}} = 0.35 \times 10^{-14}$   
 $= 3.5 \times 10^{-15}$  ✓ 1

$$h = \text{Gradient} \times e$$

$$= 3.5 \times 10^{-15} \times 1.6 \times 10^{-19}$$

$$= 5.6 \times 10^{-34}\text{Js}$$

✓ 1  
 (iii)  $W = hf_0$   
 $= 5.6 \times 10^{-34} \times 7 \times 10^{14}$  ✓ 1  
 $= 3.92 \times 10^{-19} \text{ J}$  ✓ 1

7. a) – Has infinite resistance/ does not take up any current  
 - Sensitive/ does not require heating time  
 b) i) A – Grid  
 B- Electron gun  
 ii) C – Vertical deflection of the beam  
 D- Horizontal deflection of the beam  
 iii) – By thermionic emission as heating the filament

8. a) Electrons being ejected from metal surfaces by use of electromagnetic waves

b) i) X – intercept =  $1.0 \times 10^{15}$

ii) From  $K.E = hf$

Planks constant ( $h$ ) = gradient of graph


$$= \frac{(8.2 - 0) \times 10^{-19}}{(2.5 - 1.0) \times 10^{15}}$$

$$= \frac{8.2 \times 10^{-19}}{1.5 \times 10^{15}}$$

$$H = 5.5 \times 10^{-34} \text{ JS}$$

iii) Work function,  $W_0 = 5.5 \times 10^{-34} \times 1.0 \times 10^{15}$   
 $= 5.5 \times 10^{-19} \text{ J}$

9. (a) (i)



Correct polarity

(ii) No change in the amount of photo current. Change in wavelength/frequency of the radiation does not affect the amount of photo electrons produced. It is the number of photo electrons that determines the photocurrent.

(b) (i) Total resistance = gradient

$$= \frac{7.5 - 0}{0.375 - 0} = 20\Omega$$

(ii) Combine d resistance =  $\frac{100R}{100 + R}$

$$\frac{100R}{100 + R} = 20$$

$$100R = 20R + 2000$$

$$80R = 2000 \quad R = 25\Omega$$

(c) (i) Alternating magnetic flux in the coil induces current in the core of the same coil causing eddy currents.

- Eddy currents are minimized by lamination of the core

$$(ii) \frac{V_s}{X_p} = \frac{N_s}{N_p} = \frac{\mathcal{G}}{240} = \frac{N_s}{4200} \quad N_s = \frac{4200 \times 8}{240}$$


- 10 - Intensity of the radiation  
- Energy of the radiation  
- Type of the metal

$$N_s = 140 \text{ turns}$$

11. a) i) Emission of electrons from metal surface by electromagnetic radiation falling on the surface

$$b) \quad ii) M = \frac{u}{e} = \frac{0.56 - 0}{(6.4 \times 10^{14})} = \frac{0.56}{1.4 \times 10^{14}} = 40 \times 10^{-15}$$

$$h = 4.0 \times 10^{-15} \times 1.6 \times 10^{-19} = 6.4 \times 10^{-34} \text{ J}$$



c)  $hf = Q + K.E$

$$6.4 \times 10^{-34} \times 3.0 \times 10^{15} = 6.4 \times 10^{-19} + K.E$$

$$K.E = 19.2 \times 10^{-19} - 6.4 \times 10^{-19}$$


$$= (19.2 - 6.4) \times 10^{-19}$$

$$= 12.8 \times 10^{-19}$$

$$= 1.28 \times 10^{-18} \text{ J}$$

## ELECTRICITY & Electronics

1. From acceleration  $a = V-u$  and making  $V$  the subject ;  $V = at + u$  or  $V = ut + at$
2. Polarisation ✓1 - Corrected by adding a depolarizer ✓1  
Or local adic - Corrected by amalgamation or use of pure zinc.
3.  $RT = \frac{4.5(4.5)}{9} \sqrt{\phantom{x}}$   
 $= 2.25\Omega$   
 $I = \frac{V}{R}$   
 $= \frac{6}{2.25}$   
 $= 2.667A \sqrt{\phantom{x}}$
4. i)  
ii) Straight vertical line observed since Y- gain is connected leading to vertical deflection
5. Y - Neutral Z - Live ✓
6. - Decreasing area of overlap ✓  
 - Removal of dielectric ✓  
 - Increasing separation distance ✓  
 (a) (i) For  $W$  to occupy a smaller space  
 (ii) Offers high resistance ✓
- (b) (i) To reduce power loss for long distance power transmission  
 (ii) To be able to step it up or down depending on need ✓
- (c) - High current/charge carrying capacity/density ✓  
 - Lighter ✓



(ii) Melts and breaks the current if there is an overload to protect the load connected to the main output

(e) Power consumed for 30days  
 $= 1.5 \times 2 \times 30 = 90KW$   
 Cost of the Electricity consumed  
 $= 90KW \times 6.70 = \text{Kshs.}603$

8. (a) (i) Z - Zinc plate




Y – Dilute sulphuric acid

(ii) – Local action

- Occurs due to the reaction between Zinc plate and dilute sulphuric acid thus Zinc is eaten away


(iii) Use of depolariser (Potassium dichromate)

(b) (i) Doping intrinsic semi-conductor with group III elements



9. Conductivity increases with increase in temperature. Increase in temperature makes valance electrons gain kinetic energy and jump to the conduction band

10. (all diagram correct) (2mks)



11. A1 shows a deflection while A2 doesn't. This is because diode P is forward biased while Q is reverse biased i.e it offers high resistance.

12. - Eight dry cells have a very high internal resistance compared to the car battery hence very little current can be drawn from the dry cells.

13. a) i) p-type :- it is obtained by doping an intrinsic semi conductor using a group 3 impurity.

n-type :- it is obtained by doping an intrinsic semi conductor using a group 5 impurity

ii) The figure below shows a bridge rectifier

A capacitor has been connected across the resistors as shown. Sketch on the axes below the wave form when a C-R-O is connected across the resistor; R


b)  $e.m.f = 1.5 \text{ v}$

ii) Terminal voltage =  $1.25 \text{ v}$

iii) Calculate the internal resistance of the cell

|                                  |                      |
|----------------------------------|----------------------|
| $e = I(r + R)$                   | $I = 1.25$           |
| $1.5 \text{ v} = 1r + 1.25$      | $4.8$                |
| $I r = 1.5 - 1.25$               | $= 0.2604 \text{ A}$ |
| $I r = 0.25 \text{ v}$           | $r = 0.25$           |
| $\text{But } I = \frac{1.25}{R}$ | $0.2604$             |
|                                  | $= 0.96 \Omega$      |


14. A fuse is a safety device is used to disconnect the circuit when excess current flows through it, it melts.
15. Distinguish between Topping and Dopping  
 Topping:- The addition of distilled water into a lead acid accumulator to improve on the ion concentration.  
 Dopping:- Addition of impurities to an intrinsic semiconductors to improve on its conductivity.
16. a) A is Copper B is a dilute acid (hydrochloric or sulphuric acid)  
 b) - Polarization  
 - Local Action
17. a) Cathode:- Zinc  
 b) two causes of this observation.
- Due to defects that the cell suffers. These are
  - Local action. The eating away of Zinc (cathode) by the acid.
- Polarization:- the formation of  $\text{H}_2$  bubbles at the anode insulating it.
18. - Colour televisions have three electron guns compared to one in black and white televisions
19. - Capacitance is inversely proportional to the distance of separation between the plates (1mk)  
 - Capacitance is directly proportional by the area of overlap between the plates (1mk)
20. a) i) Local action  
 ii) Polarization  
 b) - Connect the three bulbs in parallel so that their internal resistance is reduced.  
 - This arrangement increases the current making the bulbs very bright
- 21.



22. To increase the conductivity of a semi-conductor
23. (a) Forward biased  
 (b) Resistance in the circuit is varied by moving the jockey along  $R$ .  
 - A series of values of voltage for the corresponding values of current are obtained  
 (c) (iii) They are not deflected by both electric and magnetic fields  
 (iv) Alpha particles are heavy (massive) ✓  
 (v) The sheets are brought in turns between radioactive source and the counter.

- The count rate is a measure of the thickness of the metal sheet. ✓
24. a)  $QI = CV = 0.3 F \times 4.5 = 13.5c$   
 b)  $CT = C1 + C2$   
 $(0.3 + 0.5) F = 0.8F$
- c) i)  $4.5V$   
 ii) Voltmeter reads less than  $4.5V$   
 iii) The drop of p.d in C (ii) is because the charge on C1 is because is distributed to C2. Since values of C1 and C2 remain constant when Q on C1 reduces, the  $Q = CIV$  implies V must reduce also, hence reading reduced

25. (a) Ability of a capacitor to store charge  
 (b) (i) For charge distribution  
 Raised leaf



- (ii) The leaf divergence increased.  
 The potential on of increases due to reduced capacitance since distance of separation is increased.
- (iii) They are not deflected by both electric and magnetic fields  
 (iv) Alpha particles are heavy (massive)  
 (v) The sheets are brought in turns between radioactive source and the counter.  
 - The count rate is a measure of the thickness of the metal sheet.

26. a) - Area of the plates  
 - Distance of separation of the plates  
 - The electric constant
- b) - Capacitors in parallel  
 $1\mu F + 2\mu F + 3\mu F = 6\mu F$   
 - Capacitors in series:  
 $1/_{6MF} + 1/_{3MF} = 2F$

$$Q = CV$$

$$= 2.0 \times 10^{-6} F \times 10v$$

$$= 2.0 \times 10^{-5} C$$

27. a) - The process in which an impurity is introduced into a pure semi- conductor  
 b) - Intrinsic - pure semi- conductors where charge carriers come from within  
 - Extrinsic - pure semi- conductor which has been doped  
 c) Bulb  $B_2$  lights
28. a) Define Eddy currents  
 ■ These are current loops that develop in the core there is a change in the magnetic field linking with the core.
- b) i) Using Lenz's law indicate the direction of current through the galvanometer
- ii) I. The magnet is moved away from the solenoid  
 • The deflection of the galvanometer changes since direction of current is opposite the previous one
- II. The magnet is placed stationary in the solenoid.  
 ○ The galvanometer does not deflect since no current flows

c) The p.d across the primary coil is  $240V$

$$\frac{N_p}{N_s} = \frac{V_p}{V_s} = \frac{240}{V_s}$$

$$12 = \frac{240}{V_s}$$

$$V_s = \frac{240}{12} = 20V$$


$$V_{BC} = \frac{6}{9} \times 20V$$

$$= \frac{120}{9}$$

$$= 13.33$$

- d) i) It lights because during the rotation of the wheel there is an indication of part (i) the coil creating a current that flows through the bulb and it lights  
 ii) By making the wheel rotate faster or by making bicycle more faster
29. Evaporation and cell reaction cause loss of water. Distilled water does not introduce impurities to the cell


30.



### X-ray

1. Treatment of cancer, tumors
2. (i) Size of heater current/filament current  
(ii) Accelerating potential/kinetic energy of electrons/anode Voltage
3. To avoid collisions between the moving electrons and air particles
4. a) – Tungsten has high melting point and therefore it would not melt at elevated temperatures  
b) – Increasing filament voltage or heating current
5. a) Micro waves, infrared, ultra violet X – rays  
b) i) A – X – rays                      B – visible light  
ii) - X – rays – viewing bone fracture/ foreign objects in the body  
- Visible light – ordinary photography/ optical fibre

6. (a)



✓1

(b) - To direct x-rays out of the tube through the window on the shield. ✓1

(c) - Tungsten or molybdenum. ✓1

- High melting point thus it can withstand high temperature. ✓1

(d) (i) Heater current (Filament current)

(ii) Anode potential (operating potential)

(iii) - Covering with protective materials where x-rays are not required

- Minimize exposure time as much as possible

- Reduce number of exposure as much as possible (any 1-1mk)

(e) (i)  $Q = It = 10 \times 10^{-3}C (= 1.6 \times 10^{-19}C)$

$$10 \times 10^{-3}C = 1.6 \times 10^{-19} \times n$$

$$n = \frac{10 \times 10^{-3}}{1.6 \times 10^{-19}} = 6.25 \times 10^{16} \text{ electrons } \checkmark 1$$

$$\begin{aligned}
 \text{(ii)} \quad \frac{1}{2} m_e v^2 &= eV \\
 v &= \sqrt{\frac{2eV}{m_e}} \\
 &= \sqrt{\frac{2 \times 1.6 \times 10^{-19} \times 12000}{9.1 \times 10^{-31}}} \quad \checkmark 1 \\
 &= \sqrt{4.2198 \times 10^{15}} \checkmark 1 \\
 &= 6.496 \times 10^7 \text{ m/s}
 \end{aligned}$$

- (iii) - Detecting fault in metals or other structures.  
 - Controls quality of manufacturer items e.g. tyres, thickness of sheets, papers e.t.c.  
 - Analysis of gem stones. (Any one-1mk)

7. (a) A – cathode      B – Anode      C – Cooling fins  
 (b) (i) increase the p.d at the anode (B)  
 (ii) : increase the cathode heater current  
 (c) Tungsten:- It has a high melting point so the heat produced will not melt it easily  
 (d) Copper – it is used to cool/conduct heat away from the anode  
 (e) So that the electrons do not collide with gas molecules which could result in loss of energy.  
 (f) (i) Detecting fracture in bones  
 (ii) Detecting flaws in metals ✓

8. a) i) Name the part labeled C and state its function  
 ■ C is the cathode.  
 ■ It produces electron thermionically  
 iii) (Take electric charge  $e = 1.602 \times 10^{-19} \text{ C}$ , planks constant  
 $h = 6.63 \times 10^{-34} \text{ Js}$ , and speed of light  $c = 3.0 \times 10^8 \text{ m/s}$ )  
✓  $\text{Energy of X-rays } f = 8.01 \times 10^{16}$   
 $\lambda = \frac{5}{100} \times 100 \text{ kV} \times 1.602 \times 10^{-19} \text{ C}$   
 $6.63 \times 10^{-34}$   
 $= 8.01 \times 10^{-16} \text{ J} = 1.208 \times 10^{18} \text{ Hz}$   
 $\lambda = \frac{c}{f} = \frac{3.0 \times 10^8 \text{ m/s}}{1.208 \times 10^{18} \text{ Hz}}$   
 $= 2.483 \times 10^{-10} \text{ m}$


- iv)- They penetrate matter  
 -They obey properties of electromagnetic waves  
 ○ Diffraction  
 ○ Reflection  
 ○ Obey inverse square law  
 v) -Used to detect defects in metals in industries  
 -Used to sterilize medical equipment.

9. i) K- X      ii) X = 88      Y= 288

### Radioactivity

1. a) Radioactive decay is the spontaneous random emission of particles from the nucleus of an unstable nuclide  
 (b) There are 7 half lives ( $t_{1/2}$ )  
 $7t_{1/2} = 49 \text{ days}$   
 $t_{1/2} = \frac{49}{7}$   
 $= 7 \text{ days}$





- (ii) During the first half cycle, the diode is forward biased so it conducts.  
 - Current flows through  $R_L$  building a voltage which decreases as the first half cycle comes to an end.  
 - During the second half cycle, the diode is reverse biased so it does not conduct.

(b) (i)  $y = 238 - 4(1) = 242$

$X = 2g$

(ii)  $\frac{120}{20} = 6$  half lives

$0.03125 \times 26 = 2g$

(iii) They are deflected by both electric and magnetic fields

(iv) Alpha particles are heavy (massive)

(v) - The sheets are brought in turns between radioactive source and the counter.

- The count rate is a measure of the thickness of the metal sheet.

3. a) Spontaneous disintegration of unstable atoms in order to gain stability

b) i)  $a = 236 - 91 = 145$

ii)  $b = 92 - 38 = 54$

c) radioactive substances are harmful to the body when ingested

d) i) Negative

ii) A - Beta radiation

C - Alpha radiation

iii) C - more massive than A

e) i)  $A = 233 - 8 = 225$

ii)  $Z = 90 - [(2 \times 2) + (2x - 1)]$   
 $= 90 - (4 - 2)$   
 $= 90 - 2 = 88$

f) - a beta source is placed on one side of a moving sheet of paper and a G.N detector on the other side

- If the material is too thin, the count rate at the detector will be too high and vice versa

4. (i)  $S - 210$  ✓  $T - 206$

(ii) The splitting of a heavy nuclide to lighter particles (fission process)

5. State what type of radioactive decay this is. - Alpha decay

a)  $X \dots 4$   $Y \dots 2$

6. Long radio waves, AM radio waves, T.V and FM Radio waves, short Radio waves, infra red radiation, red-light, Uv radiation and X-rays.

7. No. of half lives =  $380 = 10$

$N = N_0 \left(\frac{1}{2}\right)^{t/T}$  ✓

$\frac{380}{38} = \left(\frac{1}{2}\right)^{10} = 1$  ✓

8. (a) Time taken for the activity of a sample of a radioactive material to reduce to half of the original value

(b) (i) S - scale - simple and uniform / consistent

p - Plotting at least 4 points correct


C - Line must pass through at least 3 points

(ii) -Half-life  $319 \pm 0.1$  days (1mk)

-Readings -off from the graph clearly



- ii) Now displace one end of the meter rule slightly on a horizontal plane so that when released it oscillates about a vertical axis as in the figure below.



- iii) Measure the timer for 10 oscillations and record the value in the table provided below  
 c) i) Repeat the procedure in (b) above for the values of  $d$  shown in the table (set the values of  $d$  by adjusting the positions of the loops in steps of 5cm on both sides)  
 ii) Complete the table

| D (cm) | D (m) | $1/d^2$ ( $M^{-2}$ ) | Time for 10 oscillations | Period T (s) | $T^2$ ( $S^2$ ) |
|--------|-------|----------------------|--------------------------|--------------|-----------------|
| 80     |       |                      |                          |              |                 |
| 70     |       |                      |                          |              |                 |
| 60     |       |                      |                          |              |                 |
| 50     |       |                      |                          |              |                 |
| 40     |       |                      |                          |              |                 |
| 30     |       |                      |                          |              |                 |
| 20     |       |                      |                          |              |                 |


- d) i) On the grid provided, plot a graph of  $T^2$  (y – axis) against  $1/d^2$  ( $M^{-2}$ )  
 ii) Determine the slope of your graph  
 iii) Given that  $T^2 = \frac{16K^2}{5d^2}$  where  $K$  is a constant. Use the graph to determine the value of  $K$

## QUESTION 2

This question has two parts **A** and **B**. Answer both parts.


**You are provided with the following:**

- A lens and a lens holder
- A candle
- Object consisting of a hole 2cm in diameter and parallel wires 1.5cm apart in a stiff card.  
(See diagram below)
- A screen
- A meter rule



**Proceed as follows:-**

- a) Illuminate the object with the candle flame.  
 b) Arrange the object, lens and screen in line as shown in figure 3 below



- c) Measure the distance,  $d$ , between the two parallel wire that acts as the object  
 $d = \dots\dots\dots$  cm
- d) Adjust the lens,  $u$  to 80cm
- e) Move the screen until a clear image is formed on it.
- f) Measure the distance,  $X$ , of the image, making sure that what you measure is an image corresponding to the previous reading,  $d$ .

Record these values in the table below:-

- g) Repeat your readings of  $x$  with  $u = 70, 60, 50, 40$  and  $30$ cm and complete the table

| U (cm) | 80 | 70 | 60 | 50 | 40 | 30 |
|--------|----|----|----|----|----|----|
| X(cm)  |    |    |    |    |    |    |
| d/ x   |    |    |    |    |    |    |

- h) i) On the grid provided plot a graph of  $u$  (y – axis) against  $d/x$
- ii) I. Determine the slope,  $S$  of the graph  
 II. Find the intercept on the  $u$  – axis

**PART B**

*You are provided with the following:-*

- A jockey  $J$
- An ammeter
- A voltmeter
- A switch
- 6 connecting wires,  $Z$  with crocodile clips on one end
- A resistor wire labeled  $XY$  mounted on a piece of wood having a millimeter scale
- 2 new dry cells

*Proceed as follow:*

- 1) i) Connect the circuit as shown below:




FIG. 4

- ii) Close the switch and note the voltmeter and ammeter readings when  $XJ = 10$ cm

iii) Repeat procedure (i) and (ii) above with  $XJ = 20 \text{ cm}$  and enter in the table 3 as below:  
\*KKC\*

Table 3:-

| Length XJ (cm) | P.d.V . (v) | Current, I (A) |
|----------------|-------------|----------------|
| 10             |             |                |
| 20             |             |                |


J) Given that  $\log I = n \log V + \log k$ , where  $k$  and  $n$  are constants, determine the values of  $k$  and  $n$

### KAKAMEGA EAST DISTRICT

1. You are provided with:

- Mass  $M$
- One 100g mass
- Metre rule
- Cotton thread (3 –pieces each about 30cm long)
- Retort stand and clamp
- 250cm<sup>3</sup> glass beaker
- 200cm<sup>3</sup> of water

- (a) (i) The loops of thread on solid  $M$  and the 100g mass  
(ii) Suspend the metre rule on the clamp from the 50cm mark  
(iii) Hang mass  $m$  from the mark. Balance the metre rule using the 100g mass (see fig. 1 below)



- (iv) Measure the distance  $X$  and  $X_1$  from the 50cm mark  
(v) Repeat the procedures for the values of  $X$  indicated in the table below:

| X(cm) | X <sub>1</sub> (cm) | X <sub>2</sub> (cm) | X <sub>1</sub> - X <sub>2</sub> (cm) |
|-------|---------------------|---------------------|--------------------------------------|
| 45    |                     |                     |                                      |
| 40    |                     |                     |                                      |
| 35    |                     |                     |                                      |

|    |  |  |  |
|----|--|--|--|
| 30 |  |  |  |
| 25 |  |  |  |
| 20 |  |  |  |

- (b) (i) Repeat steps (a) (iii) to (a) (iv) above, but this time, keep mass **M** totally immersed in water.  
Record distance **X<sub>2</sub>** required to balance the 100g mass in the table above.  
(ii) Complete the table for the values of (**X<sub>1</sub>**- **X<sub>2</sub>**)
- (c) (i) Plot a graph of **X<sub>1</sub>** (Vertical axis) against (**X<sub>1</sub>**-**X<sub>2</sub>**) on the grid provided  
(ii) Determine the slope **S** of your graph  
(iii) What physical property does the slope, **S** represent?  
(iv) Given that the density of water is 1000kg/m<sup>3</sup>, determine the density of mass, **M**
- (d) (i) Using the apparatus you were given, determine the mass of your metre rule  
(ii) Draw a diagram of the set-up of the apparatus you have used to work out (d) (i) above


2. **You are provided with the following:-**

- Two dry cells (size D)
- 1M long nichrome wire (S.W.G 28) labelled AB
- Ten connecting wires, one of length 70cm having a jockey
- A micrometer screw gauge
- A torch bulb
- An ammeter (0 – 10A)
- A voltmeter (0 – 3V)
- Switch

**Proceed as follows:-**

- (a) (i) Set-up the circuit below:

fig 2



- (ii) With the jockey (**J**) at **X** i.e **L = 100cm**, record the voltmeter reading, and the ammeter reading **I**. Repeat the readings for **L = 80, 60, 40, 20, and 0cm** and enter your results in the table below:

|               |     |    |    |    |    |   |
|---------------|-----|----|----|----|----|---|
| L(cm)         | 100 | 80 | 60 | 40 | 20 | 0 |
| P.d V(Volts)  |     |    |    |    |    |   |
| Current I (A) |     |    |    |    |    |   |

- (b) (i) Plot a graph of p.d **V** (y-axis) against the ammeter reading, **I**  
(ii) Determine the slope of your graph when **V = 0.5volts**  
(iii) What physical quantity does the slope in (ii) above represent?
- (c) (i) Given the apparatus above, draw a diagram of the circuit you would use to determine the current through the resistance wire, **AB** and the p.d across it when the cells are now in series  
(ii) Set-up the circuit you have just drawn and record the current **I** and p.d **V** when **L = 100cm**  
(iii) Using the micrometer screw gauge, measure the diameter **d** of the wire, **AB**  
**d = \_\_\_\_\_ mm**
- (iv) Calculate the quantity, **S**, given that:

$$S = \frac{\pi V d^2}{4 I L} \quad (\text{Take } \pi = 3.14)$$

Give the units of S

### MIGORI/NYATIKE DISTRICT

**Q.1** You are provided with the following apparatus:-

- 600 ml beaker.
- A measuring cylinder (100ml).
- Plasticine.
- A meter rule.
- Distilled water supplied in a 500 ml wash bottle.
- Candle.
- Screen.
- Tissue paper, 30 cm ruler

**Proceed as follows:-**

a i) Add 310 ml of water to the beaker and obtain **h**, the height in cm of the water above the base of the beaker.


**h** = ..... cm

ii) Calculate an approximate value of **R**, the internal radius in cm from the formula.

$$R = \sqrt{\frac{100}{h}}$$

b) i) **Fill** the beaker with more water and set up the apparatus as shown in figure 1. The vessel should be placed at around 50 cm mark on the scale.

**Figure 1**



ii) Starting with 10R, adjust the screen until you see a sharp bright vertical line.

iii) **Record** the distance **V** of the image in **table 1** whose image is sharpest. **Repeat** the experiment with values of **u** between 10R and 3R cm.

| <b>u</b>      | 3R | 4R | 5R | 6R | 7R | 8R | 9R | 10R |
|---------------|----|----|----|----|----|----|----|-----|
| <b>u (cm)</b> |    |    |    |    |    |    |    |     |
| <b>v (cm)</b> |    |    |    |    |    |    |    |     |

c) Plot the graph of **u** against **v**.

d) From the graph determine.

i) **v<sup>1</sup>** the value of **v** for which **v = u**.

ii) **u<sup>1</sup>** the value of **u** for which **u = 2v**.

e) Calculate **f**, the effective focal length of the 'lens' from the formula.

$$f = \frac{u^1 + v^1}{5}$$

f) Give the approximate value of **R/f**.



**Q2. Part A.**

**You are provided with the following apparatus:-**

- Boiling tube.
- A cork with a hole and a thermometer to fit in it.
- Complete retort stand.
- Source of heat.
- Some distilled water in a beaker.
- Stop-watch.
- Tissue paper.
- 250ml beaker.

**Proceed as follows:-**

- Heat** the water in a beaker until it boils. It is essential that the water is kept boiling throughout the experiment.
- Clamp** the boiling tube as shown in **figure 2** making sure that the tube does not touch the base of the retort stand.



- Using a 250 ml beaker, **transfer** some boiling water into the boiling tube making sure that some space is left. **Plug** the tube with the cork that carries the thermometer as shown in **figure 3**.
- Starting with the temperature of 80°C, **note** the temperature of the water every **1 minute**
  - Tabulate** your results in a table as shown below:-

|                         |   |   |   |   |   |   |   |   |   |   |    |
|-------------------------|---|---|---|---|---|---|---|---|---|---|----|
| <b>Time t (min)</b>     | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| <b>Temperature (°C)</b> |   |   |   |   |   |   |   |   |   |   |    |

- Plot a graph of temperature  $\theta$  (°C), against  $t$ , min.
- Find the gradient  $\frac{d\theta}{dt}$  of your graph at the temperature of 70°C
- Find the rate of heat loss,  $R$ , at the temperature of 70°C given that  $R = K \frac{d\theta}{dt}$  where  $K = 1.23 \times 10^4$

## Q.2 Part B


**You are provided with the following:-**

- Two New identical dry cells (size D)
- Nichrome wire labelled AB, mounted on a metre rule.
- An ammeter (0 – 1.5A) or (0 – 2.5A).
- A voltmeter 0 – 5V.
- A cell holder.
- 8 connecting wires at least 4 with crocodile clips.
- Jockey.
- A switch



**Proceed as follows:-**

a) **Connect** the circuit as shown in **figure 4** below



b) **Connect** the end **A** and **B** where **AB = 100 cm** across the terminals as shown in figure 4.

**Close** the switch and **measure** both **I** and **p.d., v** across the wire **AB**.

- i) Current **I** = ..... A
- ii) p.d. **V** = ..... V
- iii) **Measure** the e.m.f. of the cells, **E** = ..... V
- iv) Given that equation **E = V + Ir**, determine the internal resistance of the two cells.

### SOTIK DISTRICT

1. **You are provided with the following apparatus:-**

- Micrometer screw gauge
- Vernier caliper
- Water in a beaker 1000ml(should be ½ full)
- Long test-tube
- Some dry sand
- Spatula
- Millimetre scale marked on a paper strip
- Some cellotape
- 6 ball bearings

**Proceed as follows:-**

(i) Measure and record the diameter **d** of one ball bearing using micrometer screw gauge

**d** = \_\_\_\_\_ cm

(ii) Determine the volume **V** of the spherical ball bearing

**V** = \_\_\_\_\_ cm<sup>3</sup>


(iii) Measure the inside diameter **d** of the test-tube using vernier caliper. Record it below:

**d** = \_\_\_\_\_ cm

(iv) Find the cross-section area **A** of the test tube

**A** = \_\_\_\_\_ cm<sup>2</sup>

- (b) (i) Place the millimeter scale along the height of the test tube so that the zero is at the bottom
- (ii) Place the test-tube in the water carefully and add sand bit by bit until it floats while vertically upright in the water as shown:-



(iii) Note and record the height  $h_0$  of water level by use of attached millimeter scale

$$h_0 = \underline{\hspace{2cm}} \text{ cm}$$

(c) Add one ball bearing into the tube, note and record the new level  $h$  in the table of results below:

(d) Repeat step (c) with **two, three, four, five & six** ball bearings and record their corresponding  $h(\text{cm})$ .

Compute values of  $h-h_0(\text{cm})$  in the table below:-

| No. of Ball Bearing (N) | Floating level $h(\text{cm})$ | $h - h_0(\text{cm})$ |
|-------------------------|-------------------------------|----------------------|
| 1                       |                               |                      |
| 2                       |                               |                      |
| 3                       |                               |                      |
| 4                       |                               |                      |
| 5                       |                               |                      |
| 6                       |                               |                      |

(e) Plot a graph of  $h-h_0(\text{cm})$  against the number of ball bearings (N)

(f) Determine the slope  $S$ , of the graph

(g) Relative density  $P_s$ , of ball bearing is given by:

$$P_s = \frac{SA}{V}. \text{ Find } P_s$$


## 2. PART I

You are provided with the following apparatus:-


- Rectangular glass block
- Four optical pins
- Plain paper
- Soft board
- Piece of cellotape

**Proceed as follows:-**

- (a) (i) Use cello tape provided to hold the sheet of plain paper on the soft board
- (ii) Place the glass block on the middle of plain paper and with a sharp pencil, draw its outline ABCD as shown:-



- (b) (i) Construct the normal on side AB, but close to A. Use protractor and ruler to draw an incident ray with an angle of incidence =  $10^\circ$
- (ii) Insert pins  $P_1$  and  $P_2$  along the path drawn. Viewing through the glass block on side CD, locate  $P_3$  and  $P_4$  such that  $P_3$   $P_4$  appear in line with images of  $P_1$  and  $P_2$
- (iii) Produce  $P_1P_2$  to obtain a lateral displacement as shown in the figure below:-  
Measure angle of refraction  $r$ , and lateral displacement



- (c) Repeat steps (b)(ii) and (b)(iii) for angles of incidence  $i = 20^\circ, 30^\circ, 40^\circ, 50^\circ$  and  $60^\circ$ . Tabulate your results as shown in the table:-

**Note:** (You must handover your workings on the plain paper with the question paper after the session)

| Angle of incidence $i$ | $20^\circ$ | $30^\circ$ | $40^\circ$ | $50^\circ$ | $60^\circ$ |
|------------------------|------------|------------|------------|------------|------------|
| Angle of refraction    |            |            |            |            |            |
| Lateral displacement   |            |            |            |            |            |


- (d) Plot a graph of lateral displacement,  $d$ , against angle of refraction  $r$

## 2. **PART II**

**You are provided with the following:-**


- An ammeter (0- 1A)
- A voltmeter (0-2.5V or 0-5V)
- Two dry cells
- A mounted resistance wire
- Eight connecting wires, two with crocodile clips
- A three volts torch bulb in a bulb holder
- A cell holder
- A switch
- A jockey or a crocodile clip

(a) Set-up the apparatus as shown below:-



(b) With the jockey or crocodile clip at C, 30cm, 50cm and 70cm, record their corresponding  $V_1$ ,  $V_2$  and  $V_3$

(c) Replace the voltmeter with a torch bulb and an ammeter. Connect in series as shown in the circuit diagram:-



(d) Read and record the ammeter reading  $I_1$ ,  $I_2$  and  $I_3$  for the corresponding values of lengths:

$l_1 = 30\text{cm}$ ,  $I_1 = \underline{\hspace{2cm}}$

$l_2 = 50\text{cm}$ ,  $I_2 = \underline{\hspace{2cm}}$

$l_3 = 70\text{cm}$ ,  $I_3 = \underline{\hspace{2cm}}$

(e) (i) Determine voltage values across the bulb for lengths 30cm, 50cm and 70cm given that  $V = 0.025L$

(ii) Determine the average resistance of bulb during the experiment

## UGENYA/UGUNJA DISTRICT

### Question 1


#### Part 1

1. You are provided with the following apparatus:-

- Clamp, boss and stand
- Optical pin fixed on a piece of cork
- Wire (length 30cm)
- Stop watch

**Proceed as follows:**

- (a) Bend the wire in the middle so that the angle formed is  $20^\circ$ .  
 (b) Set up the apparatus as shown in figure 1 below



- (c) (i) Displace the wire horizontally and allow it to swing freely. Record time  $t$  for 10 complete oscillations.  
 (ii) Calculate the number of oscillations per second ( $f$ )  $f = 10/t$   
 (d) Repeat the procedure above for the other given angles on table 1

| Angle $\theta^\circ$ | Time( t)for 10 oscillations | $f = 10/t$ | $f^2 \text{ (Hz)}^2$ | $\theta/2$ | $\text{Cos } \theta/2$ |
|----------------------|-----------------------------|------------|----------------------|------------|------------------------|
| 20                   |                             |            |                      |            |                        |
| 40                   |                             |            |                      |            |                        |
| 60                   |                             |            |                      |            |                        |
| 80                   |                             |            |                      |            |                        |
| 100                  |                             |            |                      |            |                        |

- (e) On the grid provided plot a graph of  $f^2$  against  $\cos \theta/2$   
 (f) Find the gradient of the graph, stating its units

## **Part II**

*You are provided with the following:-*


- A rectangular glass block
- 1 white plain sheet of paper
- 4 optical pins
- 4 thumb tacks
- A soft board

You are required to determine the refractive index of the glass block

### **Proceed as follows:**

- (i) Fix the sheet of paper provided on the soft board using the thumb tacks  
 (ii) Place the rectangular glass block on the plain paper and trace its outline  
 (iii) Remove the block and draw a normal line at point O. Draw a line incident at an angle of  $15^\circ$  as shown in figure 2 below  
 (iv) Stick 2 Pins  $P_1$  and  $P_2$  along the ray in and by looking through the glass block from the opposite side, stick two other pins  $P_3$  and  $P_4$  in line with the images of  $P_1$  and  $P_2$ . Remove the glass block  
 (v) With O as the centre, draw a circle of radius 5cm, to cut both the ray incident and the refracted ray at L and M respectively.  
 (vi) Using a set square, draw the perpendicular LN and MN to the normal
- (c) Repeat the procedure (iii) to (iv) for other values of  $30^\circ$  and  $45^\circ$  as shown in figure 2

Fig. 4



(c) Make LN and MN and record your values in the table 2 below

| Angle of incidence | LN(mm) | MN (mm) | Refractive index $\frac{LN}{MN}$ |
|--------------------|--------|---------|----------------------------------|
| 15°                |        |         |                                  |
| 30°                |        |         |                                  |
| 45°                |        |         |                                  |

(d) Average value of refractive index = \_\_\_\_\_


**Question 2**

You are provided with the following apparatus:

- Bi convex lens
- Lens holder
- Screen S
- Candle
- Screen with cross wire X
- Meter rule
- Plasticine

**Proceed follows:-**

(a) Arrange the apparatus as shown in the diagram below:-



(Ensure the metre rule is firmly held on the bench by plasticine and the flame and crosswire is at a horizontal level with the centre of the lens throughout the experiment)

(b) At a point  $a = 94\text{cm}$  from the screen X, obtain a sharp image of the crosswire on screen S by adjusting the position of the lens along the metre rule. Record the values of b and c as shown in the

diagram.

- (c) Adjust the position of screen **S** along the metre rule for **a** = 90cm, obtain a sharp image of the crosswire on screen **S** by adjusting the position of the lens. Record the corresponding values of **b** and **c**


Repeat the procedure (a) and (b) above for other values of **a** = 94cm, 90cm, 86cm, 82cm, 78cm, 74cm and complete the table below

| a cm | b cm | C cm | d = a/c |
|------|------|------|---------|
| 94   |      |      |         |
| 90   |      |      |         |
| 86   |      |      |         |
| 82   |      |      |         |
| 78   |      |      |         |
| 74   |      |      |         |

- (d) Plot a graph of **b(cm)** (y-axis) against **d**  
 (e) Determine the slope **S** of the graph  
 (f) Given that the equation of the graph is  $b = \frac{(100)d}{L}$  determine the value of **L** from the equation above  
 (g) What does **L** represent?

### PART II

Arrange the apparatus as shown below:



- (a) Obtain a sharp image of flame on screen **S** by adjusting the position of screen **S**. Record the image distance,  $v = \underline{\hspace{2cm}} \text{cm}$   
 (b) Determine the constant  $m = \frac{v}{u}$   
 (c) What does **m** represent?

## NDHIWA DISTRICT


### Question 1

You are provided with the following apparatus:-

- A metre rule
- A wire of length at least 100cm
- A retort stand, boss and clamp
- A stop watch or stop clock
- A micrometer screw gauge
- An overflow can, a beaker atleast 50ml
- A 50ml measuring cylinder
- A piece of thread about 30cm
- Water in a 250ml beaker
- Two pieces of wood
- Mass labeled **m**

**Proceed as follows:-**


- (a) (i) Fill the overflow can with water to overflowing level and then allow it to drain.  
 (ii) Immerse the mass **m** into the can. Collect the overflow in a beaker as shown below in **fig 1**:-



- (iii) Using the measuring cylinder provided determine the volume **V** of the water collected in the beaker  $V = \underline{\hspace{2cm}} \text{ cm}^3$

- (iv) Calculate **I** given that  $I = \frac{10^6 m}{V}$  where  $m = 0.30\text{kg}$

- (b) Set-up the apparatus as shown in **figure 2** below. Ensure that the wire is free of kinks and the end tied to the hook is firm and the hook does not move.



- (c) Adjust the length **L**, of the wire so that  $L = 70\text{cm}$ , Give the mass **m**, a slight twist such that when



released it oscillates about the vertical axis as shown by the arrows in **figure 2** measure the time for twenty oscillations and record in table 1.

(d) Repeat the procedure in (c) above for other values of **L** shown in table 1. Complete the table

**Table 1:**

|                                     |    |    |    |    |    |    |
|-------------------------------------|----|----|----|----|----|----|
| <b>Length L(cm)</b>                 | 70 | 60 | 50 | 40 | 30 | 20 |
| <b>Length L (m)</b>                 |    |    |    |    |    |    |
| <b>Time for 20 oscillations (s)</b> |    |    |    |    |    |    |
| <b>Period T (s)</b>                 |    |    |    |    |    |    |
| <b>T<sup>2</sup>(s)<sup>2</sup></b> |    |    |    |    |    |    |

(e) On the grid provided, plot the graph of **T<sup>2</sup>(S<sup>2</sup>)** against **L (m)**

(f) Measure the diameter **d** of the wire **d = .....metres**

(g) (i) Determine the slope of the graph

(ii) Given that  $T^2 = \frac{32\pi^2 L}{Gd}$  where **G** is a constant, use the graph to determine the value of **G**

**Question 2.**

**You are provided with the following apparatus:-**


- Two new dry cells
- A resistor labelled **Q**
- A wire **AB** mounted on a millimeter scale
- 6 connecting wires with crocodile clips on one end of atleast three
- A voltmeter
- An ammeter
- A switch

**Proceed as follows:-**

(a) Connect the apparatus provided as shown in **figure 3** below:-

(b)

**fig. 3**



(i) Take the voltmeter reading when the switch **S** is open. **V<sub>1</sub> = .....Volts**

(ii) Close the switch **S**, and take the voltmeter reading **V<sub>2</sub>** and the ammeter reading **I**


**V<sub>2</sub> = .....Volts**

**I = .....Amperes**

(iii) Calculate the quantity  $P = \frac{V_1 - V_2}{I}$

(c) Set-up the circuit as shown in **figure 4:-**


fig. 4



- (i) Take the voltmeter reading  $V$  and the ammeter reading  $I$
- (ii) Determine the resistance  $R$  of  $Q$  given that  $R = \frac{V}{I}$

**N/B** :-The circuit must be left open when no reading is taken

- (c) Set-up the circuit shown in *figure 5*



- (d) Move the crocodile clip along the wire  $AB$  to a point such that  $L = 100\text{cm}$ . Note the voltmeter reading and record in table 2.
- (e) Repeat (d) above for values of  $L = 80\text{cm}, 60\text{cm}, 40\text{cm}, 20\text{cm}$  and  $0\text{cm}$ , tabulate your results

**Table 2**

| Length $L$ (cm)                         | 100 | 80 | 60 | 40 | 20 | 0 |
|---|-----|----|----|----|----|---|
| $\frac{1}{L}$ ( $\frac{1}{\text{cm}}$ ) |     |    |    |    |    |   |
| Voltmeter reading (V)                   |     |    |    |    |    |   |
| $\frac{1}{V}$ ( $\frac{1}{\text{V}}$ )  |     |    |    |    |    |   |

- (f) Plot the graph of  $\frac{1}{V}$  against  $\frac{1}{L}$
- (g) Find the slope of the graph


### MUMIAS DISTRICT

- 1. You are provided with the following apparatus:

- Ammeter (0-2.5A)
- A voltmeter (0 –5 V)
- A straight wire  $AB$  1.0m long mounted on a millimeter scale
- Two jockeys
- Connecting wires
- Micrometer screw gauge (to be shared)
- Two dry cells (size D) a cell holder and a switch
- A cell holder
- Switch

**Procedure:**

- (a) Using the micrometer screw gauge, determine the diameter **d** of the wire  
Set up the apparatus as shown in fig. 1



**Fig. 1**

With both jockeys set at  $L=10\text{cm}$  from A, measure the current **I** through the wire and voltage (**V**) across. Repeat these corresponding values of current **I** and voltage (**V**) in the table below:

|                           |    |    |    |    |    |    |    |    |    |     |
|---------------------------|----|----|----|----|----|----|----|----|----|-----|
| Length (cm)               | 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 | 90 | 100 |
| Length (m)                |    |    |    |    |    |    |    |    |    |     |
| Current I(A)              |    |    |    |    |    |    |    |    |    |     |
| Voltage V(V)              |    |    |    |    |    |    |    |    |    |     |
| $R = \frac{V}{I}(\Omega)$ |    |    |    |    |    |    |    |    |    |     |

- (i) Using the values in table I plot a graph of Resistance  $R(\Omega)$  against length  $L(\text{m})$   
(i) Determine the slope of the graph  
(ii) Determine the resistivity of the material of the wire given that;


$$R = \frac{4\rho h}{\pi d^2}$$

**2. You are provided with the following apparatus:**

- A spring balance, (0-2.5N)
- A retort stand, a boss and a clamp
- A wooden block
- One 50g mass
- A beaker and some water
- Five pieces of strings
- A vernier calipers (to be shared)

**Procedure**

- (a) Measure the length of the wooden block using a vernier calipers  $L =$
- (b) From the lower end of the wooden block, mark levels on the wood at intervals of 1cm for 5cm as shown. Mark these levels using pieces of thread
- (c) Set-up apparatus as shown below in fig 2.



(d) Lower the mass into the empty beaker provided. The mass should be as low as possible by **not** touching the base of the beaker.

Record the reading  $W_1$  of the spring balance

- (i) Slowly pour water into the beaker until the mass is completely immersed. Continue adding Water into the beaker until when the level reaches mark (thread) indicating 1cm on the wooden block. Read and record the value  $W_2$  of the spring balance (Don't disturb the wood and mass when pouring water into the beaker)
- (e) Repeat procedures in **d(ii)** above by adding water up to the line indicating 2cm, 3cm, 4cm and 5cm. Denote these lines by values **h**. Complete the table for different values of **h**

|                       |   |   |   |   |   |
|-----------------------|---|---|---|---|---|
| h(cm)                 | 1 | 2 | 3 | 4 | 5 |
| $W_2$ (N)             |   |   |   |   |   |
| $F = (W_1 - W_2)$ (N) |   |   |   |   |   |

- f. (i) Plot a graph of F against h
- (ii) Determine the gradient of the graph
- (iii) Given that  $F = \frac{Dgh}{100} + C$  Whereby **D** and **C** are constants,  $g = 10\text{N/kg}$   
use the graph to determine the values of D and C
- (iv) Determine the constant W for the wooden block given that  $W = DL$

**KISUMU WEST DISTRICT**


**Question 1**

- You are provided with the following:-
- A voltmeter
- An ammeter
- Two dry cells and a cell-holder

- A switch
- A potentiometer or Rheostat
- Six connecting wires

**Proceed as follows:-**

(a) Set-up the circuit as shown in figure 1 below:-



- (b) With the switch open, record the reading  $E_0$  of the voltmeter  
 $E_0 = \dots\dots\dots$  volts
- (c) Close the switch. Adjust the voltmeter to read 0.5volts using the potentiometer or Rheostat. Measure the corresponding current with the ammeter and record in table 1
- (d) Repeat the procedure in (c) using different voltmeter readings provided in tale 1 below, each time recording the corresponding ammeter reading. Complete the table

**Table 1**

|                                    |     |     |     |     |     |     |
|------------------------------------|-----|-----|-----|-----|-----|-----|
| Voltage (V)<br>Volts               | 0.5 | 1.0 | 1.5 | 2.0 | 2.5 | 2.7 |
| Current (I) A                      |     |     |     |     |     |     |
| $\frac{I}{I}$ ( $A^{-1}$ )         |     |     |     |     |     |     |
| $\frac{R}{I} = \frac{V}{I}$ (Ohms) |     |     |     |     |     |     |

- (e) Plot a graph of  $1/I$  (y-axis ) against R  
 (f) Determine the slope S of the graph  
 (g) (i) Find  $R_0$ , the value of R when  $1/I = 0 A^{-1}$   
 (ii) Evaluate  $1/S - E_0$

**Question 2.**


**You are provided with the following apparatus:**

- A thin lens marked L
- A lens holder
- An office pin
- An optical pin
- A white screen
- A metre rule

- Some plasticine

**Proceed as follows:-**

(a) Set-up the apparatus as in figure 3 below



Focus a distant object, say a window, by letting the lens **L** position invariant. Meanwhile slide the white screen back and forth till a sharp image of the distant object; window, appears clearly on the white screen.

Measure the distance, **f**, between the lens **L** and the white screen


**f** = .....cm

(b) Set-up the apparatus as shown in **figure 4** below. Use the office pin as an object and place it at a distance less than **f**; of the lens **L**.

Mount the office pin on plasticine to enhance its proper visibility through the lens **L**

Take the optical pin and use it as a search pin. Find a position of no parallax between the search pin and the image through the lens **L**. Measure the distance between the office pin and the image, **V** in centimeters

**fig 4**



(c) Describe the image characteristics

(d) Put the object pin near the lens **L**, that is, at **u = 3cm**. Observe through the lens **L** such that there is no -parallax between the object and the image, i.e. when **V= 3cm**

(e) Repeat the procedure using at least six different values of **u**. complete table 2 below:-

**Table 2:**

|                      |     |     |     |     |     |      |      |      |
|----------------------|-----|-----|-----|-----|-----|------|------|------|
| Object distance, Ucm | 3.0 | 4.0 | 6.0 | 7.0 | 8.0 | 10.0 | 11.0 | 12.0 |
| Image distance , Vcm |     |     |     |     |     |      |      |      |
| $M = \frac{V}{U}$    |     |     |     |     |     |      |      |      |

- (f) Plot a graph of **U (y-axis)** against **V**  
 (g) (i) from your graph, find the value of **V when  $u \approx f$**   
 (ii) State the application of this practical in real life experience

**TRANS- NZOIA WEST DISTRICT**


**PART A**

1. You are provided with the following:

- A complete retort stand
- An optical pin
- A concave mirror and a holder or a lump of plasticine
- A cork
- A candle
- A screen
- A metre rule

You are required to estimate the focal length of the mirror. Arrange the apparatus as follows in figure 1 below:

Figure 1




a) By adjusting the clamp on the stand, move the object pin up and down until the inverted image and pin itself appear to coincide (use – no- parallax method). Measure the distance

$h = \dots\dots\dots$  cm

b) Calculate the value **f** given that

$f = \frac{h}{2} \dots\dots\dots$

c) Arrange the apparatus as shown in figure 2 below



d) Place the candle at a distance  $u = 22\text{cm}$  from the mirror. Move the screen along the mirror rule until a sharp image is formed on the screen. Measure and record the image distance  $V$

e) Repeat the experiment with the values of  $u = 24\text{cm}$ ,  $26\text{cm}$ ,  $28\text{cm}$  and  $30\text{cm}$ . Record your values in table 1 below

| Object distance $u$ (cm) | Image distance $V$ (cm) | Magnification $M = v/u$ |
|--------------------------|-------------------------|-------------------------|
| 22                       |                         |                         |
| 24                       |                         |                         |
| 26                       |                         |                         |
| 28                       |                         |                         |
| 30                       |                         |                         |

f) Plot a graph of magnification,  $m$  (y- axis) against image distance,  $v$

g) Given that  $M = \frac{V}{f} - 1$ . Determine the focal length,  $f$


## **PART B**

*You are provided with the following apparatus:-*

- A lens
- A lens holder
- A white screen
- A metre rule
- Candle

### **Procedure**

i) Set up the apparatus as shown in figure 3 below



j) Starting with  $u = 30\text{cm}$  adjust the position of the screen to obtain a sharp image of the candle.

Record the value of  $V$  in the table 2 below

k) Repeat the procedure above for  $u = 20\text{cm}$  and complete the table 2 below

Table 2



| (U) cm | V (cm) | Uv (cm <sup>2</sup> ) | U+V (cm) |
|--------|--------|-----------------------|----------|
| 20     |        |                       |          |
| 30     |        |                       |          |

ii) Given that the focal length of the lens satisfies the equation,  $f = \frac{UV}{U+V}$


Determine the average value of the focal length  $f$

**Q2. You are provided with the following:**

- 2 dry cells
- A cell holder
- A torch bulb
- A bulb holder
- A voltmeter
- An ammeter
- A switch
- 8 connecting wires, one with a jockey and some with crocodile clips
- A wire attached on a metre rule or millimeter scale

**Proceed as follows :-**

a) i) Set up the circuit as shown in figure 4 below



ii) With the jockey at P (i.e.  $L = 100$  cm), take the voltmeter reading ( $v$ ) and ammeter reading ( $A$ )

Record  $V$  and  $I$  readings in table 3 below

iii) Repeat the procedure to take readings for  $L = 80$  cm,  $60$  cm,  $20$  cm and  $0$  cm respectively

Record your readings in table 3 below:

**Table 3**

|        |     |    |    |    |    |   |
|--------|-----|----|----|----|----|---|
| L (cm) | 100 | 80 | 60 | 40 | 20 | 0 |
| V (v)  |     |    |    |    |    |   |
| I (A)  |     |    |    |    |    |   |

iv) What changes do you observe on the bulb as  $L$  decreases from **P** to **Q** ?

v) Plot a graph of voltage ( $v$ ) (y- axis) against current ( $I$ )

vi) Determine the slope of graph when  $I = 0.226$  A

vii) What physical quantity is represented by the slope of the graph at any given point?

viii) Use your graph to describe how the physical quantity in (vii) above is affected as current increases. Explain

b) Using some of the apparatus in **a (i)** above, draw a diagram you would use to determine resistance of the wire. Explain how you determine the resistance

## RACHUONYO SOUTH DISTRICT

1. You are provided with the following:-


- Ammeter
- A voltmeter
- A straight wire XY mounted on a millimeter scale
- Two jockeys
- 7 connecting wires
- A micrometer screw gauge (to be shared)
- A cell holder for two dry cells
- Two dry cells
- A switch

**Proceed as follows:**

(a) Using the micrometer screw gauge, determine the diameter '*d*' of the wire XY

$d = \underline{\hspace{2cm}}$  mm

Set-up the apparatus as shown below:-



With both jockeys set at  $L = 10\text{cm}$  from X, measure current  $I$  through the wire and voltage

$V$  across it. Repeat this procedure for the other values of  $L$  and record in the table below:

|                                |    |    |    |    |    |    |     |  |
|--------------------------------|----|----|----|----|----|----|-----|--|
| Length (cm)                    | 10 | 30 | 40 | 50 | 70 | 80 | 100 |  |
| Length (m)                     |    |    |    |    |    |    |     |  |
| Current $I$ (A)                |    |    |    |    |    |    |     |  |
| Voltage $V$ (V)                |    |    |    |    |    |    |     |  |
| $R = \frac{V}{I}$ ( $\Omega$ ) |    |    |    |    |    |    |     |  |

- (b) (i) Using the values in the table above, plot a graph of  $I$ (A) against  $R$ ( $\Omega$ ) on the grid provided  
 (ii) Determine the gradient of the graph at  $R = 10\Omega$   
 (iii) Given that  $-I = \frac{\pi d^2 R}{4KL}$  where  $L = 60\text{cm}$ , find the value of  $K$

2. You are provided with a glass vessel of a minimum height of 12cm and a minimum diameter of 6cm, a retort stand and clamp, two optical pins labelled **A** and **B**, liquid **M**,  $\frac{1}{2}$ metre rule and a cork:

**Procedure**

Place a pin **A** at the bottom of the glass vessel and then pour liquid **M** upto a height of 2cm from the bottom of the glass vessel. Move pin **B** on a sliding cork adjacent to the jar up or down until there is no parallax between it and the image of pin **A**. mark on the outside of the beaker where the image pin is located


- (a) Measure **X** and record its value  $X = \underline{\hspace{2cm}}$  cm  
mk)

(Note that **X** is the distance between top level of liquid **M** and where the image of pin is located)

- (b) Repeat this procedure for different values of **y** from the initial value of 2cm in steps of 2cm. complete the table shown below:

|      |   |   |   |   |    |    |
|------|---|---|---|---|----|----|
| Y cm | 2 | 4 | 6 | 8 | 10 | 12 |
| X cm |   |   |   |   |    |    |

Arrangement of apparatus




- (c) Draw a graph of **x** (y-axis) versus **y** on the grid provided  
 (d) Calculate the slope, **S** of your graph  
 (e) Given that  $n = \frac{1}{s}$ , calculate the value of **n**

**B. You are provided with the following:**

- A half metre rule
- Two pieces of thread
- 50g mass
- A retort stand, clamp and boss

**Proceed follows**

- (i) Using a loop of thread suspend the half metre rule on the retort stand as shown below, at 41cm  
 (ii) Using the second loop suspend the 50g mass at a point that sets the half metre rule in equilibrium, as shown below. (Horizontally balanced)



Record the value of

P = .....cm


Using the equation;  $M \times q = P \times 50$ , calculate the value of M

N/B  $q + 25 = 41$

### SOTIK DISTRICT 1ST EXAM

1. *You are provided with the following apparatus:-*

- Full retort stand
- Optical pin
- 20cm copper wire
- Protractor
- Two pieces of plasticine
- Cork
- Stop watch



- A. Bend a wire in the middle. Place it on a protractor and adjust an angle  $\theta$  of  $50^\circ$ . Attach the two pieces of plasticine on the ends of the wire. Then place the wire on the optical pin as shown
- B. Displace the wire horizontally by a small angle and set it in oscillation. By use of stop watch obtain time for 20 oscillations. Repeat this procedure for values of  $\theta$ ,  $60^\circ$ ,  $70^\circ$ ,  $80^\circ$ ,  $90^\circ$  and  $100^\circ$ . Record your measurements in the table and complete it determining values of period T, frequency  $f(\text{Hz})$ ,  $f^2 (\text{Hz})^2$  and  $\cos(\theta/2)$

| Angle $\theta^\circ$ | Time t for 20 oscillations in sec | Period T (Sec) | Frequency f (Hz) | $f^2 (\text{Hz})^2$ | $\cos(\theta/2)$ |
|----------------------|-----------------------------------|----------------|------------------|---------------------|------------------|
| 50                   |                                   |                |                  |                     |                  |
| 60                   |                                   |                |                  |                     |                  |
| 70                   |                                   |                |                  |                     |                  |
| 80                   |                                   |                |                  |                     |                  |
| 90                   |                                   |                |                  |                     |                  |
| 100                  |                                   |                |                  |                     |                  |

C. (i) Draw graph of  $f^2 (\text{Hz})^2$  against  $\cos(\theta/2)$

C. (ii) Determine the gradient of the graph

C (iii) The relationship between  $f$  and  $\theta$  is  $4\pi L f^2 = 150Z \cos(\theta/2) - 10$ . Where  $L$  is 0.2m.


Determine the constant **Z** by the graph

**Question 2**

**PART A**

You are provided with;

- 1000ml measuring cylinder , Clean water, Two optical pins, Retort stand, Cork, and a Wooden block



- A. Place optical pin in the measuring cylinder. Add water to reach 400cm<sup>3</sup> mark. View the optical pin through water and locate its image.
- B. (i) Place another optical pin in cork and then in retort stand clamp as shown. Move this optical pin (search pin) up and down until it coincides with image optical pin as seen through water. Read off the volume reading of the search optical pin when it coincides with image of object optical pin. Repeat this procedure for water volume marks of 600, 700,800, 900, and 1000cm<sup>3</sup>. Enter your observations in table \*

**Table**

| Water level volume reading <b>H cm<sup>3</sup></b> | Image level volume reading <b>h cm<sup>3</sup></b> |
|--|--|
| 400  |  |
| 600  |  |
| 700  |  |
| 800  |  |
| 900  |  |
| 1000   |  |

B (ii) Draw a graph of  $H(\text{cm}^3)$  against  $h(\text{cm}^3)$


C (i) Determine slope of your graph

(ii) Determine **M** if  $H = mh + C$  where **M** and **C** are constants

(iii) If  $m = \frac{M}{M-1}$  where **M** is refractive index of water, determine **M**

### Part B

You are provided with a lens, plane mirror, retort stand, cork and optical pin and half metre rule. Set up apparatus as shown



Move the object optical pin up and down. Look for its image. When object optical pin coincides with its image measure the vertical distance from mirror to object optical pin. Repeat the experiment.

- (i) 1<sup>st</sup> attempt: Vertical height of optical pin = \_\_\_\_\_ cm  
2<sup>nd</sup> attempt: Vertical height of optical thumb pin = \_\_\_\_\_ cm
- (ii) Determine the average vertical height of optical thumb pin
- (iii) Now determine the focal length of the lens using the above measurements

## SOTIK DISTRICT 2ND EXAM

### QUESTION 1


#### Part A

You are provided with the following:

- A uniform metre rule.
- A spring balance.
- Two 15 cm long string.
- 2 complete stands.
- 30 cm or half metre rule.
- A set square.

Proceed as follows:

- a) Set the apparatus has been set ready for use as shown in the figure below. The metre rule has suspended at the 99.0 cm mark with a length of string securely tied to the clamp of the retort stand Q. So not change this position throughout the experiment.




- b) Adjust the position of the clamp of the retort stand P so that the metre rule is suspended at 0.5 cm mark and is horizontal.
- c) Note and record the distance,  $x$  from A to B and also the tension,  $T$ , of the spring balance. Enter your results in the table below.

| $x$ (cm) | $T$ (N) | $\frac{1}{x} (m^{-1})$ |
|----------|---------|------------------------|
|          |         |                        |

(3 mks)

- d) Adjust the position of the clamp of the retort stand O so that the metre rule is suspended at the 10.0 cm mark and is horizontal. Note and record the distance  $x$  and  $T$  in the above table.
- e) Repeat part (d) of the experiment with the spring balance suspended at the 15, 20, 25 and 30 cm marks. Enter your results in the table and complete the table.
- f) (i) On the grid provided, plot a graph of  $T$  ( $y$ -axis) against  $\frac{1}{x}$  (5 mks)
- (ii) Determine the slope,  $S$ , of the graph. (2 mks)
- (iii) Given that  $M = \frac{S}{4.8}$ , find a value  $M$ , the mass of the metre rule (1 mk)

- B. a) (i) Place a candle and screen about 50 cm apart. Place a lens (convex) in between the screen and the candle. Move the lens from about 10 cm from the candle towards the screen until a sharp image is focused on the screen. Mark this point  $U_1$ . Move the lens until the second sharp image of a smaller size is focused on the screen. Mark this point  $U_2$ .



(ii) Measure the displacement of the lens i.e. distance between  $U_1$  and  $U_2$  and let it be  $X_1$ . Let the distance between the screen and the candle be  $Y_1$  i.e.  $Y = 50$  cm.

(iii) Repeat the procedure in (a) above by using a value of  $Y$  of 40 cm. Let it be  $Y_2$ . Find the displacement  $X_2$

Find the value of  $x = \frac{x_1 + x_2}{2}$ , and value of  $Y = Y_1 + Y_2$ . (2 mks)


(iv) Given the equation  $4f = \frac{X^2 - Y^2}{Y}$

Calculate the focal length of the lens(f). (3 mks)

2. You are provided with the following apparatus. Two cells of 1.5v each, Nichrome wire gauge 30. An ammeter 0-5 A or 0-25 A range, cell holder, voltmeter 0.3v or 0.5 eight conductors at least 4 with crocodile clips,, A switch and a metre rule.

**Procedure**

(a) Connect the circuit as shown in the diagram below.



(b) (i) Connect the ends A and point C where AC is 100 cm across the terminal as shown. Close the switch and measure both current I and p.d. across the wire AC.

Current I = (1 mk)

p.d.(v) = (1 mk)

(ii) Measure the e.m.f. of the cells E = (1 mk)

(c) Reduce the length AC as shown 100, 70 cm, 60 cm, 50 cm, 40 cm and 20 cm. In each case record the current (I) and the corresponding p.d.(v)

(d) Enter the length in table.

| Length L (cm) | 100 | 70 | 60 | 50 | 40 | 20 |
|---------------|-----|----|----|----|----|----|
| I (A)         |     |    |    |    |    |    |
| V (Volts)     |     |    |    |    |    |    |
| E - V (V)     |     |    |    |    |    |    |

Complete the table (6 mks)

(e) (i) Plot a graph of (E-V) (V) on the y-axis against I(A) on the x-axis. (5 mks)

(ii) Determine the gradient of the graph. (3 mks)

(iii) Given the equation  $E + V + Ir$  determine the internal resistance of each cell. (3 mks)



**TRANS MARA DISTRICT**


1. **You are provided with the following:-**

\*

- ◆ Dry cell,
- ◆ An ammeter (0.25A)
- ◆ A voltmeter (0-2.5V)
- ◆ A mounted resistance wire,
- ◆ 6 connecting wires
- ◆ A jockey or a crocodile clip

**Proceed as follows:-**

(a) Set up the experiment as shown in the figure below:-



- (b) With the jockey at X, record the ammeter and voltmeter reading  
(c) Vary the length of the mounted wire through which current flows by moving away from X to a new point Y. record the ammeter and voltmeter readings obtained in the table below.

| Length (cm)           | 0.0 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 | 0.8 |
|-----------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Volmetre readings (v) |     |     |     |     |     |     |     |     |     |
| Ammeter readings (A)  |     |     |     |     |     |     |     |     |     |


- (d) On the grid provided plot a graph of V (y axis) against I  
(e) Determine the slope, **M** of the graph  
(f) The equation of the cell from which current is being drawn is  $E = V + Ir$   
Use the equation to determine:-  
(i) E.m.f of the cell  
(ii) The internal resistance of the cell

2. **You are provided with the following:-**

- ◆ A candle
- ◆ a metre rule
- ◆ A white screen
- ◆ A lens and a lens holder
- ◆ Cross wires mounted on a cardboard
- ◆ Plasticine

**Procedure:-**

- (a) Place a metre rule on a bench and hold it in position using plasticine  
Arrange the screen, the lens and the cross wires along the metre rule as shown:



- (b) Light the candle and place it next to the cross wires such that the flame is at the same level with cross wires and the cross wires coincide with the middle part of the flame.
- (c) Adjust the position of the lens so that  $U=15\text{cm}$ . Now adjust the position of screen until a sharply focused image of cross wire is obtained in the screen. Record the value of  $V$  in the table
- (d) Repeat the procedure in (c) above for values of  $U$  shown in the table

| U(cm) | V(cm) | (U + V)cm |
|-------|-------|-----------|
| 15    |       |           |
| 17    |       |           |
| 19    |       |           |
| 21    |       |           |
| 23    |       |           |
| 25    |       |           |
| 27    |       |           |
| 29    |       |           |

- (e) Plot a graph of  $(U + V)$  against  $V$
- (f) From the graph determine the values of  $V$  and  $(U+V)$  for which the graph has minimum values  
 $V$  minimum  $V_m = \underline{\hspace{2cm}}$  cm  
 $(U+V)$  minimum  $(U + V)_m = \underline{\hspace{2cm}}$  cm
- (g) Calculate the values of  $h_1$  and  $h_2$  from the equations below  
 (i)  $h_1 = \frac{V_m}{2} = \underline{\hspace{2cm}}$  cm  
 $h_2 = \frac{(U+V)_m}{4} = \text{cm } \underline{\hspace{2cm}}$
- (ii) Determine the average of  $h_1$  and  $h_2$   
 $h = \frac{h_1 + h_2}{2} = \underline{\hspace{2cm}}$  cm
- (h) Using the graph, determine  $V$  when  $U + V = 41.6\text{cm}$

### **SECTION III – ANSWERS**

#### **KAKAMEGA CENTRAL DISTRICT**

1. c)i) Repeat the procedure in (b) above for the values of  $d$  shown in the table (set the values of  $d$  by adjusting the positions of the loops in steps of 5cm on both sides)

ii) Complete the table

| D (cm) | D (m) | $\frac{L}{d^2}$ ( $M^{-2}$ ) | Time for 10 oscillations<br>$\pm 0.50\text{ s}$ | Period T (s) | $T^2$ ( $S^2$ ) |
|--------|-------|------------------------------|---|--------------|-----------------|
| 80     | 0.80  | 1.5625                       | 5.91  | 0.591        | 0.3493          |
| 70     | 0.70  | 2.04082                      | 7.66  | 0.766        | 0.5868          |
| 60     | 0.60  | 2.778                        | 8.65  | 0.865        | 0.7482          |
| 50     | 0.50  | 4.000                        | 10.44   | 1.044        | 1.0899          |

|    |      |         |       |       |        |
|----|------|---------|-------|-------|--------|
| 40 | 0.40 | 6.2500  | 12.88 | 1.288 | 1.6589 |
| 30 | 0.30 | 11.1111 | 16.94 | 1.694 | 2.8696 |
| 20 | 0.20 | 25.0000 | 25.41 | 2.541 | 6.4567 |

d) i) On the grid provided, plot a graph of  $T^2$  (y-axis) against  $1/d^2$  ( $M^{-2}$ )

ii) Determine the slope of your graph

$$\text{Slope} = \frac{(250 - 0) \times 10^{-2} S^2}{(100 - 0) \times 10^{-1} M^{-2}} \quad \left| \quad \begin{aligned} S &= 2.5 \times 10^{-1} M^2 S^2 \\ S &= 0.25 M^2 S^2 \end{aligned}$$

iii) Given that  $T^2 = \frac{16K^2}{5d^2}$  where  $K$  is a constant. Use the graph to determine the value of

$$\frac{16K^2}{5} = \text{slope} \quad \left| \quad \begin{aligned} K &= \sqrt{\frac{0.25 \times 5 M^2 S^2}{16}} \\ K &= 0.2795 \end{aligned}$$

$$K^2 = \frac{S \times 5}{16}$$

### QUESTION 2

c) Measure the distance,  $d$ , between the two parallel wire that acts as the object  $d = 1.50$  cm  
Record this value in the table below.

g) Repeat your readings of  $x$  with  $u = 70, 60, 50, 40$  and  $30$  cm and complete the table (5 marks)

|          |     |       |       |       |       |       |
|----------|-----|-------|-------|-------|-------|-------|
| $U$ (cm) | 80  | 70    | 60    | 50    | 40    | 30    |
| $X$ (cm) | 0.5 | 0.7   | 0.9   | 1.2   | 1.8   | 3.2   |
| $d/x$    | 3   | 2.143 | 1.667 | 1.250 | 0.833 | 0.469 |

i) On the grid provided plot a graph of  $u$  (y-axis) against  $d/x$

ii) I. Determine the slope,  $S$  of the graph

$$S = \frac{85 - 40}{275 - 80} \times 10^{-2} \quad \left| \quad \begin{aligned} S &= \frac{45}{195} \times 10^{-2} \\ &= 23.076 \text{ cm} \end{aligned}$$

II. Find the intercept on the  $u$ -axis

$U$ -intercept = 20.0 cm

### PART B

iii) Repeat procedure (i) and (ii) above with  $XJ = 20$  cm and enter in the table 3as below

Table 3

| Length $XJ$ (cm) | P.d.V. (v) | Current, $I$ (A) |
|------------------|------------|------------------|
| 10               | 1.1        | 0.45             |
| 20               | 1.5        | 0.25             |

J) Given that  $\log I = n \log V + \log K$ , where  $k$  and  $n$  are constants, determine the values of  $k$  and  $n$

$$\begin{aligned} \log(0.45) &= n \log(1.1) + \log k & -0.03468 &= 0.0414n \\ \log(0.25) &= n \log(1.5) + \log k & -0.6021 &= 0.1761n \\ & & 0.2553 &= -0.13469n \\ & & n &= 1.8955 \end{aligned}$$

$$\left. \begin{aligned} T. 6532 \\ -T. 3979 \\ 0.2553 \end{aligned} \right\} = \left. \begin{aligned} 0.414 \\ 0.1761 \\ T. 8653 \end{aligned} \right\} n$$

$$\left. \begin{aligned} n &= 0.2553 \\ -0.1347 \\ &= -1.8955 \end{aligned} \right|$$

$$\log(0.45) = -1.8955 \log(1.1) + \log k$$

$$\log k = \log(0.45) + 1.8955 \log(1.1)$$

$$= T. 6532 + 1.8955 \times 0.0414$$


$$= T. 6532$$

+ 0.0785      log = T.7317  
T.7317  
K = 0.53913.

**KAKAMEGA EAST DISTRICT**

| $X_{cm}$ | $X_1 C(cm)$                               | $X_2 (cm)$                                 | $X_1 - X_2 (cm)$   |
|----------|---|--|--|
| 45       | 22.5                                      | 19.25                                      | 3.25   |
| 40       | 20.0                                      | 17.75                                      | 2.25   |
| 35       | 17.5                                      | 15.35                                      | 2.20   |
| 30       | 15.0                                      | 13.10                                      | 2.00   |
| 25       | 12.5                                      | 11.00                                      | 1.25   |
| 20       | 10.5                                      | 9.50                                       | 1.00   |
|          | ½ mk each<br>Max mk – 3<br>At least 1 d.p | ½ each 1.0<br>Max mk -3<br>At least - 1d.p | Correct subtraction from<br>candidates' results<br>5-6 correct -1mk 3-4 correct – ½mk<br><3correct - 0 |

- (c) (i) Graph : Labelled axes with units..... 1mk  
: Simple and uniform scale ..... 1mk  
: Plotting (4 correctly plotted pts ..... ½mk max 2mk  
: Line (through any 3 of 4 correctly plotted)... 1mk  
.....Total 5mks)



**question 2**

(a) (ii)


|               |      |      |      |      |      |      |  |
|---------------|------|------|------|------|------|------|--|
| L(cm)         | 100  | 80   | 60   | 40   | 20   | 0    |  |
| P.d, V(volts) | 0.13 | 0.15 | 0.20 | 0.25 | 0.45 | 0.70 | ±0.02<br>½ mk each max 3mks atleast 1d.p |
| Current, I(A) | 0.07 | 0.10 | 0.11 | 0.13 | 0.15 | 0.17 | ±0.02<br>½ mk each max 3mks atleast 1d.p |

- (b) (i) Axes – labelled with units ... 1mk  
Scale – simple and uniform .... 1mk  
Plotting – 5-6 correctly plotted .....2mks

3-4 – correctly plotted ..1mk  
 < 3 correctly plotted .....0mk  
 total 5mks

(ii) Gradient  $S = \frac{\Delta V}{\Delta I}$   
 $\frac{(52 - 25) \times 10^{-2} V}{(16 - 14)}$   
 $= 13.0 \Omega \pm 2$

(iii) Resistance.....1mk



(ii) Gradient  $S$   
 (c) (ii)  $I = 0.09 \text{ A} \pm 0.01$  ..... ½ mk  
 $V = 0.80 \text{ V} \pm 0.01$  ½ mk

(iii)  $d = 0.4 \text{ mm} \pm 0.02$  ..... 1mk  
 (iv)  $S = \frac{3.142 \times 0.80 \times (4.4 \times 10^{-1})^2}{4 \times 0.09 \times 100}$

Answer either in  $\Omega\text{m}$  or  $\Omega\text{cm}$

**MIGORI/NYATIKE DISTRICT**

1. (a) (i)  $h = 5.0 \text{ cm}$   
 (ii)  $R = \sqrt{\frac{100}{h}}$  ✓1  
 $= \sqrt{\frac{100}{5}}$  ✓1  
 $= 4.47 \text{ cm}$  ✓1

(b) (iv)

| U     | 3R    | 4R    | 5R    | 6R    | 7R    | 8R    | 9R   | 10R   |
|-------|-------|-------|-------|-------|-------|-------|------|-------|
| U(cm) | 13.40 | 17.90 | 22.40 | 26.80 | 31.30 | 35.80 | 40.2 | 44.70 |
| V(cm) | 21.0  | 15.0  | 14.0  | 12.5  | 12.0  | 11.5  | 11.0 | 10.5  |

Consistency in values ✓1 ✓1 ✓1 ✓1

(c) graph plotting  
 3 and 4 points } P2  
 5-8 points }  
 scale S1  
 curve C1  
 axis A1  
 ✓2

(d) (i)  $V^l = 17.5 \text{ cm}$   
 (ii)  $U^l = 2.26.25$  ✓2  
 Give reading off from the graph \*UGU\* each

(e)  $f = \frac{U^l + V^l}{5}$   
 $= \frac{17.5 + 27.5}{5}$  ✓1  
 $= \frac{45}{5} = 9 \text{ cm}$  ✓1

$$f) \frac{R}{f} = \frac{4.47}{9} \quad \checkmark 1$$

$$= 0.496667$$

$$= 0.50 \quad \checkmark 1$$

(total 20mks)

2. Part A

(d)

|                  |      |      |      |      |      |      |      |      |      |
|------------------|------|------|------|------|------|------|------|------|------|
| Time t (min)     | 0    | 1    | 2    | 3    | 4    | 5    | 6    | 7    | 8    |
| Temperature (°C) | 80.0 | 77.0 | 74.0 | 71.0 | 68.5 | 66.5 | 64.0 | 62.0 | 60.5 |

(i) trend  $\checkmark 1$   
 decimal place acquired  $\checkmark 1$   
 filling the table  $\checkmark 2$

(ii) Plotting P2  $\checkmark 1$

scale S  $\checkmark 1$

Axis A  $\checkmark 1$

curve C  $\checkmark 1$

(iii) Tangent line drawn at 70°C and identifying two points

(5,66 and (1,45, 74)

$$\frac{d\theta}{dt} = \frac{74 - 66}{1.45 - 5} \quad \checkmark 1$$

$$= \frac{8}{-3.15} \quad \checkmark 1$$

$$= -2.540 \quad \checkmark 1$$

(iv) Heat loss,  $R = K \frac{d\theta}{dt}$

$$= 1.23 \times 10^4 \times 2.540 \quad \checkmark 1 = 31,242J$$

### SOTIK DISTRICT

1. (i)  $d = 0.635cm \ 0.05$

(ii)  $V = \frac{4}{3} (0.635)^3 = 0.134cm^3 \ 0.13 \text{ allow } 0.14$

(iii)  $d = 1.68 \ 0.01$

(iv)  $A = r^2 = 3.14 \times (1.68)^2 = 2.21 \ (2.21 \ - \ 2.30)$

=

(b) (iii)  $h_o = 8.6cm \pm 0.2cm \ 8.0$

(d)

| N | h    | h-h <sub>o</sub> |
|---|------|------------------|
| 1 | 9.0  | 0.4              |
| 2 | 9.7  | 1.1              |
| 3 | 10.3 | 1.7              |
| 4 | 10.9 | 2.3              |
| 5 | 11.5 | 2.9              |
| 6 | 12.0 | 3.4              |

(1mk for mark for each h value upto a max. of 4mks )

1mk for atleast 5 correct differences

½ mk for 3 and 4 correct differences

2 or less correct differences 0mks

(e) Axes – must be labeled the units (mark both)


Scale – be simple & uniform \*

Plotting = ½ mk for each correctly plotted points up to 2mks

> 4 correct plotted 2mks

> 2 & 3 correctly plotted 1mk

<2 correctly plotted 0mk



Line must pass at least 3 correctly plotted points

(f)  $S = 0.527\text{cm}$  – Identifying the pts on the graph

-Correct substitution ( $\frac{1}{2}$  mk)

-Correct answer to 2d.p ( $\frac{1}{2}$  mk)

(g)  $O_s = \frac{0.527 \times 2.21}{0.134}$  (correct sub)

$0.134$

$= 8.69$  (no units) (correct evaluation to 2d.p –

2. PART I

(c)


|     |            |            |            |            |            |                        |
|-----|------------|------------|------------|------------|------------|------------------------|
| $i$ | $10^\circ$ | $20^\circ$ | $30^\circ$ | $40^\circ$ | $50^\circ$ | $60^\circ$             |
| $r$ | $8^\circ$  | $13^\circ$ | $18^\circ$ | $24^\circ$ | $30^\circ$ | $36^\circ \pm 1^\circ$ |
| $d$ | $0.5$      | $0.9$      | $1.3$      | $1.9$      | $2.8$      | $3.2 \pm 0.1\text{cm}$ |

$\frac{1}{2}$  mk for each correct value of both  $r$  and  $d$

(d)

- Axes (Quantity and units- 1mk)
- Scale (simple and uniform 1mk)
- Plotting ( $\frac{1}{2}$  mk for each correctly plotted point up to 2mks)
- The graph's curve as shown
- Maximum = 6mks
- Smooth curve with correct shape as shown 1mk

Note:- The workings on the plain paper must be seen before marking this section



2. Part II

(b)  $t_1 = 30\text{cm}$ ,  $V_1 = 0.75 \pm 0.05$

$t_2 = 50\text{cm}$ ,  $V_2 = 1.225$

$t_3 = 70\text{cm}$ ,  $V_3 = 1.75$

(d)  $t_1 = 30\text{cm}$ ,  $I = 0.12\text{A} \pm 0.01$

$t_2 = 50\text{cm}$   $t_2 = 0.16\text{A}$

$t_3 = 70\text{cm}$   $F_3 = 0.20\text{A} * \text{STK}^*$

e(i)  $V_1 = 30 \times 0.0025 = 0.75\text{V}$

$V_2 = 50 \times 0.025 = 1.25\text{V}$

$V_3 = 70 \times 0.025 = 1.75$

3 values of  $V$  correctly calculated 1mk

2 values of  $V$  correctly calculated  $\frac{1}{2}$  mk

0.01 value of  $V$  correctly calculated 0mk

(ii)  $R_1 = \frac{V_1}{I_1} = \frac{0.75}{0.12} = 6.25\Omega$

$R_2 = \frac{V_2}{I_2} = \frac{1.25}{0.16} = 7.81\Omega$

$R_3 = \frac{V_3}{I_3} = \frac{1.75}{0.2} = 8.75\Omega$

3 values correctly calculated to 2d.p 1mk

2 values correctly calculated to 2d.p  $\frac{1}{2}$  mk

0 or 1 value correctly calculated to 2d.p 1mk 0mk

Average  $R = \frac{R_1 + R_2 + R_3}{3} = \frac{6.25 + 7.85 + 8.75}{3} = \frac{22.85}{3} = 7.617\Omega$

Substitution of 3 values of  $R$   $\frac{1}{2}$  mk

Correct evaluation of average  $R$  value to 3d.p  $\frac{1}{2}$  mk

**UGENYA/UGUNJA**

1. (d)

(a) Repeat the procedure above for the other given angles on table 1

| Angle $\theta^\circ$ | Time (t) for 10 oscillations | $f = 10/t$ | $f^2$<br>(Hz) <sup>2</sup> | $\theta/2$ | Cos ( $\theta/2$ ) |
|----------------------|------------------------------|------------|----------------------------|------------|--------------------|
| 20                   | 6.28                         | 1.592      | 2.54                       | 10         | 0.99               |
| 40                   | 6.47                         | 1.546      | 2.39                       | 20         | 0.94               |
| 60                   | 6.63                         | 1.508      | 2.28                       | 60         | 0.87               |
| 80                   | 7.28                         | 1.374      | 1.89                       | 40         | 0.77               |
| 100                  | 7.87                         | 1.271      | 1.62                       | 50         | 0.64               |

(f) Gradient of  $\Delta y = (2.4475 - 1.75) \text{ Hz}^2$

$$\begin{aligned} \Delta x &= (0.96 - 0.68) \\ &= 0.28 \\ &= 2.589 \text{ Hz}^2 \end{aligned}$$

(c) (i)  $f = \frac{10}{t} = 10$   
 $t = 6.28$   
 $= 1.592 \text{ Hz}$

**PART B**

(c)

| Angle of incidence | LN (mm) | MN (mm) | Refractive index $LN/MN$ |
|--------------------|---------|---------|--------------------------|
| 15°                | 13      | 7       | 1.857                    |
| 30°                | 25      | 17.5    | 1.429                    |
| 45°                | 36      | 24      | 1.500                    |

(d) Average =  $\frac{1.875 + 1.429 + 1.500}{3}$

$$= 1.60 \text{ (2 dp)}$$

Results of question 2

| a cm | b cm | c cm | d = a/c |
|------|------|------|---------|
| 94   | 67   | 27   | 3.481   |
| 90   | 62   | 28   | 3.214   |
| 86   | 56   | 30   | 2.867   |
| 82   | 50   | 32   | 2.563   |
| 78   | 42   | 36   | 2.167   |
| 74   | 37   | 37   | 2.00    |

Trend 2  $d=1$   
 Correct = 3  $4 \text{ correct} = 2$

total 06mks

(d) graph  $A_2 S_1 P_3 L_1 = 7 \text{ mks}$

(e) Slope  $\frac{\Delta b}{\Delta d} = \frac{2.5 \text{ cm}}{1.3}$

$$S = 19.23 \text{ cm}$$

(f)  $L = \frac{100}{5}$

$$= \frac{100}{5} = 19.23 = 5.2 \text{ Dioptres } (5.0 \pm 0.2)$$

(g) Power of the lens

**NDHIWA DISTRICT**

1.

(a) (iii)  $V = 45 \text{ cm}^3$

(iv)  $I = \frac{106 \times 0.30}{45}$  for substitution

$$= 6,666.7 \text{ kg/m}^3 \quad \checkmark$$



(d)

|                                 |       |       |       |        |       |       |
|---------------------------------|-------|-------|-------|--------|-------|-------|
| Length L(cm)                    | 70    | 60    | 50    | 40     | 30    | 20    |
| Length L (m)                    | 0.70  | 0.60  | 0.50  | 0.40   | 0.30  | 0.20  |
| Time for 20 oscillations (s)    | 61.01 | 56.08 | 51.36 | 46.006 | 39.81 | 32.80 |
| Period T (s)                    | 3.057 | 2.804 | 2.568 | 2.303  | 1.991 | 1.640 |
| T <sup>2</sup> (s) <sup>2</sup> | 9.309 | 7.862 | 6.595 | 5.304  | 3.960 | 2.690 |

- Correct conversion of length L in m

- time for 20 oscillations ✓ (½mk) for each value within range (maximum of 3mks)

- correct evaluation of period ✓

- correct evaluation of T<sup>2</sup> ✓

(e)

Axes : labelled with units

Scale : simple and uniform ✓

Plotting : Each correctly plotted (½mk) to a max of 2mks

line – passing through at least 3 points correctly plotted. ✓

(f)  $d = 0.34 \times 10^{-3}$

$= 3.4 \times 10^{-4}m$

(g) (i)  $\frac{\Delta T^2}{\Delta L} = \frac{8.6 - 2.0}{65 - 15}$

$\frac{6.6}{50} = 0.1320$

(ii)  $T^2 = \frac{32\pi^2 L}{Gd}$  ✓

Gradient =  $\frac{32\pi^2}{Gd}$

$G = \frac{322 \times (3.14)^2}{3.4 \times 10^{-4} \times 0.132}$   
 $= 7.03898 \times 10^7$

Q2. (a) (i)  $V_1 = 3V$

(ii)  $V_2 = 2.80V$

$I = 0.24A$

(ii)  $P = \frac{3.0 - 2.8}{0.2} = 1\Omega$

(b) (i)  $V = 2.5V$

$I = 0.25A$  ✓

$R = \frac{2.5}{0.25} = 10\Omega$

- Correct conversion ½

- Voltmeter reading ½ each

|                                  |       |        |        |       |       |          |
|----------------------------------|-------|--------|--------|-------|-------|----------|
| Length L (cm)                    | 100   | 80     | 60     | 40    | 20    | 10       |
| $\frac{I}{L}$ ( $\frac{A}{cm}$ ) | 0.01  | 0.0125 | 0.0167 | 0.025 | 0.05  | $\infty$ |
| Voltmeter reading (v)            | 1.8   | 1.3    | 1.1    | 0.9   | 0.6   | 0.0      |
| $\frac{I}{V}$ ( $\frac{A}{V}$ )  | 0.667 | 0.769  | 0.909  | 1.111 | 1.667 | $\infty$ |

**MUMIAS DISTRICT**

a) diameter  $d = 0.28mm + 0.02$

|               |      |      |      |      |      |      |       |       |       |       |
|---------------|------|------|------|------|------|------|-------|-------|-------|-------|
| Length (cm)   | 10   | 20   | 30   | 40   | 50   | 60   | 70    | 80    | 90    | 100   |
| Length (m)    | 0.1  | 0.2  | 0.3  | 0.4  | 0.5  | 0.6  | 0.7   | 0.8   | 0.9   | 1.0   |
| Current I (A) | 0.70 | 0.54 | 0.44 | 0.32 | 0.28 | 0.26 | 0.22  | 0.20  | 0.18  | 0.16  |
| Voltage V (V) | 1.20 | 1.80 | 1.80 | 2.20 | 2.40 | 2.50 | 2.50  | 2.60  | 2.70  | 2.74  |
| $R = v/I$ (x) | 1.71 | 3.33 | 4.09 | 6.88 | 8.57 | 9.62 | 11.36 | 11.36 | 13.00 | 17.13 |

b) GRAPH

i) Axes- labeled with correct units

Scale – simple, uniform, and consistent and accommodate all values

Points – 4 correctly plotted points within an error of one small square

Line- straight line passing through at least 3 correctly plotted points

ii) Gradient =  $\frac{R}{L} = \frac{16.4 - 0}{1.04} = 15.7692 \text{ nm}^{-1}$

iii)  $R = \frac{4p^2}{d^2}$

$$M = \frac{4p}{d^2}$$

$$P = \frac{md^2}{4}$$

$$= \frac{15.7692 \times (0.28 \times 10^{-4})^2}{4} = 9.7265 \times 10^{-8} \text{ RM}$$

2. a)  $L = 8.0 \text{ cm}$

d) i)  $0.80 + 0.2$

|                    |     |      |      |      |      |
|--------------------|-----|------|------|------|------|
| $H(\text{cm})$     | 1   | 2    | 3    | 4    | 5    |
| $W_2(\text{N})$    | 0.8 | 0.7  | 0.6  | 0.5  | 0.4  |
| $F = (W_1 - W_2)N$ | 0.2 | 0.30 | 0.40 | 0.50 | 0.60 |

f) i) Labelling ✓

Appropriate scale ✓

Plotting ✓ ✓

Straight/ smooth line ✓

$$\begin{aligned} \text{ii) } M &= \frac{DF}{DH} \\ &= \frac{0.6 - 0.2}{5-1} \\ &= \frac{0.4}{4} \\ &= 0.1/\text{cm} \end{aligned}$$

$$\begin{aligned} \text{iii) } D_g/100 &= m \\ D \times 10^3/100 &= 0.1/0.01 \\ &= 100 \text{ Kg/m} \end{aligned}$$

$$\text{iv) } W = 100 \text{ Kg/m} \times 0.075 \text{ m} = 7.5 \text{ Kg}$$

**KISUMU DISTRICT**

Question 1.

(b)  $E_o = 3.2 \pm 0.1 \text{ volts}$

|                                  |                                    |      |      |      |      |             |
|----------------------------------|------------------------------------|------|------|------|------|-------------|
| Voltage (V)<br>Volts             | 0.5                                | 1.0  | 1.5  | 2.0  | 2.5  | 2.7         |
| Current (I) A                    | 0.67                               | 0.48 | 0.36 | 0.24 | 0.12 | 0.06 ± 0.02 |
| $\frac{I}{I}$                    | Candidate's values correct to 3d.p |      |      |      |      |             |
| $\frac{R}{I} = V \text{ (Ohms)}$ | Candidate's values correct to 3d.p |      |      |      |      |             |

✓ each upto  
max. 5

✓ all values  
1mk

(e) graph

(f) Slope =  $\frac{3.0 - 1.50}{5.0 - 0}$

$$= \frac{1.5}{5.0}$$

$$= 0.3 \text{ A}^{-1} \Omega^{-1}$$

(g) (i) When  $\frac{1}{I} = 0$ ,  $r = R_o = 5 \Omega \pm 1$  \*KSM\*

(ii)  $\frac{1}{I} - E_o$

$$s = \frac{1}{0.3} - 3.2$$

$$= 0.1 \pm 0.01 \quad V$$

**Question 2**

(a)  $f = 20 \text{ cm}$

(c) (i) erect (ii) virtual (iii) magnified

|                                 |                                   |     |     |      |      |      |      |                |
|---------------------------------|-----------------------------------|-----|-----|------|------|------|------|----------------|
| Object Distance, $U \text{ cm}$ | 3.0                               | 4.0 | 6.0 | 7.0  | 8.0  | 10.0 | 11.0 | 12.0           |
| Image distance, $V \text{ cm}$  | 3.0                               | 5.0 | 7.8 | 10.0 | 13.3 | 18.5 | 22.0 | $29.1 \pm 0.2$ |
| $M = \frac{V}{U}$               | Candidates correct values to 1d.p |     |     |      |      |      |      |                |

✓ each upto max. 7

(f) graph ...

total – 8mks

(g) (i) when  $u \approx f$ ,  $V = \text{Infinitely large number (undefined)}$

N/B: i.e when object is at  $F$ , the image is formed at infinity

(ii) Application: Magnifying lens

**TRANS NZOIA WEST DISTRICT**

1. a)  $h = 30 \text{ cm}$

b)  $f = \frac{h}{2} = \frac{30}{2} = 15 \text{ cm}$

c) Table of results

| Object distance $U \text{ cm}$ | Image distance $V \text{ (cm)}$ | Magnification $M = \frac{v}{u}$ |
|--------------------------------|---------------------------------|---------------------------------|
| 22                             | 10.9                            | 0.4955                          |
| 24                             | 10.2                            | 0.4250                          |
| 26                             | 10.0                            | 0.3846                          |
| 28                             | 9.6                             | 0.3429                          |
| 30                             | 9.5                             | 0.3167                          |

Accuracy =  $\pm 0.2$

- For  $V$  values award 3 marks for all values within range see accuracy above, otherwise award  $\frac{1}{2}$  mark for each correct
- For  $M$  award 2 marks for all values correct but 3 -4 values correct award 1 mark and less than 3 values correct award 0 marks
- N.B If a school did not use the specified focal length of the mirror it should be indicated so that candidate are not penalized

f) Graph on graph paper

Axes (1 mark) - Both quantity and unit on both

Scale (1 mark) – Simple and uniform

Plotting (2 marks) – All points correctly plotted award 2 marks

3 – 4 points correctly plotted (1 mark)

Line (1 mark) line to pass in at least three correctly plotted points

NB The line MUST cut the negative part of the  $Y$ - axis for a mark

g) Gradient/ slope =  $\frac{1}{f}$

$$f = \frac{1}{\text{slope}}$$

$$= 15 \pm 0.1 \text{ cm}$$

**PART B**

**TABLE 2 RESULTS**

K

i)

| U (cm) | V (cm) | UV (cm <sup>2</sup> ) | U + v cm |
|--------|--------|-----------------------|----------|
| 20     | 20.0   | 400.00                | 40.0     |
| 30     | 15.5   | 465.00                | 45.5     |

$$ii) f = \frac{uv}{u+v} = \frac{400 + 465}{40 + 45.5} = \frac{865}{85.5} = 10.12cm$$

2.

a)

iii)

**Table of results**

|           |      |      |      |      |      |      |
|-----------|------|------|------|------|------|------|
| L (cm)    | 100  | 80   | 60   | 40   | 20   | 0    |
| V (volts) | 1.20 | 1.35 | 1.50 | 1.75 | 2.00 | 2.50 |
|           | 0.18 | 0.20 | 0.22 | 0.23 | 0.24 | 0.28 |

Award ½ mark for each correct reading See accuracy above

iv) It is brighter

v) Graph on graph paper

Scale 1 mark (simple and uniform)

Plotting 2 marks (each ½ mark max 4 points correctly plotted)

Axes 1 mark (Both quantity and unit on both axes)

Curves 1 mark – passing through at least 3 correctly plotted points – No mark for straight line

$$vi) \text{ Slope} = \frac{DV}{DI} = \frac{1.9 - 1.5}{0.25 - 0.22} = 13.33$$

Tangent 1 mark (Tangent line must be continuous)

vii) Electrical resistance of the bulb

viii) - Resistance increases with increase in current

- There is increased resistance to flow of electric current as temperature increases

b) Resistance of wire  $R = \frac{V}{I}$

Award for R as shown above or if values have been substituted by the candidate

**RACHUONYO SOUTH DISTRICT**

1.

a)  $d = 0.3mm \checkmark$

|                            |      |      |      |      |       |      |       |
|----------------------------|------|------|------|------|-------|------|-------|
| Length (m)                 | 0.1  | 0.3  | 0.4  | 0.5  | 0.7   | 0.8  | 1.0   |
| I (A)                      | 0.7  | 0.44 | 0.36 | 0.28 | 0.22  | 0.19 | 0.16  |
| $\frac{V}{I} V(V)$         | 1.2  | 1.80 | 2.10 | 2.40 | 2.5   | 2.6  | 2.74  |
| $\frac{V}{I} = R (\Omega)$ | 1.17 | 4.09 | 5.83 | 8.57 | 11.36 | 13.7 | 17.13 |

b)

i) Scale

$S_1$

Axes

$A_1$

Plotting

$P_2$

Curve

$C_1$

ii) Tangent  $\checkmark$

$$= \frac{0.47 - 0.2}{12 - 0}$$

Co-ordinates  $\checkmark$

$$12 - 0$$

Correct working  $\checkmark$

$$= 0.0225$$

$$iii) -I = \frac{\pi R d^2}{4KL}$$

$$I = \frac{\pi d^2}{4KL}$$

$$R = 4KL \text{ (Slope at } R = 10\Omega) \checkmark$$

$$-0.0225 = \frac{-\pi \times (0.3 \times 10^{-3})^2}{4 \times K \times 0.6} \checkmark$$

$$K = 3.927 \times 10^{-4}$$

$$\Omega m^{-1} A^{-1}$$

2. a)  $x = 1.5\text{cm} \checkmark$

b)

|      |     |   |     |   |     |    |
|------|-----|---|-----|---|-----|----|
| y cm | 2   | 4 | 6   | 8 | 10  | 12 |
| x cm | 1.5 | 3 | 4.5 | 6 | 7.5 | 9  |

c)  $A_1 \checkmark$        $S_1 \checkmark$        $P_2 \checkmark \checkmark$        $L_1 \checkmark$

d)  $S = \frac{6 - 4.5}{8 - 6} \checkmark \checkmark = 0.75 \checkmark$

e)  $n = 1.333 \checkmark \checkmark$

$P = 8\text{cm} \checkmark$

$M \times 16 = 8 \times 50 \checkmark$

$M = 25\text{g} \checkmark$

**SOTIK DISTRICT 1ST EXAM**

**Question 1**


|                                   |       |           |           |           |           |           |  |
|-----------------------------------|-------|-----------|-----------|-----------|-----------|-----------|--|
| Angle $\theta^\circ$              | 50    | 60        | 70        | 80        | 90        | 100       |  |
| Time t sec for 20 oscillations    | 11    | 12        | 13        | 14        | 16        | 18        | T sec, 1mk for each correct value max 5mks |
| Period $T = \frac{t}{20}$ sec     | 0.55  | 0.60      | 0.65      | 0.7       | 0.8       | 0.9       | 1mk for at least 4 correct evaluations     |
| Frequency $f = \frac{1}{T} (H_2)$ | 1.82  | 1.67      | 1.58      | 1.43      | 1.25      | 1.11      |  |
| $f^2 (H_2)^2$                     | 3.312 | 2.78<br>9 | 2.49<br>6 | 2.04<br>5 | 1.56<br>3 | 1.23<br>2 | 1mk for at least 4 correct evaluations     |
| $\cos(\frac{\theta}{2})$          | 0.906 | 0.86<br>7 | 0.81<br>9 | 0.76<br>6 | 0.70<br>7 | 0.64<br>2 | 1mk for at least 4 correct evaluations     |

**N/B**

(i) – Period  $T = \frac{t}{20}$  sec should be correctly evaluated to 2 d.p

- Frequency should be correctly evaluated to 2 d.p
- $f^2 (H_2)^2$  should be correctly evaluated to 3 d.p
- $\cos(\theta/2)$  should be correctly read to 3 d.p

**C(i) Axes 1mk**



Scale (1mk)

Simple and uniform

**Plotting** (2mks)

Plot with accuracy of small lines. Each correctly plotted point award 1/2mk up to max of (2mks)

**Line** (1mk)

+ve slope line with negative y-intersect

**C (ii)** Award gradient mark only when line is correct

Intervals (1mk)  
 Substitution 1mk  
 Evaluation 1mk  
 Gradient =  $\frac{3.31 - 2.05}{0.91 - 0.77} = \frac{1.26}{0.14} = 9$


(iii)  $4\pi L f^2 = 150Z \cos \theta/2 - 10$   
 $f^2 = \frac{150Z \cos \theta/2 - 10}{4\pi L}$   
 Slope =  $\frac{150Z}{4\pi L}$  ✓  
 $= \frac{150Z}{4\pi L} = 9$   
 $Z = \frac{4\pi L \times 9}{150}$   
 $= 4 \frac{22}{7} \times \frac{0.2 \times 9}{150}$   
 $= 0.1509$

N/B:- Pick point from line substitute in equation and solve for Z

**Question 2**

| Water level volume reading Hcm <sup>3</sup> | Image level volume reading hcm <sup>3</sup> |
|---|---|
| 400   | 100   |
| 600   | 150   |
| 700   | 175   |
| 800   | 200   |
| 900   | 225   |
| 1000  | 250   |

B. (ii) Axes ✓ (1mk)



Scale (1mk)

Simple and uniform

Plotting (2mks)

Correctly plotted point (½mk) for max of 2mks

Line ✓ 1mk

Straight line with +ve gradient passing through at least 3 correctly plotted points

C. (i) Without getting mark for line award (0 marks) for slope

Intervals ✓ 1mk

Evaluation ✓ (1mk)

Accuracy ✓ 1mk

$\frac{600-400}{150-100} = \frac{200}{50} = 4$  (no units)

(ii) M = slope = 4 (1mk)

✓

$$\begin{aligned} \text{(iii)} \quad 4 &= \frac{M}{M-1} \\ 4M - 4 &= M \\ 3M &= 4 \\ \Rightarrow M &= \frac{4}{3} \end{aligned}$$

$$M = 1.33 \quad (1\text{mk})$$

**N/B:-** You can also pick two points on the time to solve for M by substituting in equation and solving Simultaneously.

### SOTIK DISTRICT

**Q 1. PART A:**

| $x$ (cm) | $T$ (m) | $\frac{1}{x} (m-1)$ |
|----------|---------|---------------------|
| 94       | 0.8     | 1.06                |
| 89       | 0.7     | 1.12                |
| 84       | 0.6     | 1.19                |
| 79       | 0.5     | 1.27                |
| 75       | 0.4     | 1.33                |
| 70       | 0.3     | 1.43                |

(3 mks)

- f(i) Graph
- Labeling axes. ✓1
  - Appropriate Scale. ✓1
  - Plotting: 5 – 6 pts 2 ✓1 ✓1  
                  : 3 -4 pts ✓1
  - Straight line ✓1

(5 mks)

- (ii) Slope - ✓1 (2 mks)  
Evaluation ✓1  
Answer ✓ (0.48 ± 0.05)

(2 mks)

(ii) Given  $M = \frac{S}{4.8}$  when  $S = 0$

$$M = \frac{0.48}{4.8} = 0.1 \text{ kg } \checkmark 1$$

#### QUESTION 1: PART B

f(i)  $X_1 = 7.5 \text{ cm } \checkmark 1 \text{ mk}$   
 $X_2 = 50 \text{ cm } \checkmark 1 \text{ mk}$   
(ii)  $X_2 = 6 \text{ cm } \checkmark 1$   
 $Y_2 = 40 \text{ cm } \checkmark 1$   
 $X = \frac{7.5 + 6}{2} = 6.75 \text{ cm } \checkmark 1$   
 $Y = \frac{40 + 40}{2}$   
 $= 40 \text{ cm } \checkmark 1$   
(iii)  $4f = \frac{(6.75)^2 - (40)^2}{40} \checkmark 1$   
 $f = 10 \text{ cm } \checkmark 1$   
9

#### QUESTION 2

b) (i)  $I = 0.12A \pm 0.01 \checkmark 1$   
 $V = 2.6V \pm 0.1 \checkmark 1$

(ii)  $E = 3.0\text{v}$  (Max. range) ✓1

d) Table

|          |      |      |      |      |      |      |
|----------|------|------|------|------|------|------|
| $L$ (cm) | 100  | 70   | 60   | 50   | 40   | 20   |
| $I$ (A)  | 0.12 | 0.19 | 0.20 | 0.24 | 0.28 | 0.42 |
| $V$ (v)  | 2.60 | 2.50 | 2.40 | 2.35 | 2.30 | 2.00 |
| $E-V$    | 0.40 | 0.50 | 0.60 | 0.65 | 0.70 | 1.00 |

(7 mks)

e) (i) Graph

- Labeling axis ✓1 mk
- Appropriate scale ✓1 mk
- Plotting: 5 – 6 pts ✓2 mks
- : 3 – 4 pts ✓1 mk
- Straight line ✓1

(ii) Slope - ✓1 mk Evaluation - ✓1 mk

Answer - ✓1 mk

(iii) Slope of the graph =  $r$ , internal /resistance ✓1 3 mks

### TRANS MARA DISTRICT

1. (b)  $I = 1.00\text{Amps}$

$V = 1.00\text{Volts}$

(c)

|                           |      |      |      |      |      |      |      |      |      |
|---------------------------|------|------|------|------|------|------|------|------|------|
| Length $L$ (m)            | 00   | 0.1  | 0.2  | 0.3  | 0.4  | 0.5  | 0.6  | 0.7  | 0.8  |
| Voltmeter reading $V$ (V) | 1.00 | 1.10 | 1.20 | 1.25 | 1.30 | 1.35 | 1.40 | 1.45 | 1.45 |
| Ammeter reading $I$ (A)   | 1.00 | 0.80 | 0.60 | 0.50 | 0.40 | 0.35 | 0.30 | 0.25 | 0.20 |

Increasing  $V$  1mk      Decreasing 1mk

7-9 correct pairs = 2mks

5-6 correct pairs = 1 ½mks

3-4 correct pairs = 1mk

2 correct pairs = ½ mk

Total = 7mks

(d.) Graph

Labelling – both axes quantity and units shown -1mk

Scale-

Plotting – (½mk each maximum) - 2mks

Straight line – through 3 correctly plotted points- 1mk

Total 5mks

(e) Gradient  $\frac{\Delta V}{\Delta I} = \frac{1.3 - 1.2}{0.6 - 0.4} = \frac{0.1}{0.2} = \frac{1}{2} = 0.5$

$$\frac{\Delta V}{\Delta I} = \frac{1.3 - 1.2}{0.6 - 0.4} = \frac{0.1}{0.2} = \frac{1}{2} = 0.5$$

Gradient = 0.5

(f) (i)  $V = Ir + E$

when  $I = 0$ ,  $V = E$ , y intercept

$$E = 1.48 \pm 0.1\text{Volts}$$

(ii) Gradient  $S = -r$

$$r = -S$$

$$= (0.5) = 0.5\Omega$$

2. (d)

| $U$ cm | $V$ cm | $U + V$ cm |
|--------|--------|------------|
| 15     | 30.5   | 45.5       |
| 17     | 24.0   | 41.3       |
| 19     | 21.0   | 40.0       |
| 21     | 19.0   | 40.0       |
| 23     | 17.6   | 40.6       |
| 25     | 16.6   | 41.6       |
| 27     | 15.8   | 42.8       |
| 29     | 15.3   | 44.3       |

(e) On the graph paper – Labelling both axes – 1mk

- scale – 1mk



(f)  $Vm = 20\text{cm}$

$$(U + V)m = 39.9$$

(g) (i)  $h_1 = \frac{Vm}{2} = \frac{20}{2} = 10\text{cm} \pm 0.2\text{cm}$

$$h_2 = \frac{(U + V)m}{4} = \frac{39.9}{4}$$

$$= 9.975\text{cm} \pm 0.2$$

(ii)  $h = \frac{h_1}{2} + \frac{h_2}{2} = 10 + 9.975 = 9.986\text{cm} \pm 0.2$

(h) Value  $V$  when  $U + V = 41.6$

$$V_1 = 16.5\text{cm} \pm 0.2$$

$$V_2 = 24.5\text{cm} \pm 0.2$$